



Development and Characterization of Novel EST-SSR Markers for *Speranskia tuberculata* (Euphorbiaceae)

Authors: Fu, Yi, Ju, Miao-Miao, Ma, Huan-Cheng, Xin, Pei-Yao, He, Cheng-Zhong, et al.

Source: Applications in Plant Sciences, 4(10)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1600067>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

DEVELOPMENT AND CHARACTERIZATION OF NOVEL EST-SSR MARKERS FOR *SPERANSKIA TUBERCULATA* (EUPHORBIACEAE)¹

YI FU², MIAO-MIAO JU², HUAN-CHENG MA², PEI-YAO XIN², CHENG-ZHONG HE²,
DONG-RUI JIA^{3,5}, AND BIN TIAN^{2,4,5}

²Key Laboratory of Biodiversity Conservation in Southwest China, State Forestry Administration, Southwest Forestry University, Kunming 650224, People's Republic of China; ³School of Ecology and Environmental Science, Yunnan University, Kunming 650091, People's Republic of China; and ⁴Key Laboratory of Biodiversity and Biogeography, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, People's Republic of China

- **Premise of the study:** The first set of expressed sequence tag–simple sequence repeat (EST-SSR) markers were developed and characterized for *Speranskia tuberculata* (Euphorbiaceae), a traditional medicinal plant endemic to northern China, to explore the effects of recent habitat fragmentation on the genetic diversity and structure of this species.
- **Methods and Results:** In this study, a total of 18 novel polymorphic microsatellite (EST-SSR) markers were developed for *S. tuberculata* using high-throughput transcriptome sequencing. Analysis of 24 individuals of *S. tuberculata* from four natural populations revealed their robust polymorphic reliability. The number of alleles per locus ranged from two to 11, while the expected and observed heterozygosity per marker varied from 0.187 to 0.827 and 0.042 to 0.917, respectively. Of these markers, 13 showed good amplification results in the closely related species *S. cantonensis*.
- **Conclusions:** These newly generated SSR markers are expected to provide novel tools for genetic studies of *S. tuberculata*, which will contribute to the conservation and sustainable use of the species' wild genetic resources.

Key words: Euphorbiaceae; expressed sequence tag–simple sequence repeat (EST-SSR); *Speranskia cantonensis*; *Speranskia tuberculata*; transcriptome sequencing.

Speranskia Baill. (Euphorbiaceae) is a small genus endemic to China, comprising three herbaceous perennial species: *S. tuberculata* (Bunge) Baill., *S. cantonensis* (Hance) Pax & K. Hoffm., and *S. yunnanensis* S. M. Hwang (Hwang, 1989). *Speranskia tuberculata* is endemic to northern China and occurs on grassy slopes, grasslands, and thickets. The entire plant is commonly used for Chinese traditional medicine (Mazzio et al., 2014). Although *S. tuberculata* is not listed in the IUCN Red List, it is exhibiting a general decreasing trend or even disappearing completely in many distributional areas because of agricultural intensification and over-exploitation of natural population resources. To explore the genetic consequences of recent habitat fragmentation for this medicinal plant and generate useful information to facilitate the conservation and sustainable use of wild genetic resources, we developed the first set of 18 polymorphic expressed sequence tag–simple sequence repeat (EST-SSR) markers for *S. tuberculata* using high-throughput transcriptome sequencing. We also tested these developed markers in *S. cantonensis*, a closely related species (Hwang, 1989), to identify their cross-species utility.

METHODS AND RESULTS

Fresh leaves of *S. tuberculata* seedlings were gathered in Beijing (39°59'06"N, 116°02'04"E; voucher specimen accession no. TB2013079,

¹Manuscript received 3 June 2016; revision accepted 21 July 2016.

This study is supported by the National Natural Science Foundation of China (NSFC; 31260050).

⁵Authors for correspondence: tianbinlzu@gmail.com, jia.drui@ynu.edu.cn

doi:10.3732/apps.1600067

deposited at the Herbarium of Southwest Forestry University [SWFC], Kunming, China) and immediately frozen in liquid nitrogen, and then stored at –80°C. RNA extraction, cDNA library construction, and transcriptome sequencing were conducted following the procedures previously described by Ju et al. (2015). After removing adapter sequences and low-quality sequences, a total of 86,138,489 nonredundant unigenes were assembled from 95,791,418 raw reads. A high-quality reference genome with nonredundant unigenes was then generated by performing de novo transcriptome assembly using Trinity with the parameter of full clean up (Grabherr et al., 2011) and clustering similar contigs using CD-HIT with default parameters (Fu et al., 2012). Furthermore, we used MISA Perl script (MicroSatellite identification tool; Thiel et al., 2003) to screen for SSR motifs from all unigenes, and the minimum numbers of repeats were set as seven, five, five, five, and five for di-, tri-, tetra-, penta-, and hexanucleotide repeats, respectively. MISA recovered a total of 26,202 SSR motifs, of which 30 were randomly selected for primer design using Primer3 software (Rozen and Skaletsky, 1999). The major parameters for primer pair design were set as follows: primer length of 15–25 bases, PCR product size of 100–400 bp, and annealing temperatures of 55–60°C.

The 30 target EST-SSR markers were initially tested for amplification using DNA from 24 *S. tuberculata* individuals from four natural populations located in different provinces across the distributional range in northern China (populations YA, XZ, YT, and KQ; Appendix 1). Total genomic DNA was extracted from silica gel–dried leaves using the Ezup DNA Extraction Kit (Sangon Biotech, Shanghai, China) following the manufacturer's protocol. PCRs were performed using the S1000 Thermal Cycler (Applied Biosystems, Foster City, California, USA) in a 25- μ L total volume with 1 μ L (~10 ng) of genomic DNA, 12.5 μ L of *Taq* PCR Mix (Sangon Biotech), 9.5 μ L of ddH₂O, and 1 μ L (5 pmol) of each primer. The PCR program consisted of 10 min of initial denaturation at 95°C; followed by 35 cycles of denaturation at 94°C for 45 s, annealing at specific temperature (58–60°C; Table 1) for 1 min, extension at 72°C for 1 min; and a final extension at 72°C for 10 min. All PCR products were run on 1% agarose gels to check for successful amplification. Twenty primer pairs produced clear amplicons of the expected size ranges. Multiplex-Ready PCR technology (Hayden et al., 2008) was then applied for fluorescence-based SSR genotyping. Forward primers for the 20 successfully amplified loci were labeled with three different fluorescent dyes (6-FAM, HEX, and NED; Applied Biosystems; Table 1) and used for

Applications in Plant Sciences 2016 4(10): 1600067; <http://www.bioone.org/loi/apps> © 2016 Fu et al. Published by the Botanical Society of America. This work is licensed under a Creative Commons Attribution License (CC-BY-NC-SA).

TABLE 1. Characteristics of the 18 polymorphic microsatellite markers developed for *Speranskia tuberculata*.

Locus	Primer sequences (5'–3')	Repeat motif	Allele size range (bp)	T _a (°C)	Fluorescent dye	GenBank accession no.	BLAST top hit description [organism]	BLAST top hit accession no.	E-value
8852	F: GTGCTCATCCGAAAT R: CAACAGCAGCAAAAACAA	(TGC) ₅	361–370	60	6-FAM	KT285024	Probable methyltransferase PMT27 [<i>Ricinus communis</i>]	XP_002533655.1	2E-10
9441	F: CCAAAAAGCTAAACCACTCG R: CTGCTGCTGTGTTTTG	(GCA) ₅	225–240	59	6-FAM	KT285026	Trihelix transcription factor GTL1 [<i>Ricinus communis</i>]	XP_002516129.1	0.008
9832	F: CTTGCACCTCCAACCTCCG R: AGCTTGAGCATGACGGAGA	(TCA) ₅	192–201	60	6-FAM	KT285027	Uncharacterized protein At1g65710 [<i>Ricinus communis</i>]	XP_002521410.1	0.0000003
10026	F: TGCATTTGATTTGACATTTG R: CACGGTCTCAAAGACC	(TGA) ₆	154–172	59	6-FAM	KT285028	No hit	—	—
10117	F: CCTCAATCCATGCCAC R: CGGGAGTTTCGGAGAA	(CAA) ₅	203–209	60	NED	KT285029	Pentatricopeptide repeat-containing protein At3g49240 [<i>Ricinus communis</i>]	XP_002516677.1	2.00E-05
10128	F: TCCAGGGTCGAGATTTGG R: GCAAAACCAAGAAAGCCGT	(TTC) ₅	150–159	59	NED	KT285030	Rho GTPase-activating protein 3-like isoform X3 [<i>Populus euphratica</i>]	XP_0111039002.1	1.00E-23
10809	F: GAAGAGCTGAAAAGGCAACCT R: TTCCTTTGCCCTCAGCTT	(TGTGG) ₆	196–231	60	HEX	KT285032	22.7 kDa class IV heat shock protein [<i>Ricinus communis</i>]	XP_002521274.1	4.00E-06
10960	F: GGCATCTCTTATTCCTCC R: CTGAAAAGAAAACGGAGCG	(TCCTT) ₅	211–236	59	NED	KT285033	Uncharacterized protein LOC8265384 [<i>Ricinus communis</i>]	XP_002528323.1	0.072
10997	F: TCAAGCATGCCAARAAGGT R: TGCATGAACAAGGTGCC	(AGA) ₇	233–242	60	NED	KT285034	Probable BOI-related E3 ubiquitin-protein ligase 3 [<i>Atropa curcas</i>]	XP_012081782.1	1.00E-32
16226	F: TGGCATAAGAGTGCAACCA R: TGATGATGTGAAAACCTCCA	(CAT) ₇	232–238	58	6-FAM	KT285035	No hit	—	—
16859	F: CACAACACACACACCAACCA R: TTTGAAAATTTGGAAACCCA	(TAC) ₇	160–178	59	HEX	KT285036	No hit	—	—
22194	F: CCCTGTTCTGTGGTCTG R: GAAGAAGAGTGTGATGTC	(TTTGG) ₆	226–241	59	HEX	KT285037	Amino acid permease 6 [<i>Ricinus communis</i>]	XP_002510013.1	6.00E-05
23632	F: GCGACCAAGGCGATGAA R: TCTTCTGCCCTCAGCATTT	(AGTG) ₆	224–254	60	HEX	KT285038	Dynamamin-related protein 3A isoform X1 [<i>Ricinus communis</i>]	XP_015572520.1	2.00E-13
24490	F: AAGGTAAGGTTGCCACG R: CAAGAGGCTCATCCACC	(TGGA) ₇	180–208	60	6-FAM	KT285039	No hit	—	—
25334	F: CACAACCTCCACCGATCA R: ACCGCTAGAACTCGCTGC	(CCT) ₉	175–193	59	6-FAM	KT285040	No hit	—	—
25439	F: TCACCGGATTTGACGA R: CAGRAACCCACCTAGAGAA	(TGAGGC) ₇	147–177	59	HEX	KT285041	No hit	—	—
26221	F: ATGGGACATGATGTTGG R: GCCTTTGTTCTGTTGAGAA	(TGA) ₇	153–189	60	NED	KT285042	Transcription factor PIF7 isoform X2 [<i>Vitis vinifera</i>]	XP_010663294.1	2.00E-07
26474	F: TGGACCATCACCATCAC R: CCCCTCAACTCAATCCATCA	(AG) ₁₁	220–226	60	HEX	KT285043	Conserved hypothetical protein [<i>Ricinus communis</i>]	EEF49157.1	3.00E-12

Note: T_a = annealing temperature.

TABLE 2. Genetic properties of the 18 novel polymorphic EST-SSR markers developed in four populations of *Speranskia tuberculata*.^a

Locus	YA (N = 6)			XZ (N = 6)			YT (N = 6)			KQ (N = 6)			Total	Mean	
	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e		A	H _o
8852	4	0.500	0.708	3	0.167	0.292	1	0	0	4	0.333	0.597	4	0.250	0.626
9441	5	0.833	0.736	1	0	0	2	0	0.278	2	0.167	0.153	5	0.250	0.386
9832	2	0.167	0.486	1	0	0	1	0	0	1	0	0	2	0.042	0.187
10026	3	0.250	0.656	3	0.333	0.486	4	0.333	0.681	2	0.333	0.278	7	0.318	0.705
10117	1	0	0	3	0.333	0.500	2	0.167	0.153	1	0	0	3	0.125	0.192
10128	1	0	0	4	0.333	0.597	2	0	0.278	1	0	0	4	0.087	0.271
10809	2	0.833	0.486	4	0.500	0.597	3	0.333	0.292	2	0.167	0.153	5	0.458	0.556
10960	3	0.667	0.611	3	0.500	0.569	4	0.167	0.625	4	0.667	0.694	6	0.500	0.752
10997	3	0.500	0.569	2	0.167	0.375	4	0.667	0.653	3	0.500	0.542	4	0.458	0.588
16226	1	0	0	2	0	0.278	2	0.833	0.486	1	0	0	2	0.208	0.305
16859	4	0.500	0.514	3	0.333	0.292	3	0.667	0.569	2	0.167	0.375	5	0.417	0.531
22194	3	0.333	0.500	3	0.167	0.292	1	0	0	4	0.667	0.708	4	0.292	0.440
23632	5	0.667	0.681	2	0.167	0.486	2	0	0.320	2	0.400	0.480	6	0.318	0.675
24490	6	0.833	0.806	4	0.667	0.681	3	0.500	0.500	3	0.500	0.625	8	0.625	0.827
25334	2	0.167	0.375	4	0	0.667	4	0.833	0.653	3	0.667	0.500	7	0.417	0.806
25439	3	0.833	0.653	4	1.00	0.708	3	0.833	0.667	4	1.000	0.681	6	0.917	0.748
26221	4	0.500	0.653	5	0.333	0.750	7	0.833	0.819	4	0.500	0.625	11	0.542	0.826
26474	4	0.667	0.681	4	0.200	0.700	2	0	0.500	4	0.333	0.597	4	0.304	0.717

Note: A = number of alleles per locus; H_e = expected heterozygosity; H_o = observed heterozygosity; N = number of individuals sampled.

^aLocality and voucher information are provided in Appendix 1.

amplifications with the same protocol. The labeled PCR products were analyzed on an ABI 3730 DNA Analyzer with a GeneScan 500 LIZ Size Standard (Applied Biosystems). Allele sizes were called using GeneMarker version 2.6.0 (SoftGenetics, State College, Pennsylvania, USA). Number of alleles per locus (A), observed heterozygosity (H_o), and expected heterozygosity (H_e) were calculated using GenAlEx version 6.2 (Peakall and Smouse, 2006).

Eighteen of the 20 candidate markers showed polymorphisms among the four populations of *S. tuberculata*. The corresponding sequences of these markers were deposited in GenBank (Table 1). The number of alleles per locus ranged from two to 11, H_e ranged from 0.187 to 0.827, and H_o ranged from 0.042 to 0.917 (Table 2).

Cross-species amplification of the 18 newly developed polymorphic markers was tested in 24 *S. cantonensis* individuals from a single population (Ruyuan, Guangdong; Appendix 1), using the same procedures described above. Thirteen loci (72.22%) were successfully amplified in all *S. cantonensis* individuals tested, of which six showed polymorphisms (Table 3).

CONCLUSIONS

These 18 novel polymorphic SSR markers will be used to evaluate impacts of recent habitat fragmentation on the genetic

TABLE 3. Polymorphisms at the 13 successfully cross-amplified EST-SSR markers in single population samples of *Speranskia cantonensis* (N = 24).^a

Locus	A	H _o	H _e	GenBank accession no. ^b
8852	1	0	0	KT312943
9441	4	0.042	0.666	KT312944
9832	1	0	0	KT312945
10026	1	0	0	KT312946
10117	2	0	0.375	KT312947
10128	1	0	0	KT312948
10960	1	0	0	KT312949
10997	1	0	0	KT312950
16226	1	0	0	KT312951
23632	4	0	0.663	KT312952
24490	2	0	0.500	KT312953
25334	3	0.250	0.288	KT312954
25439	3	0	0.625	KT312955

Note: A = number of alleles per locus; H_e = expected heterozygosity; H_o = observed heterozygosity; N = number of individuals sampled.

^aLocality and voucher information are provided in Appendix 1.

^bGenBank accession numbers are for the cross-amplified markers in *Speranskia cantonensis*.

diversity and structure of *S. tuberculata*, and to develop suitable conservation strategies for the species. Of these SSR markers developed in *S. tuberculata*, 13 were successfully amplified in single population samples of the related species *S. cantonensis*, extending their potential usefulness for future research in the genus *Speranskia* (e.g., comparisons of genetic diversity).

LITERATURE CITED

FU, L. M., B. F. NIU, Z. W. ZHU, AND W. LI. 2012. CD-HIT: Accelerated for clustering the next-generation sequencing data. *Bioinformatics (Oxford, England)* 28: 3150–3152.

GRABHERR, M. G., B. J. HAAS, M. YASSOUR, J. Z. LEVIN, D. A. THOMPSON, I. AMIT, X. ADICONIS, ET AL. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology* 29: 644–652.

HAYDEN, M. J., T. M. NGUYEN, A. WATERMAN, G. L. McMICHAEL, AND K. J. CHALMERS. 2008. Application of multiplex-ready PCR for fluorescence-based SSR genotyping in barley and wheat. *Molecular Breeding* 21: 271–281.

HWANG, S. M. 1989. A notes on genera *Speranskia* in China (Euphorbiaceae). *Bulletin of Botanical Research* 9: 37–40.

JU, M. M., H. C. MA, P. Y. XIN, Z. L. ZHOU, AND B. TIAN. 2015. Development and characterization of EST-SSR markers in *Bombax ceiba* (Malvaceae). *Applications in Plant Sciences* 3: 1500001.

MAZZIO, E., R. BADISA, N. MACK, S. DEIAB, AND K. F. A. SOLIMAN. 2014. High throughput screening of natural products for anti-mitotic effects in MDA-MB-231 human breast carcinoma cells. *Phytotherapy Research* 28: 856–867.

PEAKALL, R., AND P. E. SMOUSE. 2006. GenAlEx 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6: 288–295.

ROZEN, S., AND H. SKALETSKY. 1999. Primer3 on the WWW for general users and for biologist programmers. In S. Misener and S. A. Krawetz [eds.], *Methods in molecular biology*, vol. 132: Bioinformatics methods and protocols, 365–386. Humana Press, Totowa, New Jersey, USA.

THIEL, T., W. MICHALEK, R. K. VARSHNEY, AND A. GRANER. 2003. Exploiting EST databases for the development and characterization of gene-derived SSR-markers in barley (*Hordeum vulgare* L.). *Theoretical and Applied Genetics* 10: 411–422.

APPENDIX 1. Locality information for the sampled populations of *Speranskia tuberculata* and *S. cantonensis* tested in this study.

Population	Species	Collection locality	<i>N</i>	Geographic coordinates	Altitude (m)	Voucher no. ^a
YA	<i>S. tuberculata</i> (Bunge) Baill.	Yan'an, Shaanxi	6	36°35'N, 109°29'E	1061	TB2014087
XZ	<i>S. tuberculata</i>	Xinzhou, Shanxi	6	39°19'N, 113°34'E	1160	TB2014117
YT	<i>S. tuberculata</i>	Yantai, Shandong	6	37°17'N, 121°44'E	120	TWYT02
KQ	<i>S. tuberculata</i>	Chifeng, Inner Mongolia	6	42°57'N, 118°59'E	631	TB2013153
RU	<i>S. cantonensis</i> (Hance) Pax & K. Hoffm.	Ruyuan, Guangdong	24	24°59'N, 113°08'E	650	FL2014098

Note: *N* = number of individuals.

^aVoucher specimens deposited at the Herbarium of Southwest Forestry University (SWFC), Kunming, China.