

## **A New, Unusual Rhynchonellide Brachiopod with a Strophic Shell from the Silurian of Iran**

Authors: Popov, Leonid E., Hairapetian, Vachik, Pour, Mansoureh Ghobadi, and Modzalevskaya, Tatiana L.

Source: *Acta Palaeontologica Polonica*, 60(3) : 747-754

Published By: Institute of Paleobiology, Polish Academy of Sciences

URL: <https://doi.org/10.4202/app.00030.2013>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# A new, unusual rhynchonellide brachiopod with a strophic shell from the Silurian of Iran

LEONID E. POPOV, VACHIK HAIRAPETIAN, MANSOUREH GHOBADI POUR,  
and TATIANA L. MODZALEVSKAYA



Popov, L.E., Hairapetian, V., Ghobadi Pour, M., and Modzalevskaya, T.L. 2015. A new, unusual rhynchonellide brachiopod with a strophic shell from the Silurian of Iran. *Acta Palaeontologica Polonica* 60 (3): 747–754.

A new, unusual rhynchonellide brachiopod *Jafarirhynchus alatus* assigned to the newly established family Jafarirhynchidae is described from the Silurian (Telychian) of the Boghu Mountains in east-central Iran. It forms a low diversity association with the spiriferide *Striispirifer? ocissimus*, which exhibits well preserved calcified brachial supports. A strophic shell, well-developed ventral interarea and liberossessile mode of life make this taxon unique among Palaeozoic rhynchonellide brachiopods. In spite of a superficial similarity to spiriferides and the atrypide family Davidsonioidea, *Jafarirhynchus* retains the typical rhynchotrematoid cardinalia with a septalium supported by the median septum, a septiform cardinal process and long, raduliform crura. It is considered as an offshoot of the local lineage, which includes two successive species of *Stegocornu* (family Rhynchotrematidae) which proliferated in Central Iran and adjacent Afghanistan during Aeronian time.

**Key words:** Rhynchonellida, morphology, taxonomy, Silurian, Telychian, Iran.

Leonid E. Popov [leonid.popov@museumwales.ac.uk], Department of Geology, National Museum of Wales, Cathays Park, Cardiff CF10 3NP, UK.

Vachik Hairapetian [vachik@khuisf.ac.ir], Department of Geology, Khorasgan (Isfahan) Branch, Islamic Azad University, PO Box 81595-158, Isfahan, Iran.

Mansoureh Ghobadi Pour [mghobadipour@yahoo.co.uk], Department of Geology, Faculty of Sciences, Golestan University, Gorgan 49138-15739, Iran.

Tatiana L. Modzalevskaya [Tatiana\_Modzalevskaya@vsegei.ru], Department of Stratigraphy and Palaeontology, Russian Geological Research Institute (VSEGEI), 74 Sredny prospect, St. Petersburg 199106, Russia.

Received 2 October 2013, accepted 24 January 2014, available online 28 January 2014.

Copyright © 2015 L.E. Popov et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Introduction

The existing record of Iranian Silurian brachiopods remains sparse and incomplete apart from the rich Aeronian brachiopod faunas of Central Iran and Kopet-Dagh, which are well documented in publications by Cocks (1979), Brice (1999), and Hairapetian et al. (2012). The major focus of this paper is to describe a small, oligotaxic brachiopod association from the Telychian deposits of the Boghu Mountains in the vicinity of the town of Kashmar, which are situated at the northern margin of the Tabas block in the north-eastern part of the Central Iran plate. The most distinctive component of this association is an unusual rhynchonellide with a strophic shell and a well-developed ventral interarea assigned to a new genus and species *Jafarirhynchus alatus*.

**Institutional abbreviations.**—AEU, Azad University, Esfahan, Iran; NMW, Department of Geology, National Museum

of Wales, Cardiff, UK; CNIGR, F. N. Chernyshev Central Research and Geological Exploration Museum, St Petersburg, Russia.

**Other abbreviations.**—Lv, Ld, maximum length of ventral and dorsal valves; max, maximum observed size; min, minimum observed size; N, number of specimens; S, standard deviation from the mean; SLw, SLh, maximum width and height of the sulcus/median folds; W, T; maximum width, thickness of the shell; X, mean.

## Geological and geographical setting

The Boghu Mountains are situated about 25 km south-west of the town of Kashmar (Fig. 1). An outline of the Lower Palaeozoic geology of the area can be found in publications by




Fig. 1. **A.** Geographic and geologic setting of the studied material showing position of studied section (modified after Taheri 2001). **B.** Schematic map showing position of studied locality (asterisk). **C.** Stratigraphical column of the Boghu Section showing the informal lithostratigraphical subdivision of the Niur Formation, position of fossil samples, and stratigraphical distribution of brachiopods and conodonts. Arrows below and above the log indicate that the section continues in both directions (for a complete log see Hairapetian et al. 2012). Arrows at the lowermost of distribution line of some taxa indicates that these also occur in lower strata.

Ghavidel-Syooki (2003) and Hairapetian et al. (2012). The best exposures are situated on the southern flank of the Boghu Mountains (Fig. 1A). There the Silurian Niur Formation rests unconformably on siliciclastic sediments (green shale member) of the Ordovician Katkoyeh Formation (Fig. 1B). The base of the measured section is situated at 34°4'6.2"N, 58°15'48.8"E, altitude 1222 m. The Niur Formation comprises intercalating beds of argillite, sandstone, siltstone, and bioclastic limestone. The Aeronian part of the section is characterised by a conodont assemblage including *Distomodus staurogathoides* (Walliser, 1964), *Wurmiella* ex. gr. *excavata* (Branson and Mehl, 1933), *Ozarkodina* spp. and *Oulodus?* sp., and a low diversity brachiopod fauna dominated by the rhynchonellides *Stegocornu procerum* Dürkoop, 1970 and *Stegocornu denisae* Popov, Modzalevskaia, and Ghobadi Pour in Hairapetian et al., 2012 (Fig. 1B).

The brachiopods *Jafarirhynchus alatus* gen. et sp. nov. and *Striispirifer? ocissimus* Popov, Modzalevskaia, and Ghobadi Pour in Hairapetian et al., 2012 were sampled from the bed of grey bioclastic limestone at 12 m above the base of unit 22 (Fig. 1; sample K33/1). It is separated from the last occurrence of *Wurmiella* ex. gr. *excavata* (Fig. 1B; sample K32) by a barren sandstone unit about 61.8 m thick (Fig. 1B; Unit 21). A conodont assemblage collected from sample K36 at 23.3 m above sample K33/1 includes *Distomodus staurogathoides* and *Arianagnathus jafariani* Männik et al. (2015). The latter was known from the Derenjal Mountains, east-central Iran associated with several taxa including *Pterospathodus amorphognathoides lennarti* Männik, 1998, suggesting the *P. amorphognathoides lennarti* Conodont Zone of mid Telychian age (Männik et al. 2013, 2015).

The uppermost conodont occurrences in the studied succession are in samples K37 and K38 (43.4 m above sample K33/1; Fig. 1B). They contain *Kockelella ranuliformis* (Walliser, 1964), suggesting an early Sheinwoodian (early Wenlock) age (Hairapetian et al. 2012).

## Systematic palaeontology

Phylum Brachiopoda Dumeril, 1806

Subphylum Rhynchonelliformea Williams, Carlson, Brunton, Holmer, and Popov, 1996

Class Rhynchonellata Williams, Carlson, Brunton, Holmer, and Popov, 1996

Order Rhynchonellida Kuhn, 1949

Superfamily Rhynchotrematoidea Schuchert, 1913

Family Jafarirhynchidae nov.

*Diagnosis.*—Aberrant Rhynchotrematoidea with a coarsely costate, alate, strophic shell, a wide, straight hinge line, a well-developed ventral interarea, a strong ventral sulcus and a dorsal median fold extending from the beak. Anterior commissure strongly bisulcate. Dental plates and septalium

supported by a dorsal medium septum; cardinal process septiform.

*Remarks.*—The family Jafarirhynchidae is unique within the order Rhynchonellida in having a strophic shell with a wide, straight hinge line and a well-developed ventral interarea. These features in combination with cyrtomatodont teeth are considered diagnostic for spiriferides (Jaanusson 1971); however, the family Jafarirhynchidae has dorsal cardinalia with a septalium supported by the median septum, a septiform cardinal process and long, raduliform crura, typical for the rhynchonellide superfamily Rhynchotrematoidea. The serial sections demonstrate the lack of calcified spiralia.

### Genus *Jafarirhynchus* nov.

*Type species:* *Jafarirhynchus alatus* sp. nov.; monotypic; see below.

*Etymology:* After Professor Mohammad-Ali Jafarian (1935–2013) in appreciation of his valuable contributions to the Palaeozoic palaeontology and stratigraphy of Iran.

*Diagnosis.*—As for the family.

*Stratigraphic and geographic range.*—Silurian, Llandovery, Telychian; Niur Formation of east-central Iran.

### *Jafarirhynchus alatus* sp. nov.

Figs. 2, 3A–C; Table 1.

*Etymology:* After the alate shape of the shell.

*Type material:* Holotype: AEU1500, a pair of conjoined valves. Paratypes: AEU1501, 1501–1510, NMW 2011.11G.400–456, CNI-GR136/12600, total 62 articulated shells and four ventral valves from sample K-33/1.

*Type locality:* Boghu Mountains about 25 km south-west of the town of Kashmar, Iran.

*Type horizon:* Sample K-33/1, lower–middle Telychian, Llandovery, Silurian.

*Measurements* (in mm): Paratype AEU1501, Lv = 14.0, Ld = 12.3, W = 25.6, T = 10.0, SLw = 7.6, SLt = 8.3; AEU1503, Lv = 17.6, Ld = 15.6, W = 28.1, T = 13.6, SLw = 9.9, SLt = 9.4.

*Diagnosis.*—As for the genus.

*Description.*—Shell variably asymmetrical, strophic, dors-ibiconvex, strongly alate, transverse, subtriangular in outline about three-fifths as long as wide, with maximum width at or slightly anterior to hinge line, and two-thirds as thick

Table 1. Basic measurements (in mm) of 13 shells of *Jafarirhynchus alatus* gen. et sp. nov.; Llandovery, Telychian, Niur Formation, sample K33/1, Boghu Mountains. Abbreviations: Lv, Ld, maximum length of ventral and dorsal valves; max, maximum observed size; min, minimum observed size; n, number of specimens; s, standard deviation from the mean; SLw, SLh, maximum width and height of the sulcus/median folds; W, T; maximum width, thickness of the shell; x, mean.

	Lv	Ld	W	T	SLw	SLh	Lv/ W	Ld/ W	T/ Lv	SLw/ W
n	13	13	13	13	13	13	13	13	13	13
x	15.7	14.1	26.8	10.2	8.7	9.8	58.5%	52.6%	65.3%	32.5%
s	1.20	1.21	1.35	1.17	1.15	1.67	4.4	4.5	6.2	4.9
min	14.2	12.25	24.7	8.7	6.8	7.4	48.5%	43.4%	55.8%	23.2%
max	17.8	15.8	29.25	13.5	10.3	14.2	63.6%	58.9%	75.8%	41.7%




Fig. 2. Rhynchonellide brachiopod *Jafarirhynchus alatus* gen. et sp. nov. from Llandoverly, Telychian, Niur Formation, sample K33/1, Boghu Mountains, CNIGR136/12600. A. Transverse serial sections (L = 18.4; W = 23.2; T = 10.5). B. Sagittal section of the shell. C. Dorsal valve interior. Numbers indicate distance in mm from the tip of the ventral umbo.

as long. Cardinal extremities slightly rounded. Lateral commissures almost straight, converging anteriorly. Anterior commissure strongly bisulcate. Ventral valve gently convex with a sulcus originating at the umbo and strongly deepening anterior to mid-length, bearing a single rib. Ventral interarea, planar, almost orthocline, with a broad, triangular delthyrium completely covered by the medially merged deltidial plates. Lateral profile of dorsal valve strongly convex with maximum height between mid-length and the anterior valve margin or at the anterior margin. Median fold with steep lateral slopes originating at the umbo, bearing two strong, high ribs, occupying about one-third maximum valve width at the anterior margin. Radial ornament of 17–22 strong, tubular ribs separated by interspaces of equal size. Two prominent subangular ribs on the dorsal median fold and a pair of subangular ribs on the flanks of ventral sulcus about twice as high and wide as other ribs. Concentric ornament of fine, regularly spaced concentric lamellae, becoming more prom-

inent and densely crowded close the anterior margin of the mature individuals.

Ventral interior with strong, oblique teeth supported by the short, strongly thickened dental plates. Ventral muscle field strongly impressed with adductor scars completely enclosed by larger diductor scars. Strongly thickened pedicle callist mainly occupying the floor of the small delthyrial cavity. Dorsal interior with massive hinge plates and a small septalium supported by a strongly thickened, short median septum. Cardinal process septiform. Crura faint, radulifer, gently curved dorsolaterally. No calcified brachial supports.

*Stratigraphic and geographic range.*—Niur Formation, Aeronian, Llandoverly, Silurian, Central Iran.

Order Spiriferida Waagen, 1883

Suborder Spiriferidina Waagen, 1883

Superfamily Cyrtioidea Frederiks, 1924




Fig. 3. Rhynchonellide brachiopod *Jafarirhynchus alatus* gen. et sp. nov. (A–C) and spiriferide *Striispirifer?* *ocissimus* Popov, Modzalevskaya, and Ghobadi Pour in Hairapetian et al., 2012 (D, E) from Niur Formation, Telychian, Llandoverly, Silurian, sample K33/1, Boghu Mountains. A. AEU1500, holotype, a pair of conjoined valves, in dorsal (A<sub>1</sub>), ventral (A<sub>2</sub>), lateral (A<sub>3</sub>), and anterior (A<sub>4</sub>) views, enlarged ventral interarea (A<sub>5</sub>). B. AEU1501, a pair of conjoined valves, in anterior (B<sub>1</sub>), ventral (B<sub>2</sub>), dorsal (B<sub>3</sub>), and lateral (B<sub>4</sub>) views of a strongly asymmetrical specimen. C. AEU1502, dorsal view of a pair of conjoined valves (C<sub>1</sub>), enlarged surface ornament (C<sub>2</sub>). D. AEU1511, a pair of conjoined valves, in ventral (D<sub>1</sub>), lateral (D<sub>2</sub>), and dorsal (D<sub>3</sub>) views. E. AEU1512, ventral valve exterior. Scale bars 2 mm.

## Family Cyrtioidea Frederiks, 1924

## Subfamily Eospiriferinae Schuchert, 1929

Genus *Striispirifer* Cooper and Muir-Wood, 1951

*Type species: Delthyris niagarensis* Conrad, 1842; Niagara Group, Wenlock, Silurian, Lockport, New York, USA.

*Striispirifer? ocissimus* Popov, Modzalevskaya, and Ghobadi Pour in Hairapetian et al., 2012

Figs. 3D, E, 4.

2012 *Striispirifer? ocissimus* Popov, Modzalevskaya, and Ghobadi Pour 2012; Hairapetian et al. 2012: 101, figs. 6R–T, 11.

*Material*.—AEU1510–1512, NMW 2011.11G.457–469. Total six pairs of conjoined valves and nine ventral valves.

*Remarks*.—Specimens from the Telychian have spiralia with up to seven whorls. Except for their smaller size, these shells are morphologically identical to those of *Striispirifer? ocissimus* from the underlying Aeronian deposits (Hairapetian et al. 2012) and are considered here as conspecific.

*Stratigraphic and geographic range*.—Niur Formation, Aeronian, Llandowery, Silurian, Central Iran.

## Discussion

There is a general consensus that the earliest “astrophic spire-bearers” with calcified brachial supports evolved monophyletically from a rhynchonellide ancestor (Copper and Gourvennec 1996; Alvarez et al. 1998; Popov et al. 1999), while the astrophic shell can be considered as plesiomorphic feature inherited by the early spire-bearers from rhynchonellides.

Rong and Zhan (1996) also published convincing evidence, subsequently supported by a study of the early spire-bearing brachiopods from Kazakhstan (Popov et al. 1999) that spiriferides evolved from an atrypide (probably lissatrypoid) ancestor with a simple, centrally directed or planospiral spirulum. Therefore the strophic shell in spiriferides is an apomorphic character. Remarkably, the strophic shell evolved repeatedly several times in “astrophic spire-bearers”, including the atrypide superfamily Davidsonioidea, the Devonian athyridide genera *Anathyrella* Khalifin in Gratsianova et al., 1961, and *Anathyris* Peetz, 1901, the Permian subfamily Comelicaninae and the Mesozoic superfamily Koninckinoidea (for details see Alvarez and Rong 2002). Wright (1979) argued in favour of the origin of spiriferides from orthides, based on a superficial spiriferide appearance of some genera of the family Platystrophiidae; however, this view was not supported by subsequent studies by Rong and Zhan (1996).

There is no previous record of a similar strophic shell in rhynchonellides through their entire evolutionary history from the Ordovician to Recent. Hence the family Jafarirhynchidae is unique within the order Rhynchonellida in having a strophic shell with a wide, straight hinge line and a well-developed ventral interarea. These features, in combination with cyrtomatodont dentition, are considered diagnostic for the spiriferides (Jaanusson 1971); however, brachiopods assigned to the family Jafarirhynchidae lack calcified brachial supports and show the dorsal cardinalia with a septalium supported by the median septum, a septiform cardinal process and long, raduliform crura, typical for the rhynchonellide superfamily Rhynchotrematoidea. From other side, shells of *Striispirifer? ocissimus*, which occur at the same locality, have well preserved calcified spiralia (Fig. 4).




Fig. 4. Transverse serial sections of a cyrtoid spiriferid brachiopod *Striispirifer? ocissimus* Popov, Modzalevskaya, and Ghobadi Pour in Hairapetian et al., 2012; NMW 2011.11G.457 from Llandowery, Telychian, Niur Formation, sample K33/1, Boghu Mountains. Numbers indicate distance in mm from the tip of the ventral umbo.

*Jafarirhynchus* itself can be linked to the Aeronian lineage of rhynchotrematid *Stegocornu* represented by two succeeding species *S. procerum* Dürkoop, 1970 and *S. denisae* Popov, Modzalevskaya, and Ghobadi Pour in Hairapetian et al., 2012 in a number of localities across Central Iran (Hairapetian et al. 2012; Popov and Cocks 2013). Indeed, *S. denisae* shows an erect beak, an almost straight hinge line that is only slightly shorter than maximum shell width and a delthyrium mainly covered by converging deltidial plates (Hairapetian et al. 2012: fig. 6S, T). In comparison with the ancestral taxon *S. procerum*, *S. denisae* is also characterized by its wider, dorsoventrally compressed shell and narrower fold and sulcus in relation to the maximum shell width. All these tendencies are progressed further in *Jafarirhynchus alatus*, which possibly represents the last member of that lineage.

The delthyrium in *Jafarirhynchus alatus* is completely sealed by the medially merged deltidial plates and there is no functional pedicle foramen in mature individuals. This suggests a liberossessile mode of life for this species, which is unusual in rhynchonellides. *Jafarirhynchus alatus* is a dominant species in the oligotaxic rhynchonellid association with *Striispirifer? ocissimus* as a minor component. In the early Telychian of the Kashmar area this association replaced the *Stegocornu* association characteristic of the Aeronian (Hairapetian et al. 2012). Both associations inhabited the shallow shelf environments on a carbonate ramp and can be assigned to the Benthic Assemblage Zone 2 of Boucot (1975).

*Jafarirhynchus* shows a remarkable similarity to atrypides of the superfamily Davidsonioidea in the morphology of the ventral interarea and delthyrial covers (Copper 2002: fig. 937.2); however, that similarity is superficial and represents no more than homoplasy, since *Jafarirhynchus* retains typical rhynchonellide cardinalia with a septalium. Recent rhynchonellides have a spirolophe lophophore with dorsally directed apices of cones. They lack calcified supports, but it is likely that their basic morphology remained unchanged since the early Palaeozoic. The strongly transverse shell renders this type of lophophore dysfunctional in *Jafarirhynchus*, therefore the eversion model of lophophore cones developed from the laterally directed axis of spiralia, proposed by Rong and Zhan (1996) for spiriferides, was probably applicable also for *Jafarirhynchus*.

## Conclusions

A strophic shell with well-developed ventral interarea convergently similar to those of spiriferides, and a liberossessile mode of life make *Jafarirhynchus* and the family Jafarirhynchidae unique among the Palaeozoic rhynchonellide brachiopods. In spite of its unusual morphology and life style, previously unknown in the early Palaeozoic representatives of the order, *Jafarirhynchus* retains characteristic rhynchotrematoid cardinalia and can be considered as a terminal member of a local lineage, which evolved in Central Iran

during the Aeronian, where it is represented by two succeeding species of *Stegocornu* (Hairapetian et al. 2012).

## Acknowledgments

We are grateful to Peep Männik (Tallinn University of Technology, Estonia) and C. Giles Miller (Natural History Museum, London, UK) for conodont identifications and to Hasan Hejazi (AEU) and Amir Akbari (Esfahan, Iran) for assistance in fieldworks. We thank L. Robin M. Cocks (Natural History Museum, London, UK) and David Harper (Durham University, UK), for their insightful reviews. Leonid Popov acknowledges support from NMW. Mansoureh Ghobadi Pour thanks the NMW for logistical support on her visits to Cardiff; her work in Iran is supported by the Golestan University. This paper is a contribution to IGCP591 Project: The Early to Middle Paleozoic Revolution.

## References

- Alvarez, F., Rong, J., and Boucot, A.J. 1998. The classification of athyridid brachiopods. *Journal of Paleontology* 72: 827–855.
- Alvarez, F. and Rong, J. 2002. Order Athyridida. In: R.L. Kaesler (ed.), *Treatise on Invertebrate Paleontology. Part H, Brachiopoda (Revised)*, 1475–1614. Geological Society of America and the University of Kansas Press, Boulder.
- Boucot, A.J. 1975. *Evolution and Extinction Rate Controls*. 426 pp. Elsevier Scientific Publishing Company, New York.
- Branson, E.B. and Mehl, M.G. 1933. Conodonts from the Bainbridge Formation (Silurian) of Missouri. *University of Missouri Studies* 8: 39–52.
- Brice, D. 1999. Middle(?) Silurian rhynchonellid and spiriferid brachiopod faunas from eastern central Iran. *Annales de la Société géologique du Nord* 7: 5–12.
- Cocks, L.R.M. 1979. A silicified brachiopod fauna from the Silurian of Iran. *Bulletin of the British Museum (Natural History), Geology* 32: 25–42.
- Conrad, T.A. 1842. Observations on the Silurian and Devonian Systems of the United States, with descriptions of new organic remains. *Journal of the Academy of Natural Sciences of Philadelphia* 8: 228–280.
- Cooper, G.A. and Muir-Wood, H.M. 1951. Brachiopod homonyms. *Journal of Washington Academy of Sciences* 41: 195–196.
- Copper, P. 2002. Atrypida. In: R.L. Kaesler (ed.), *Treatise on Invertebrate Paleontology, Part H, Brachiopoda, Revised* 4, 1377–1474. Geological Survey of America and University of Kansas Press, Boulder.
- Copper, P. and Gourvenec, R. 1996. Evolution of spire bearing brachiopods (Ordovician–Jurassic). In: P. Copper and J. Jin (eds.), *Brachiopods. Proceedings of the Third International Brachiopod Congress. Sudbury, Ontario, Canada, 2–5 September, 1995*, 81–88. A.A. Balkema, Rotterdam-Brookfield.
- Duméril, A.M.C. 1806. *Zoologie analytique ou méthode naturelle de classification des animaux*. xxiv + 344 pp. Allais, Paris.
- Dürkoop, A. 1970. Brachiopoden aus dem Silur, Devon und Karbon in Afghanistan. *Palaeontographica, series A* 134: 153–225.
- Frederiks, G.N. 1924. Paleontological studies. 2. On Upper Carboniferous spiriferids from the Urals [in Russian]. *Izvestiâ Geologičeskogo Komiteta* 38 (for 1919): 295–324.
- Ghavidel-Syooki, M. 2003. Palynostratigraphy and paleogeography of Lower Paleozoic Strata at Kuh-e-Boghrou, Southwest of Kashmar City, at Eastern Central Iran. *Iranian International Journal of Science* 4: 181–207.
- Gratsianova, R.T. [Gracânova, R.T.], Zinchenko, B.A. [Zinčenko, B.A.], and Kul'kov, N.K. 1961. Type Brachiopoda [in Russian]. In: L.L. Khal'fin (ed.), *Biostratigrafiâ paleozoâ Sajno-Altajskoj gornoj oblasti. Tom 2. Paleozoj. Trudy SNIIGGIMS (Sibirskii Naučno-Issledovatel'skij Institut Geologii, Geofiziki i Mineralogii Sibiri)* 20: 1–850.



- Hairapetian, V., Ghobadi Pour, M., Popov, L., and Modzalevskaya, T.L. 2012. *Stegocornu* and associated brachiopods from the Silurian (Llandovery) of Central Iran. *Estonian Journal of Earth Sciences* 61: 82–104.
- Jaanusson, V. 1971. Evolution of the brachiopod hinge. *Smithsonian Contributions to Paleobiology* 3: 33–46.
- Kuhn, O. 1949. *Lehrbuch der Paläozoologie*. 326 pp. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart.
- Männik, P. 1998. Evolution and taxonomy of the Silurian conodont *Pterospaathodus*. *Palaeontology* 41: 1001–1050.
- Männik, P., Miller, C.G., and Hairapetian, V. 2013. Conodonts from the Niur Formation (Silurian) from the Derenjal Mountains, Central Iran. *Geological Magazine* 150: 639–650.
- Männik, P., Miller, C.G., and Hairapetian, V. 2015. A new early Silurian prioniodontid conodont with three P elements from Iran and associated species. *Acta Palaeontologica Polonica* 60: 733–746.
- Rong, J. and Zhan, R. 1996. Brachidia of late Ordovician and Silurian eospiriferines (Brachiopoda) and the origin of the spiriferides. *Palaeontology* 39: 941–977.
- Peetz, H. von 1901. Étude sur la faune de l'étage de Malevkon-Mourayevnia. *Travaux de la Société des Naturalistes de St.-Petersbourg, Section de Géologie et de Mineralogie* 22 (2): 29–118.
- Popov, L.E., Nikitin, I.F., and Sokiran, E.V. 1999. The earliest atrypides and athyridides (Brachiopoda) from the Ordovician of Kazakhstan. *Palaeontology* 42: 625–661.
- Popov, L.E. and Cocks, L.R.M. 2013. The radiation of early Silurian spiriferide brachiopods, with new taxa from the Llandovery of Iran. *Alcheringa* 4: 558–564.
- Schuchert, C. 1929. Classification of brachiopod genera, fossils and recent. In: C. Schuchert and C.M. LeVene (eds.), *Fossilium Catalogus I: Animalia* 42. *Brachiopoda (Generum et Genotyporum Index et Bibliographia)*, 10–25. W. Junk, Berlin.
- Taheri, J. 2001. *Geological Map of Quadrangle Kashmar, scale 1:100,000*. Geological Survey of Iran, Tehran.
- Waagen, W.H. 1883. Salt-Range fossils, Vol. 1, Part 4, *Productus* Limestone fossils, Brachiopoda. *Memoirs of the Geological Survey of India, Palaeontologia Indica (series 13)* 2: 391–546.
- Walliser, O.H. 1964. Conodonten des Silurs. *Abhandlungen der Hessischen Landesamtes Bodenforschung* 41: 1–106.
- Williams, A., Carlson, S.J., Brunton, C.H.C., Holmer, L.E., and Popov, L.E. 1996. A supra-ordinal classification of the Brachiopoda. *Philosophical Transactions of the Royal Society, Biological Sciences* 35: 1171–1193.
- Wright, A.D. 1979. The origin of the spiriferidine brachiopods. *Lethaia* 12: 29–33.