

The Pustulated Temnospondyl Revisited—a Plagiosternine Plagiosaurid from the Lower Triassic of Brazil

Authors: Dias-Da-Silva, Sérgio, and Milner, Andrew R.

Source: *Acta Palaeontologica Polonica*, 55(3) : 561-563

Published By: Institute of Paleobiology, Polish Academy of Sciences

URL: <https://doi.org/10.4202/app.2009.0131>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



The pustulated temnospondyl revisited—a plagiosternine plagiosaurid from the Lower Triassic of Brazil

SÉRGIO DIAS-DA-SILVA and ANDREW R. MILNER

A recent contribution published in this journal (Dias-da-Silva and Ilha 2009) reported a dermal skull fragment indicating the presence of a putative plagiosauroid temnospondyl in the Lower Triassic Sanga do Cabral Formation of the Paraná Basin, Southern Brazil. The taxonomic assignment of this specimen was necessarily tentative as it was based on circumstantial evidence, specifically the presence of a dense pustular ornamentation over four partial dermal skull bones, consideration of the described taxa known to bear such ornamentation, and the stratigraphic and paleobiogeographic range of such taxa. Therefore, Dias-da-Silva and Ilha (2009) could not be totally certain about the plagiosauroid affinities of the new specimen and ascribed it to ?Plagiosauridae. It was particularly difficult to make a precise osteological identification of the specimen and six alternative osteological interpretations were made in comparison to both *Gerrothorax* and *Peltobatrachus* (see Dias-da-Silva and Ilha 2009: fig. 2). In spite of the poor taxonomic resolution, the new specimen raised interesting questions regarding the presence of plagiosauroid stereospondyls in western Gondwana, as well as their evolutionary patterns, biostratigraphic and paleobiogeographic implications. After Dias-da-Silva and Ilha's (2009) contribution was published, new data from Damiani et al. (2009) raised the possibility of narrowing down the taxonomic identity of the plagiosauroid from Brazil. Accordingly, this brief report provides a more precise taxonomic assignation for this material.

Introduction

After Dias-da-Silva and Ilha (2009) went to press, Damiani et al. (2009) published a description of new material of the German plagiosaur *Plagiosuchus pustuliferus*. Incorporated in the comparative section of this paper was a new reconstruction of the skull of another German plagiosaur, *Plagiosternum granulosum*, prepared by Stephanie Gastou (as part of an unpublished degree project; Gastou 2008). This reconstruction, based on new material, included details of the bone arrangements around the cheek and suspensorium that were not previously available (see Warren 1995; Hellrung 2003), and thus allows a more precise taxonomic assignation of the Brazilian plagiosauroid material (under collection number MCN PV 1999) as a plagiosternine plagiosaurid.

Institutional abbreviations.—MCN PV, Paleovertebrates collection, Museu de Ciências Naturais da Fundação Zoobotânica do Estado do Rio Grande do Sul, Porto Alegre, Brazil; SMNS, Staatliches Museum für Naturkunde, Stuttgart, Germany.

Systematic paleontology

Temnospondyli von Zittel, 1888
Stereospondylomorpha Yates and Warren, 2000
Stereospondyli von Zittel, 1888
Plagiosauridae Abel, 1919
Plagiosterninae Shishkin, 1986
Incertae sedis

Locality and horizon: 6.4 km from the beginning of the BR 158 (a federal highway), that connects the municipalities of Santa Maria and Rosário do Sul (Coordinates S29°42'59.45" W53°54'01.76"). *Lystrosaurus* Assemblage Zone, Sanga do Cabral Formation, Lower Triassic, Brazil.

Material.—MCN PV 1999, a dermal skull fragment including four dorsoventrally compressed partial bones (Fig. 1C).

Discussion

Morphological identity.—The new published figure of the skull of *Plagiosternum granulosum* (Damiani et al. 2009: fig. 8C) shows the shape and sutures of the cheek region and suspensorium for the first time in this taxon. Previous authors had to speculate about the exact relationships of the bones in this region (Warren 1995; Hellrung 2003). Comparison of the bone arrangement of the right cheek of *P. granulosum* with the elements in MCN PV 1999 shows an almost exact correspondence (Fig. 1). MCN PV 1999 comprises apparently the almost complete quadratojugal, the posterior half of the jugal, the anterior half of the squamosal, and the lateral part of the postorbital. The lateral margin of the jugal and most of the entire margin of the quadratojugal are preserved, while the remaining edges of the specimen are incomplete. Therefore, bone structure in the preserved fragment allows clear identification as part of a plagiosternine skull roof. However, there still remains the problem of the distribution of the dermal ornament types within the Plagiosauridae.




Fig. 1. Plagiosaurid *Plagiosternum granulosum* Frass, 1889 from the Middle Triassic of Germany. **A.** Schematic line drawing, modified from Damiani et al. (2009); the gray area shows the presumable anatomic correspondence between MCN PV 1999 and *P. granulosum*. Not to scale. **B.** Line drawing of MCN PV 1999, Plagiosterninae indet. from the Lower Triassic Sanga do Cabral Formation (Paraná Basin, Southern Brazil). **C.** Photograph of MCN PV 1999 with its presumable anatomic correspondence to *P. granulosum*.

Dermal ornamentation.—Dias-da-Silva and Ilha (2009) reported MCN PV 1999 as a plagiosaurid based upon its pustular ornamentation, but could not assign the specimen to any subfamily within the Plagiosauridae. Recent workers (Shishkin 1987; Schoch and Werneburg 1998; Warren 2000; Hellrung 2003) have recognized up to three plagiosaur subfamilies and, according to Hellrung (2003), two of these, Plagiosaurinae and Plagiosuchinae, were characterized by pustular ornamentation, while the Plagiosterninae retained reticulate ornamentation—the general temnospondyl condition. Based upon this statement, Dias-da-Silva and Ilha (2009) suggested that MCN PV 1999 might belong to either Plagiosaurinae or Plagiosuchinae, since this Brazilian specimen clearly shows a pustular pattern of ornamentation. However, observations by Stephanie Gastou (personal communication 2009) and ARM on material housed in the collections at SMNS, indicate that the pattern of ornamentation within plagiosaurids is not as strict within subfamilies as stated by Hellrung (2003). In *Plagiosternum*, the ornamentation of the central region of the skull is primitively reticulate, but towards the suspensorium there are pustules on the intersections between the reticular ridges. Moreover, some specimens seem to be more pustular whereas others are more reticulate. For instance, the ornamentation present in *Plagiosternum* specimen SMNS 82002 includes both reticulate and pustular, the latter particularly developed in the cheek region, but its posterolateral bones are an almost perfect match to MCN PV 1999, assuming the above morphological interpretation is correct. The quadratojugal of *Melanopelta antiqua* (Shishkin 1967: fig. 2), interpreted as a plagiosternine from the Lower Triassic of Russia

(Shishkin 1987) also shows pustular ornamentation superimposed on a reticulate base. Nevertheless, MCN PV 1999 does differ from the described European and Russian specimens of *Plagiosternum* in the greater extent of the pustular ornamentation. The fragment is insufficiently diagnostic to permit critical generic diagnosis, for example between *Plagiosternum* and *Melanopelta*, and we therefore restrict its identification as a Plagiosterninae *incertae sedis*. The subfamily Plagiosterninae Shishkin, 1986 comprises *Plagiosternum* (Middle Triassic of Germany and Russia), *Melanopelta*, and *Aranetsia* (Lower and Middle Triassic of Russia, respectively). Unfortunately, plagiosternine specimens are “largely represented by scant cranial and postcranial remains” (Damiani et al. 2009: 349). The stratigraphic age of MCN PV1999 falls within this range (Lower to Middle Triassic) although the paleogeographical range is a significant extension from that known previously for this subfamily. The paleobiogeographic implications of the presence of plagiosauroids in Western Gondwana have already been discussed by Dias-da-Silva and Ilha (2009) and repetition is unnecessary. Even so, depositional units from South America are providing new information regarding the distribution of temnospondyls across the Permo-Triassic boundary, mainly because of an increase in prospecting efforts, particularly in the last ten years. In conclusion, field work in South American units from Argentina, Uruguay, and Brazil must be an ongoing task in order to find more complete and diagnostic temnospondyls in this region of Gondwana.

Acknowledgements.—SDS would like to thank Ana Maria Ribeiro and Jorge Ferigolo (both Fundação Zoobotânica do Rio Grande do Sul,

Brazil) for allowing the study of MCN PV 1999. Rainer Schoch and Stéphanie Gastou (both Staatliches Museum für Naturkunde Stuttgart, Stuttgart, Germany) for the photographs of comparative plagiosternine material deposited in German collections. We are greatly indebted to the peer reviewers Claudia Marsicano (Universidad de Buenos Aires, Buenos Aires, Argentina), Dhurjati Sengupta (Indian Statistical Institute, Kolkata, India) and Florian Witzmann (Humboldt University, Berlin, Germany) for their suggestions that greatly improved the manuscript. The Conselho Nacional de Desenvolvimento Científico e Tecnológico (Brazil) provided the financial support (CNPq Research Grant 500919/2009-7).

References

- Abel, O. 1919. *Die Stämme der Wirbeltiere*. 914 pp. De Gruyter, Berlin.
- Damiani, R., Schoch, R.R., Hellrung, H., Werneburg, R., and Gastou, S. 2009. The plagiosaurid temnospondyl *Plagiosuchus pustuliferus* (Amphibia: Temnospondyli) from the Middle Triassic of Germany: anatomy and functional morphology of the skull. *Zoological Journal of the Linnean Society* 155: 348–373. <http://dx.doi.org/10.1111/j.1096-3642.2008.00444.x>
- Dias-da-Silva, S. and Ilha, A.L.R. 2009. On the presence of a pustulated temnospondyl in the Lower Triassic of southern Brazil. *Acta Palaeontologica Polonica* 54: 609–614. <http://dx.doi.org/10.4202/app.2008.0073>
- Fraas, E. 1889. Die Labyrinthodonten der Schwäbischen Trias. *Palaeontographica* 36: 1–158.
- Gastou, S. 2008. A Revision of the Plagiosaurid *Plagiosternum granulosum* Fraas (Amphibia: Temnospondyli) from the Middle Triassic of Germany. 76 pp. Unpublished DES Report, Université Paris 6, Paris.
- Hellrung, H. 2003. *Gerrothorax pustuloglomeratus*, ein Temnospondyle (Amphibia) mit knöcherner Branchialkammer aus dem Unteren Keuper von Kupferzell (Süddeutschland). *Stuttgarter Beiträge zur Naturkunde Serie B (Geologie und Paläontologie)* 330: 1–130.
- Schoch, R.R. and Werneburg, R. 1998. The Triassic labyrinthodonts from Germany. *Zentralblatt für Geologie und Paläontologie Teil I* 1998: 629–650.
- Shishkin, M.A. [Šiškin, M.A.] 1967. Plagiosaurs from the Triassic of the USSR [in Russian]. *Paleontologičeskij žurnal* 1967: 86–92.
- Shishkin, M.A. 1986. New data on plagiosaurs from the Triassic of the USSR [in Russian]. *Bulleten' Moskovskogo Obščestva Ispytatelej Prirody, Otdel Geologičeskij* 61: 97–102.
- Shishkin, M.A. 1987. Evolution of early amphibians (Plagiosauroida) [in Russian]. *Trudy Paleontologičeskogo Instituta* 225: 1–143.
- Warren, A. 1995. *Plagiosternum granulosum* E. Fraas: a plagiosaurid temnospondyl from the Middle Triassic of Crailsheim, Germany. *Stuttgarter Beiträge zur Naturkunde Ser. B* 229: 1–8.
- Warren, A. 2000. Secondarily aquatic temnospondyls of the Upper Permian and Mesozoic. In: H. Heatwole and R.L. Carroll (eds.), *Amphibian Biology, Vol. 4, Palaeontology: The Evolutionary History of Amphibians*, 1121–1149. Surrey Beatty and Sons, Chipping Norton.
- Yates, A.M. and Warren, A.A. 2000. The phylogeny of the “higher” temnospondyls (Vertebrata: Choanata) and its implications for the monophyly and origins of the Stereospondyls. *Zoological Journal of the Linnean Society* 128: 77–121. <http://dx.doi.org/10.1111/j.1096-3642.2000.tb00650.x>
- Zittel, K. von 1887–1890. *Handbuch der Paläontologie. I. Abteilung, Paläozoologie. 3. Vertebrata (Pisces, Amphibia, Reptilia)*. 1890 pp. Verlag R. Oldenbourg, München.

Sérgio Dias-da-Silva [sergiosilva@unipampa.edu.br], Laboratório de Paleobiologia, Campus de São Gabriel, Universidade Federal do Pampa, Avenida Antônio Trilha, 1847, Bairro Centro, CEP 97.300-000, São Gabriel, Rio Grande do Sul, Brazil;
Andrew R. Milner [andrew.milner@nhm.ac.uk], Department of Palaeontology, The Natural History Museum, Cromwell Road, London, SW7 5BD, United Kingdom.

Received 18 November 2009, accepted 18 March 2010, available online 25 March 2010.