

First Observations of Mating Behavior in Captive Kuroiwa's Ground Gecko (*Goniurosaurus kuroiwa*)

Authors: Yamazaki, Kei, Sasai, Takahide, Ashida, Hiroshi, and Kawazu, Isao

Source: Current Herpetology, 44(1) : 88-93

Published By: The Herpetological Society of Japan

URL: <https://doi.org/10.5358/hsj.44.88>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First Observations of Mating Behavior in Captive Kuroiwa's Ground Gecko (*Goniurosaurus kuroiwa*)

KEI YAMAZAKI^{1,*}, TAKAHIDE SASAI^{1,2}, HIROSHI ASHIDA¹, AND
ISAO KAWAZU^{1,2}

¹Okinawa Churaumi Aquarium, Ishikawa 424, Motobu, Okinawa 905–0206, JAPAN

²Okinawa Churashima Research Institute, Ishikawa 888, Motobu, Okinawa 905–0206, JAPAN

Abstract: Captive breeding programs are used worldwide to elucidate the natural history of endangered species and to recover and conserve their field populations. Kuroiwa's ground gecko, *Goniurosaurus kuroiwa* (Namiye, 1912) is an endangered species occurring in southern Japan; however, its mating behavior has not yet been described. We thus observed and quantified mating behavior of captive *G. kuroiwa*. Breeding trials with two males and three females resulted in 12 successful matings. Mating included precopulatory, copulatory, and postcopulatory stages, with 13 distinct mating behaviors observed. Behaviors indicative of successful copulation included the male preferentially biting the female's neck and the female approaching the male to initiate copulation. Mating behavior was initiated more than 10 h at night after cohabitation in some instances. These results suggest that at least one night of cohabitation is required for captive breeding of this species. Our findings thus contribute to conservation efforts by informing captive breeding programs.

Key words: Eublepharidae; Ex situ conservation; *Goniurosaurus kuroiwa*; Mating behavior

INTRODUCTION

Wild populations of Kuroiwa's ground gecko, *Goniurosaurus kuroiwa*, have declined because of habitat loss caused by development, predation by alien species such as mongoose, *Urva auropunctata*, and illegal collection (Toda and Tanaka, 2017). The species is currently listed as “vulnerable” on the IUCN Red List of Threatened Species (Kidera and Ota, 2017) and is in CITES Appendix III (CITES, 2023). Conservation guidelines advocate ex

situ methods such as captive breeding to supplement wild populations temporarily, which may be crucial for species preservation (Ministry of the Environment, Government of Japan, 2011). A thorough understanding of reproductive behaviors, including mating, is fundamental for the efficient implementation of captive breeding programs; however, no previous studies on the breeding behavior of *G. kuroiwa* exist. This species is nocturnal and solitary, complicating observing breeding behavior in situ. This study was conducted to observe and quantify the mating behaviors of *G. kuroiwa* in captivity to improve the understanding of its reproductive biology to inform and improve captive breeding techniques.

* Corresponding author.

E-mail address: k-yamazaki@okichura.jp

MATERIALS AND METHODS

Study animals

Five geckos (two males and three females) were captured at the Ocean Expo Park (Motobu, Okinawa Prefecture, Japan) in August and September 2020 and May 2021. Only sexually mature individuals (snout-vent length >77 mm) were used in this study (Tanaka and Nishihira, 1987) (Table 1). Each individual's sex was determined based on the presence or absence of hemipenal bulges (Kurita and Toda, 2013). All experiments were performed in accordance with the ethical guidelines for animal exhibition and research of the Japanese Association of Zoos and Aquariums (Japanese Association of Zoos and Aquariums, 2017).

Experimental conditions and behavioral observations

Each gecko was kept in a separate plastic container (21.5×36.5×15.0 cm) containing peat

moss, red ball soil, a plastic shelter, and a water dish. Rearing conditions were maintained at 18–31°C and >80% humidity; under a 10L:14D photoperiod. Every two days, geckos were fed crickets dusted with vitamin supplements. For mating trials, the geckos were placed in larger containers (34.0×43.0×28.0 cm) containing peat moss and red ball soil. For each trial, one male and female among the five captured individuals were randomly selected and were simultaneously placed together in a breeding container at 21.5–29.8°C. Mating behavior was observed between May and September. Four combinations of male-female pairs were tested during the trials (Table 1). Each trial lasted 16 hours, from 1800 to 1000 h, with 40 lx illumination. Behaviors were recorded using a digital video camera (Hero7 Black, GoPro Inc.) and were categorized into precopulatory, copulatory, and postcopulatory stages. In total, 15 trials were conducted.

RESULTS AND DISCUSSION

TABLE 1. Pairing of male and female Kuroiwa's ground geckos (*Goniurosaurus kuroiwa*) during mating trials. Black circles indicate the successful insertion of the male hemipenis into the female cloaca.

Date	Male ID	Female ID	Insertion of genitalia
11 May 2021	4	2	●
24 May 2021	5	3	●
29 May 2021	5	1	●
5 Jul 2021	4	1	●
21 Jul 2021	5	3	●
5 Aug 2021	4	2	●
15 Aug 2021	5	3	●
20 Aug 2021	4	2	●
22 Aug 2021	4	1	●
26 Aug 2021	5	3	●
7 Sep 2021	4	2	●
9 Sep 2021	5	3	●
12 Sep 2021	5	3	●
14 Sep 2021	5	1	●
19 Sep 2021	5	1	●

Insertion of the male genitalia into the female cloaca (defined as mating success) was observed in 12 (80%) of the 15 trials (Table 1; Fig. 1). The male and female precopulatory, copulatory, and postcopulatory behaviors are described in Fig. 1, Table 2; photographs of the behaviors are shown in Fig. 2.

The first step in precopulatory behavior is the sex identification of the mating partner. Previous studies indicate that in many gecko species, mating behavior is typically initiated by males (Brillet, 1993; Regalado, 2003; Todd, 2005). In *Eublepharis macularius*, *Hemidactylus mabouia*, and *Sphaerodactylus vincentii*, males lick females to obtain chemical signals for sex identification (Regalado, 2003; Marcum et al., 2008; Mason and Parker, 2010); however, in the present study, *G. kuroiwa* males licked females in only five cases (33%), suggesting the possibility that they recognize sex via other cues than non-volatile chemicals on the body surface of females. Additionally, males frequently initiated neck biting (67%; 10




FIG. 1. Mating behavioral sequence in Kuroiwa's ground gecko (*Goniurosaurus kuroiwa*) were categorized into precopulatory, copulatory, and postcopulatory patterns for both sexes. The behaviors are presented in the chronological order as displayed during mating.

cases) rather than tail biting, and neck biting was observed in all successful matings. This behavior, also observed in other geckos (e.g., *He. mabouia* and *Hoplodactylus maculatus*) (Regalado, 2003; Todd, 2005), has been described as a mechanism for female restraint during copulation. Therefore, neck biting by males likely contributes to successful copulation and mating success in *G. kuroiwa*. Furthermore, females approached males in five cases (33%), all of which resulted in successful copulation. This behavior, coupled with females raising their tails, would be a reliable indicator of female readiness for copulation. Although the time between introduction to the container and the onset of interaction varied in the present study (147–38, 117 s), mating was successful even in the cases where the outset of interaction was delayed. This suggests that if mating behavior does not begin immediately,

cohabitation at least overnight is required for captive breeding of *G. kuroiwa*.

Among copulatory behaviors, successful mating occurred in all cases that progressed to the “copulatory position.” Additionally, “gaping mouth” and “contortion” were observed, which, to our knowledge, have not been reported in other geckos. Among postcopulatory behaviors, licking of genitalia, as observed after copulation in the present study, may be a further indicator of ejaculation; similar behaviors have been observed in various gecko species (e.g., *E. macularius*, *Thecadactylus rapicauda*, *He. maculatus*, and *He. mabouia*) (Regalado, 2003; Todd, 2005; Quesnel, 2006; Huang, 2013). Further studies with larger sample sizes are needed to evaluate the interspecific differences in the reproductive behavior and the functional significance of licking of genitalia between this and the other gecko species.

TABLE 2. Definitions and quantitative data on mating behaviors observed in Kuroiwa's ground geckos (*Goniurosaurus kuroiwae*) presented in chronological order.

Behavior classification	Behavior type	Definition	Quantitative data (mean±SD; range, n)
Precopulatory behaviors	Female licks male	The female licks the male body surface before the male bites her.	6.8±9.8 times (1–24, n=5)
	Male licks female	The male licks the female body surface before biting her.	8.2±15.5 times (1–36, n=5)
	Male bites female, except on the neck	The male bites the lower trunk region of the female (e.g., lateral body, hind limb, or tail).	—
	Time from the start of cohabitation to the neck bite	—	2,619.5±4,162.5 s (387–12589 s, n=8)
	Male bites female's neck	Male bites female's neck while copulating.	—
	Time from the start of cohabitation to the neck bite	—	5,033.0±10,082.1 s (147–38117 s, n=14)
	Female raises tail	The female raises its tail in a copulatory position.	—
	Time from the start of the neck bite to the female raising tail	—	102.2±35.9 s (42–176 s, n=12)
	Copulatory position	After the female raises its tail, the male assumes a position where they contact each other's cloacal regions at the ventral side of the female.	—
	Insertion of genitalia (pausing)	The male inserts his genitalia into the female cloaca and pauses briefly.	—
Postcopulatory behaviors	Pausing	—	8.2±2.4 s (6–15 s, n=12)
	Lumbar thrust	The male thrusts its lumbar region after intromission.	4.4±1.3 s (2–6 s, n=12), 2.0±0.7 times/s (1.0–3.3 times/s, n=12)
	Gaping mouth	After the lumbar thrust, the male pauses and gapes its mouth, followed by a re-bite on the female's neck.	2.9±0.6 s (2–5 s, n=8)
	Contortion	The male shakes his lower trunk region very slightly after lumbar thrust.	23.7±2.9 s (18–28 s, n=11), 3.2±0.5 times/s (2.4–3.9 times/s, n=11)
	Time from insertion of the genitalia to the end of contortion	—	42.7±6.7 s (34–57 s, n=11)
	Separation of genitalia	Separation of male's hemipenis from the female's cloaca.	—
	Time from the end of contortion to separation	—	90.1±81.9 s (40–325 s, n=11)
	Releases bite	Terminating copulation, the male releases the female neck.	—
	Male bite	—	305.4±105.0 s (194–529 s, n=12)
	Male licks genitalia	Terminating copulation, the male licks its genitalia before retracting the hemipenis into the hemipenal bulges.	—




FIG. 2. Photographs depicting mating behaviors of Kuroiwa's ground gecko (*Goniurosaurus kuroiuae*): (A) male licks female, (B) male bites female, except on the neck, (C) male bites female neck and female raises its tail, (D) copulatory position, (E) gaping mouth, and (F) male licks its own genitalia.

This is the first report on the mating behavior of *G. kuroiuae* in captivity. The observed behavioral patterns and quantitative data across precopulatory, copulatory, and postcopulatory stages serve as indicators of mating success and provide valuable insights for informing captive breeding programs and improving the mating success rates. However, due to the small number of individuals used, these observations unlikely cover the entire repertoire of reproductive behavior of this species. Future studies with a larger number of individuals and populations are warranted.

ACKNOWLEDGMENTS

This study was approved by the Ministry of the Environment, the Government of Japan (reference no. 2007313 and 2104263 to K.Y., 2020–2021), the Okinawa Prefecture Board of Education (reference no. 36 and 20 to K.Y., 2020–2021) and the Okinawa General Secretariat (reference no. 34 and 16 to K.Y., 2020–2021). We thank Editage (www.editage.jp) for English language editing.

LITERATURE CITED

- BRILLET, C. 1993. Behavioural cues in sex recognition by two species of nocturnal lizards: *Eublepharis macularius* and *Paroedura pictus*. *Amphibia-Reptilia* 14: 71–82.
- CITES. 2023. Cites Appendices I, II and III. <https://cites.org/sites/default/files/eng/app/2023/E-Appendices-2023-11-25.pdf> (accessed 3 January 2024)
- HUANG, V. 2013. Ontogenetic and mechanistic explanations of within-sex behavioral variation in a lizard with temperature-dependent sex determination. *Unpublished doctoral dissertation*. University of Texas, Austin.
- JAPANESE ASSOCIATION OF ZOOS AND AQUARIUMS. 2017. *Ethical guidelines for animal exhibition and research*. <https://www.jaza.jp/assets/document/about-jaza/document/2016rinriyokou.pdf> (accessed 3 November 2024).
- KIDERA, N. AND OTA, H. 2017. *Goniurosaurus kuroiwaie*. The IUCN Red List of Threatened Species 2017: e.T98152257A96877452. <http://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T98152257A96877452.en> (accessed 20 April 2022)
- KURITA, T. AND TODA, M. 2013. Validation and application of skeletochronology for age determination of the Ryukyu ground gecko, *Goniurosaurus kuroiwaie* (Squamata: Eublepharidae). *Asian Herpetological Research* 4: 233–241.
- MARCUM, M. A., POWELL, M. A., MUENSCH, A. J., ARNOLD, H. F., AND POWELL, R. 2008. Social behaviour of the dwarf gecko *Sphaerodactylus vincenti vincenti* on St. Vincent, Lesser Antilles. *Salamandra* 44: 15–22.
- MASON, R. T. AND PARKER, M. R. 2010. Social behavior and pheromonal communication in reptiles. *Journal of Comparative Physiology A* 196: 729–749.
- MINISTRY OF THE ENVIRONMENT, GOVERNMENT OF JAPAN. 2011. *Basic policy on the reintroduction of endangered species of wild fauna and flora to the wild*. <https://www.env.go.jp/content/900519785.pdf> (accessed 7 May 2023)
- QUESNEL, V. C. 2006. Reproductive behaviour of the Neotropical gecko *Thecadactylus rapicauda* (Houttuyn). *Living World, Journal of the Trinidad and Tobago Field Naturalists' Club* 2006: 38–43.
- REGALADO, R. 2003. Roles of visual, acoustic, and chemical signals in social interactions of the Tropical house gecko (*Hemidactylus mabouia*). *Caribbean Journal of Science* 39: 307–320.
- TANAKA, S. AND NISHIHARA, M. 1987. A field study of seasonal, daily, and diel activity patterns of *Eublepharis kuroiwaie kuroiwaie*. *Herpetologica* 43: 482–489.
- TODA, M. AND TANAKA, S. 2017. *Goniurosaurus kuroiwaie kuroiwaie* (Namiye, 1912). p. 191–192. In: Department of Environmental Affairs, Okinawa Prefectural Government (ed.), *Threatened Wildlife in Okinawa, Third Edition (Animals)*. Red Data Okinawa. Department of Environmental Affairs, Okinawa Prefectural Government, Naha.
- TODD, A. C. 2005. The social mating system of *Hoplodactylus maculatus*. *New Zealand Journal of Zoology* 32: 251–262.

Accepted: 23 September 2024