

SOME BLOOD VALUES OF THE WESTERN DIAMOND-BACK RATTLESNAKE (*Crotalus atrox*) FROM SOUTH TEXAS

Authors: PARKER, O. S., and McCOY, R. H.

Source: Journal of Wildlife Diseases, 13(3) : 269-272

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-13.3.269>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SOME BLOOD VALUES OF THE WESTERN DIAMOND-BACK RATTLESNAKE (*Crotalus atrox*) FROM SOUTH TEXAS

O. S. PARKER, Department of Biology, Texas A & I University, Kingsville, Texas 78363, USA

R. H. McCOY, Box 278, Crowell, Texas 79227, USA

Abstract: Hematologic values of the western diamond-back rattlesnake (*Crotalus atrox*) from south Texas were determined by standard techniques. The means of the principal blood measurements were 0.68 ± 0.20 million erythrocytes/mm³, $31.9 \pm 4.6\%$ hematocrit and 10.0 ± 1.8 gm% hemoglobin. A comparison of hematologic values of snakes is given.

INTRODUCTION

Numerous researchers maintain colonies of venomous snakes as sources of enzymes.¹¹ Personnel involved in the care of snakes may encounter situations where disease compromises the health of specimens. Although present methods in reptile care incorporate quantitative hematology in the diagnosis of disease,^{7,11} few reports are available on the blood values of poisonous species. These data are presented to help establish the range of normal values for rattlesnake blood.

MATERIALS AND METHODS

Western diamond-back rattlesnakes (*Crotalus atrox*) were collected in the south Texas area around Duval County by participants in the Freer Jaycee Rattlesnake Roundup, 2-4 April 1976. Snakes 0.9 to 1.8 m long were decapitated and bled into sterile tubes. Sequester-Sol[†] (1 drop/5 ml blood) was added as the anticoagulant. Hemoglobin was measured as cyanmethemoglobin using Drabkin's

diluent.⁹ Samples in Drabkin's diluent were stored at 4 C until analyzed with a Beckman Model 25 spectrophotometer[‡] set at 540 nm. Cyanmethemoglobin Certified Standard[‡] was used to prepare the standard curve. Non-heparinized capillary tubes were filled with blood, sealed on one end with clay, and centrifuged for 5 min at 11,500 rpm in an Adams Microhematocrit centrifuge.[‡] A modified diluent of Natt and Herrick⁶ was used for erythrocyte counts. Gentian violet (0.1 gm/1) was substituted for methyl violet 2B in the diluent. Techniques for erythrocyte counts given by Seiverd⁹ were followed.

Blood was added to Drabkin's diluent within 10 min post-collection. Erythrocyte counts and hematocrits were performed within 4 hr post-collection.

Standard methods³ were used to calculate the mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC). Statistical analyses followed Mendenhall.⁵

[†] Cambridge Chemical Products, Inc., Fort Lauderdale, Fla. 33309.

[‡] Beckman Instruments, Inc., Anaheim, Cal. 92806.

[‡] Hycel Inc., Houston, Tex. 77001.

[‡] Clay-Adams, Inc., New York, N.Y. 10519.

RESULTS

The number of samples, range, mean, and standard deviation for erythrocyte (RBC) count, hematocrit (PCV), hemoglobin (Hb), MCV, MCH, and MCHC are listed in Table 1. The means of the measurements were: 0.68 ± 0.20 mil-

lion erythrocytes/mm³; $31.9 \pm 4.6\%$ hematocrit; 10.0 ± 1.8 gm% hemoglobin; 480 ± 152 μm^3 , MCV; 146 ± 46 pg, MCH; and $31.9 \pm 5.4\%$, MCHC. A comparison of the values reported for other snakes is found in Table 2.

TABLE 1. Number of samples (n), range, mean (\bar{X}), and standard deviation (S.D.) of blood values of the western diamond-back rattlesnake.

	n	range	\bar{X}	S.D.
RBC/mm ³ (X10 ⁶)	14	0.39-1.01	0.68	0.20
PCV %	71	20.0-43.0	31.9	4.6
Hb gm%	69	6.0-13.6	10.0	1.8
MCV μm^3	14	311-652	480	152
MCH pg	13	95-258	146	46
MCHC %	68	18.8-58.5	31.9	5.4

DISCUSSION

The mean hemoglobin concentration of rattlesnake blood is at the upper range of values reported for other snakes (Table 2). This could be due to species differences, environmental factors, or methods of measurement. Hemoglobin can be measured in several forms: oxyhemoglobin, carboxyhemoglobin, methemoglobin, and cyanmethemoglobin.⁵ The most accurate assay of total hemoglobin involves cyanmethemoglobin.⁴

Avian blood diluents have been used to count erythrocytes of reptiles.¹ Natt and Herrick⁶ used methyl violet 2B in an avian blood diluent that was stable

for long periods and simple to prepare. Methyl violet 2B was unavailable so gentian violet was substituted. This diluent was satisfactory for diluting rattlesnake blood as it prevented clumping and lysis, and allowed the differentiation of erythrocytes from leukocytes and thrombocytes.

Possible injury due to capture, stress induced by captivity and crowding, sex, and age of specimens were not considered in this study. Parasitic infestation as well as seasonal and dietary changes have been reported to produce wide fluctuations in hematological parameters of reptiles.⁷

Acknowledgements

The authors are grateful for the assistance offered by the Jaycees of Freer, Texas. D. Curling, T. Urban, J. Karges, T. Garza, and F. Chiles aided in the collection of blood samples. Our thanks are also extended to the faculty of the Department of Biology, Texas A&I University, for providing facilities and equipment.

TABLE 2. Comparison of mean hematologic values of snakes. Names in parentheses are those used in the reference. Values in parentheses are ranges.

species	RBC/mm ³ (X10 ⁶)	PCV %	Hb gm%	MCV μm ³	MCH pg	MCHC %	reference
<i>Crotalus atrox</i>	0.68 (0.39-1.01)	31.9 (20-43)	10.0 (6.0-13.6)	480 (311-652)	146 (95-258)	31.9 (18.8-58.5)	
<i>Constrictor c. imperator</i>	—	29.3 (22-37)	—	—	—	—	10
<i>Pituophis catenifer annectans</i>	—	29.5 (27-33)	—	—	—	—	10
<i>Pituophis melanoleucus sayi</i> (<i>P. sayi</i>)	1.095	—	8.8	—	80.5*	—	8
<i>Thamnophis sirtalis</i> (<i>Eutamia sirtalis</i>)	1.05 (0.71-1.39)	28.0 (19-37)	8.5 (5.8-11.3)	267 (266-268)	82.0	30.4*	2
<i>Heterodon platyrhinos</i> (<i>H. contortrix</i>)	0.57 (0.50-0.63)	18.7 (13-24)	5.6 (3.7-7.5)	325 (266-383)	95.5 (74-119)	29.9*	2
<i>Natrix sipedon</i>	0.77	35.5	10.0	465	131.0	28.2*	2
<i>Natrix natrix</i>	0.85	—	—	—	—	—	7

*Calculated from available data.

LITERATURE CITED

1. ALTLAND, P. D., B. HIGHMAN and B. WOOD. 1951. Some effects of x-irradiation on turtles. *J. Exptl. Zool.* 118: 1-14.
2. ——— and D. S. DITTMER (eds.). 1961. *Blood and Other Body Fluids*. Federation of American Societies for Experimental Biology. Bethesda, Md. 540 pp.
3. DAVIDSOHN, I. and B. B. WELLS. 1962. *Clinical Diagnosis by Laboratory Methods*. W. B. Saunders Co., Philadelphia, Penn. 1020 pp.
4. Hycel Inc. December, 1973. Cyanmethemoglobin reagent for the quantitative determination of hemoglobin in blood. Hycel publication No. 5717. Hycel Inc., Houston, Tex. 4 pp.
5. MENDENHALL, W. 1964. *Introduction to Statistics*. Wadsworth Publishing Co., Inc., Belmont, Cal. 305 pp.
6. NATT, M. P. and C. A. HERRICK. 1952. A new blood diluent for counting the erythrocytes and leucocytes of the chicken. *Poult. Sci.* 31: 735-738.
7. REICHENBACH-KLINKE, H. and E. ELKAN. 1965. *The Principal Diseases of Lower Vertebrates*. Academic Press, N.Y. 600 pp.
8. RYERSON, D. L. 1949. A preliminary survey of reptilian blood. *J. Ent. Zool.* 41: 49-55.
9. SEIVERD, C. E. 1964. *Hematology for Medical Technologists*. Lea & Febiger, Philadelphia, Penn. 643 pp.
10. THORSON, T. B. 1968. Body fluid partitioning in reptilia. *Copeia* 1968: 592-601.
11. WALLACH, J. D. 1969. Medical care of reptiles. *J. Am. vet. med. Ass.* 155: 1017-1034.

Received for publication 31 May 1976
