

CHRONIC DERMATITIS IN NUTRIA IN LOUISIANA

Authors: CHABRECK, ROBERT H., THOMPSON, ROGER B., and ENSMINGER, ALLAN B.

Source: Journal of Wildlife Diseases, 13(3) : 333-334

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-13.3.333>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

CHRONIC DERMATITIS IN NUTRIA IN LOUISIANA

ROBERT H. CHABRECK, Louisiana State University, School of Forestry and Wildlife Management, Baton Rouge, Louisiana 70803, USA

ROGER B. THOMPSON, Louisiana State University, School of Veterinary Medicine, Baton Rouge, Louisiana 70803, USA

ALLAN B. ENSMINGER, Louisiana Wildlife and Fisheries Commission, Refuge Division, New Orleans, Louisiana 70130, USA

Abstract: Chronic dermatitis in nutria (*Myocastor coypus*) in Louisiana was traced to secondary bacterial and fungal infection resulting from the penetration of achene awns of smooth beggartick (*Bidens laevis*) into the skin.

CASE REPORT

Reports from fur animal trappers in Louisiana of lesions on nutria (*Myocastor coypus*) resulting in or initiating damage to the pelts prompted an examination of infected animals. On 27 February 1974, a visit was made to the Salvador Wildlife Management Area (WMA) in southeastern Louisiana, origin of several of the reports. The area is situated 32 km southwest of New Orleans in a freshwater marsh and contains a high population of these rodents. Numerous nutria were observed on small resting platforms scattered throughout the marsh and 5 animals were collected. Two nutria contained lesions on the ventral portion of the body, in the neck and chest regions, and upper portion of the front legs.

A closer examination disclosed loss of epidermis in many areas of the skin as a result of necrosis. In affected areas, the surface was covered by necrotic debris containing numerous inflammatory cells along with colonies of bacteria and groups of fungal hyphae. The skin was extremely thickened because of proliferation of connective tissue in the dermis and the infiltration of inflammatory cells. Numerous thorn-like structures containing small barbs also were present and had penetrated deeply into the dermis. The structures were surrounded by neutrophils and appeared to

be the initial source of lesions with bacterial and fungal infections as secondary sources.

The barbed thorn-like structures were identified as awns of the achenes of smooth beggartick (*Bidens laevis*). Several complete achenes were found on the nutria with awns attached and penetrating the skin. Awns are usually paired, each containing 2 rows of retrorse barbs, and serve as a means of attachment to animals for achene dissemination. Smooth beggartick is a common plant on Salvador WMA and dies back during late fall, after the achenes have matured. The marsh usually becomes inundated during the winter and nutria collect dead stems of this plant for use in construction of resting platforms. Many mature achenes remain attached to the dead stems, come in direct contact with nutria on the platforms, and initiate the chronic dermatitis.

Nutria are extremely sensitive to foreign bodies and conditions similar to those reported herein were noted by Evans *et al.*² during tagging and marking studies. Tags, collars, or other devices attached to the animals soon produced infection.

Trappers received an average price of \$4.50 for dried nutria pelts during the 1974-75 season.³ Lesions caused by the awns weaken nutria skin, often causing it to tear during the skinning process. If

skins are successfully removed from affected animals, the lesions often occur as holes in the dried pelt, reducing the value as much as 75%.

Smooth beggartick is restricted to freshwater marshes in Louisiana and in only a few locations do sizable communities of the plant occur. Studies in 1968 disclosed that this species made up less than 0.1% of the total species composition of the freshwater marsh.¹ However, in most areas where dense stands occur, chronic dermatitis in nutria has been reported and complaints are becoming more numerous each year.

Smooth beggartick is a pioneer plant and will invade sites where succession has retrogressed. The dominant species

in the freshwater marshes is maidencane (*Panicum hemitomon*);¹ however, large die-offs of maidencane, as a result of excessive flooding, have taken place since 1973. Should smooth beggartick invade the areas formerly occupied by maidencane or other species, the problem of chronic dermatitis in nutria would increase. Habitat potentially suitable for smooth beggartick in southeastern Louisiana includes about 160,000 ha with an annual nutria harvest of approximately 250,000.

Future studies are needed to determine the present distribution of smooth beggartick and the factors affecting local abundance. Also, methods of control of the plant should be investigated.

LITERATURE CITED

1. CHABRECK, R. H. 1972. Vegetation, water and soil characteristics of the Louisiana coastal region. La. Agr. Exp. Sta. Bull. 664. 72 p.
2. EVANS, J., J. O. ELLIS, R. D. NASS and O. L. WARD. 1972. Techniques for capturing, handling, and marking nutria. Proc. 25th Ann. Conf. of the Southeast. Assoc. of Game and Fish Commissioners. p. 295-315.
3. O'NEIL, T. and G. LINScombe. 1975. The fur animals, the alligator, and the fur industry in Louisiana. La. Wildlife and Fisheries Comm., Wildlife Education Bull. 106. 66 p.

Received for publication 12 July 1976