

GASTRO-INTESTINAL HELMINTHS IN WHITE-TAILED DEER (*Odocoileus virginianus*) OF ILLINOIS

Authors: COOK, T. W., RIDGEWAY, B. T., ANDREWS, R., and HODGE, J.

Source: Journal of Wildlife Diseases, 15(3) : 405-408

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-15.3.405>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

GASTRO-INTESTINAL HELMINTHS IN WHITE-TAILED DEER (*Odocoileus virginianus*) OF ILLINOIS

T. W. COOK, □ B. T. RIDGEWAY, R. ANDREWS and J. HODGE, Department of Zoology, Eastern Illinois University, Charleston, Illinois 61920, USA.

Abstract: Two deer populations, one in northern Illinois the other in southern Illinois, were examined by necropsy (n = 44 and 40 respectively) for helminth parasites of the gastro-intestinal tract and abdominal cavity. Both herds were parasitized by *Apteragia odocoilei*, *Haemonchus contortus*, *Gongylonema pulchrum*, *Setaria yehi*, *Trichuris ovis*, and *Moniezia benedeni*. *Nematodirus* sp. was found only in deer of northern Illinois. *Ostertagia mossi*, *Capillaria* sp., *Cooperia* sp., and *Oesophagostomum* sp. were found only in deer of southern Illinois.

INTRODUCTION

White-tailed deer (*Odocoileus virginianus*) have recovered to harvestable levels within the last 20 years in Illinois. Since no published reports of helminth fauna of deer are available for Illinois, a survey of the gastro-intestinal and abdominal helminths is clearly warranted. This is a report of the helminths recovered at necropsy from deer located in two northern and five southern Illinois counties.

MATERIALS AND METHODS

White-tailed deer killed by hunters were brought to check stations in Carroll and Jo Daviess counties in northern Illinois and in Hardin, Johnson, Massac, Pope, and Saline counties in southern Illinois during November and December, 1971.

The gastro-intestinal tract was removed and divided into eight sections for study: esophagus, rumen, reticulum, omasum, abomasum, small intestine, cecum and large intestine. The surface of viscera also was examined at this time for *Setaria*.

The contents and scrapings from the mucosal linings were preserved in 10% formalin. This material was then placed into shallow black pans; helminths were located with the aid of a dissecting microscope. Nematodes were preserved in 10% formalin, later cleared in lactophenol, and identified using keys of Becklund and Walker,^{2,3,4} Levine,⁵ Skryabin *et al.*,¹⁰ and Yamaguti.¹³ Cestodes were fixed in hot alcohol-acetic acid-formalin, and later stained, cleared, and mounted on slides for identification using keys of Wardle and McLeod¹¹ and Yamaguti.¹² Voucher specimens were sent to the United States National Helminthological Collection in Beltsville, Maryland (Nos. 74971-74980).

RESULTS AND DISCUSSION

A total of 10 species of nematodes and one species of cestode was recovered from deer herds in northern and southern Illinois (Table 1). Six species of nematodes were found in both populations. *Capillaria* sp., *Cooperia* sp., and *Oesophagostomum* sp. were found in deer in southern Illinois, while

□ Present address: Department of Biology, Centenary College of Louisiana, Shreveport, Louisiana 71104, USA.

TABLE 1. Gastro-intestinal and abdominal helminths recovered from deer in two northern and five southern counties in Illinois.

Helminth species	Site*	No. infected/ No. examined		%		Intensity			
		N	S	N	S	Range		Mean	
						N	S	N	S
NEMATODA									
<i>Gongylonema pulchrum</i>	E	4/34	18/39	18.8	46.3	1-3	1-19	1.5	4
<i>Aptergia odocoilei</i>	A	27/44	27/40	61.5	67.5	1-84	1-238	14	30
<i>Haemonchus contortus</i>	A	10/44	9/40	22.7	22.4	1-13	1-13	4	3
<i>Ostertagia mossi</i>	A	0/44	14/40	0	35	0	1-38	0	7
<i>Capillaria</i> sp.	SI	0/44	2/40	0	5	0	2	0	2
<i>Cooperia</i> sp.	SI	0/44	2/40	0	5	0	1	0	1
<i>Nematodirus</i> sp.	SI	11/44	0/40	25	0	1-52	0	8	0
<i>Trichuris</i> sp.	LI	2/44	2/40	4.5	5	1-5	1-2	3	1.5
<i>Oesophagostomum</i> sp.	C	0/44	1/40	0	2.5	0	1	0	1
<i>Setaria yehi</i>	AC	6/44	7/40	13.6	17.5	1-4	1-8	2	2
CESTODA									
<i>Moniezia benedeni</i>	SI	8/44	6/40	18.2	15	1	1-3	1	1.3

*A = abomasum, AC = abdominal cavity, C = cecum, E = esophagus, LI = large intestine, and SI = small intestine.
N = N. Ill. and S = S. Ill.

Nematodirus sp. was found only in deer in northern Illinois. Using the t test, there was no significant difference ($P < 0.05$) in prevalence between sexes in either deer herd.

The variety and prevalence of helminth species recovered were rather

typical of recent surveys.^{1,6,7,8,9} Although some of the genera identified (*Capillaria*, *Cooperia*, *Haemonchus*, *Oesophagostomum*, and *Trichuris*) also parasitize cattle, according to Prestwood, *et al.*,⁶ it is unlikely that these deer are acting as reservoir hosts for livestock.

Acknowledgements

Facilities for processing deer tracts were provided in southern Illinois by M.E. Mansfield and the Dixon Springs Agricultural Center and in northern Illinois by Forrest Loomis of the Illinois Department of Conservation. This study was supported, in part, by Federal Aid to Wildlife Restoration, Illinois Project W-63-R, Illinois Department of Conservation.

LITERATURE CITED

1. BEAUDOIN, R.L., W.M. SAMUEL and C.P.A. STROME. 1970. Comparative study of the parasites of populations of white-tailed deer. *J. Wildl. Dis.* 6: 56-63.
2. BECKLUND, W.W. and M.L. WALKER. 1967. Redescrptions of the nematodes *Ostertagia bisonis* Chapin, 1925, of cattle and wild ruminants, and *Ostertagia mossi* Dikmans, 1931, of deer. *J. Parasit.* 53: 1273-1280.
3. ———. 1968. *Ostertagia dikmansii* sp. n. (Nematoda: Trichostrongylidae) from deer, *Odocoileus virginianus*, with a key to the species of medium stomach worms of *Odocoileus* in North America. *J. Parasit.* 54: 441-444.
4. ———. 1969. Taxonomy, hosts, and geographic distribution of the *Setaria* (Nematoda: Filaroidea) in the United States and Canada. *J. Parasit.* 55: 359-368.
5. LEVINE, N.D. 1968. *Nematode Parasites of Domestic Animals and of Man*. Burgess Publishing Co., Minneapolis, Minn. 600 pp.
6. PRESTWOOD, A.K., F.E. KELLOG, S.R. PURSGLOVE and F.A. HAYES. 1975. Helminth parasitisms among intermingling insular populations of white-tailed deer, feral cattle, and feral swine. *J. Am. vet. med. Ass.* 166: 787-789.
7. SAMUEL, W.M. 1969. Parasites of white-tailed deer (*Odocoileus virginianus*) in south Texas. Ph.D. Dissertation. University of Wisconsin. 196 pp.
8. ——— and R.L. BEAUDOIN. 1966. Evaluation of two survey methods for detection of helminth infections in white-tailed deer (*Odocoileus virginianus*). *Bull. Wildl. Dis. Ass.* 2: 100-107.
9. ——— and D.O. TRAINER. 1969. A technique for survey of some helminth and protozoan infections of white-tailed deer. *J. Wildl. Manage.* 33: 888-894.
10. SKRYABIN, K.I., N.P. SHIKHOBALOVA and R.S. SCHULZ. 1954. *Essentials of Nematodology. Trichostrongylids of Animals and Man*. Academy of Sciences of the USSR, Moscow. 704 pp.
11. WARDLE, R.A. and J.A. McLEOD. 1952. *The Zoology of Tapeworms*. University of Minnesota Press, Minneapolis. 780 pp.
12. YAMAGUTI, S. 1959. *Systema Helminthum*. Vol. II. *The Cestodes of Vertebrates*. Interscience Publ., New York. 860 pp.

-
13. ———. 1961. *Systema Helminthum*. Vol. III. *The Nematodes of Vertebrates*. Interscience Publ., New York. 1261 pp.

Received for publication 19 April 1978
