

BOOK REVIEW

Source: Bulletin of the Wildlife Disease Association, 2(3): 51-52

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-2.3.51

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Pemberton, R. T. 1959. Life cycle of Cyathostoma lari Blanchard 1849 (Nematoda, Strongyloidea). Nature 184, 1423.

Ressler, N. and Zak, B. 1956. Electrophoresis in a fluid film. Clin. chim Acta I, 392-99.

Sanders, E., Huddleson, I. F. and Schiabble, P. J. 1944. An electrophoretic study of serum and plasma from normal and leucosis affected chickens. J. biol. Chem. 155, 469-78.

Stoll, N. 1929. Studies with the strongyloid nematode, Haemonchus contorta. 1. Acquired resistance of hosts under natural re-infection conditions out of doors. Am. J. Hyg. 10, 384-418.

Sturkle, P. D. 1954. Avian Physiology. Comstock Publ. Assoc., Ithaca, New York. Threlfall, W. 1965. Studies on Helminth Parasites of Herring Gulls. Ph.D. Thesis, Univ. of Wales.

Twisselmann, N. M. 1939. A study of the cell content of blood of normal chickens with supra vital dyes. Poult. Sci. 30, 240-50.

White, J. C., Beaver, G. H., and Ellis, M. 1956. The analysis of human haemoglobins by paper electrophoresis. CIBA Foundation Symposium: Paper Electrophoresis. Churchill, London.

Wickware, A. B. 1947. The differential blood picture in chickens before and after administration of embryonated eggs of *Heterakis gallinae* with notes on pathogenicity. Can. J. comp. Med. II, 78-83.

Wild, A. E. 1963. Protein of Rabbit foetal fluids. Ph.D thesis. U.C.N.W., Bangor Yakimoff, W. L. and Rastegiaeff, 1929. Sur la question des variations cytologiques du sang des poules. Bull. Soc. Path. exot. 22, 766-69.

BOOK REVIEW

MYXOMATOSIS, by Frank Fenner and F. N. Ratcliffe. Cambridge University Press, London, New York and Ibadan. 1965. 379 pp.

Myxomatosis is a disease caused by a myxoma virus. It was first recognized in a laboratory rabbit in South America in 1898 and since then has been reported to occur sporadically in laboratory rabbits in several areas in the southern half of the western hemisphere. This virus is remarkably host specific, causing a mild infection in Sylvilagus spp. sometimes evidenced by a benign, localized, often persistent tumor in the skin, but no apparent mortality. In Oryctolagus spp., the European rabbit that has been used extensively for laboratory and commercial purposes throughout the world, the myxoma virus causes a much more extensive infection, usually fatal. No other susceptible hosts are known.

Myxomatosis has been reported in commercial rabbitries in California and Oregon since 1930. In 1954 myxomatosis was deliberately introduced into wild Oryctolagus in Tierra del Fuega and has since caused extensive outbreaks there and further north in Chile. The major impact of myxomatosis has occurred (a) in Australia where it was deliberately introduced in 1950 to control the established wild population of the European rabbit and (b) in Europe where it was introduced on a private estate in France in 1952 to control a local rabbit population but has since spread over much of Europe and the British Isles.

(Continued on page 77)

- CASSIE, R. M. 1954. Some uses of probability paper in the analysis of size frequency distributions. Australian J. Freshwater Res. 5: 514-517.
- DAVIS, D. E. 1959. The sex and age structure of roosting starlings. Ecology 40: 136-139.
- DAVIS, D. E. and C. ZIPPIN. 1954. Planning wildlife experiments involving percentages. J. Wild. Man. 18: 170-178.
- ERICKSON, H. R. 1963. Reproduction, growth, and movement of muskrats inhabiting small water areas in New York state. New York Fish and Game J. 10: 90-117.
- HILL, A. B. 1950. Principles of Medical Statistics. The Lancet Limited, London, 5th ed ix \pm 282 pp.
- HUNTER, W. S. and T. L. QUAY, 1953. An ecological study of the helminth fauna of Macgillivray's seaside sparrow (Ammospiza maritima macgilli vraii) (Audubon). Amer. Mid. Nat. 50: 407-413.
- KNIGHT, I. M. 1951. Diseases and parasites of the muskrat (Ondatra zibethica) in British Columbia. Can. J. Zool. 29: 188-214.
- LORD, R. D., Jr. 1959. The lens as an indicator of age in cottontail rabbits. J. Wild. Man. 23: 358-361.
- MEYER, M. C. and J. R. REILLY. 1950. Parasites of muskrats in Maine. Amer. Mid. Nat. 44: 467-477.
- OLIVER, L. 1962. Studies on natural resistance to Taenia taeniaeformis in mice. II. The effect of cortisone. J. Parasitol. 48: 758-762.
- PEARL, R. 1940. Introduction to Medical Biometry and Statistics. W. B. Saunders Co., Philadelphia. 3d ed. xv +537 pp.
- TAKOS, M. J. 1940. A review of literature on disease and parasites of the muskrat, Maine Cooperative Wildlife Research Unit. 14 pp.

BOOK REVIEW CONTINUED FROM PAGE 51-

Fenner and Ratcliffe are well qualified to write this volume because of their close association with the Australian studies. Fenner is Professor of Microbiology at the Australian National University and Ratcliffe was formerly Officer-in-charge of the Wildlife Survey Section of the Australian Commonwealth Scientific and Industrial Research Organization. They and their staffs were responsible for most of the research on myxonatosis in Australia. The history of myxomatosis during the past decade and a half establishes a classic of the development of a disease agent as a control mechanism of a wild animal population. The first part of the book outlines the history of the spread of the wild European rabbit in Australia and presents a detailed account of recent research on the biology of the wild rabbit. This is followed

by a technical description of the myxoma virus, its transmission by mosquitoes and other insects, the course of the disease in the European rabbit, and subsequent changes in the virulence of the virus and in the resistance of the rabbits to it. The final chapters document the history of the disease in the four continents where it occurs, and discuss its continuing evolution.

The authors have done an excellent job of telling this story in a concise and interesting fashion. It is well documented with a complete bibliography and a helpful index. Illustrations, charts and tables increase the value of the book. The reader's interest is enhanced by inclusion of photographs of many of the personalities who were involved in the research and operations that provided the data for this monumental work. Carleton M. Herman.