

SERUM CHEMISTRY OF THE COLLARED PECCARY (TAYASSU TAJACU)

Authors: Lochmiller, Robert L., and Grant, W. E.

Source: Journal of Wildlife Diseases, 20(2) : 134-140

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-20.2.134>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SERUM CHEMISTRY OF THE COLLARED PECCARY (*TAYASSU TAJACU*)

Robert L. Lochmiller and W. E. Grant

Department of Wildlife and Fisheries Sciences, Texas A&M University,
College Station, Texas 77843, USA

ABSTRACT: Values of serum biochemistry were obtained for 33 adult (16 male and 17 female) and six juvenile (four male and two female) collared peccaries collected by trapping and drugging animals from southern Texas during the period July through September 1982. Only cholesterol and aspartate aminotransferase concentrations differed significantly with respect to sex. Juvenile peccaries had significantly lower concentrations of total protein and globulins, but had higher concentrations of glucose, triglycerides, alkaline phosphatase, alanine aminotransferase, and calcium, and a higher albumin/globulin ratio than did adults. Effects of method of capture on biochemical attributes of serum from five gunshot and nine trapped adult peccaries collected from a single herd during March 1983 also were examined. Trapped peccaries had significantly higher levels of serum urea nitrogen, urea nitrogen/creatinine ratio, glucose, and chloride concentrations. Levels of uric acid, calcium, and potassium were significantly lower among the trapped animals.

INTRODUCTION

Serum biochemistry may be useful in diagnosing health and condition of populations (Bandy et al., 1957; Franzmann, 1972; Seal et al., 1972, 1978; Buckland, 1974; LeResche et al., 1974; Bahnak et al., 1979; Warren et al., 1981, 1982). Normal physiological values of serum metabolites have been reported for most North American ungulates, but information is practically nonexistent for the collared peccary. Zervanos and Hadley (1973) studied the effects of dehydration on serum levels of urea nitrogen and electrolytes among captive peccaries. Wallach et al. (1971) presented values of nine biochemical properties of captive adult peccaries (2 male, 2 female) collected by a gunshot to the head.

The peccary occupies arid habitats of the southwestern United States where prolonged periods of drought are not uncommon. Nutritional stresses during these periods have had profound influences on herd recruitment (Low, 1970; Bissonette, 1982) and serum metabolic constituents

may prove to be useful indices of nutritional stress in the collared peccary. The objective of this research was to establish baseline data on the serum biochemistry of the peccary from southern Texas. Influences of sex and age on metabolic profiles were examined. Information also was obtained on the effects of capture method (gunshot versus immobilization with drugs) on adult serum attributes.

MATERIALS AND METHODS

Thirty-three adult (16 male and 17 female) and six juvenile (4 male and 2 female) collared peccaries were trapped from the Chaparrosa Ranch, Zavala County, Texas during July through September 1982. These peccaries were captured in modified Clover deer traps baited with corn. Traps were baited and set around 2100 hours and checked the following morning by 0700 hours. Peccaries caught in traps were immobilized prior to blood-sampling with a dosage of 20 mg/kg of ketamine hydrochloride (Ketaset, Bristol Laboratories, Syracuse, New York 13201, USA) administered intramuscularly by blow-gun syringe (Lochmiller and Grant, 1983). Nine additional adult peccaries were captured using the same method on the Chaparral Wildlife Management Area, Dimmit and LaSalle counties, Texas during March 1983. Six adult animals were collected concurrently from the same herd, through killing by gunshot, in

Received for publication 8 September 1983.

order to compare their serum biochemical profiles to those herd members trapped and subsequently drugged. These two techniques are the most frequently used collection methods for collared peccaries (Neal, 1959). Shot animals were killed with a single rifle-shot to the cervical area of the spine, and blood samples usually were obtained prior to death. All animals collected from the Chaparral Wildlife Management Area were mature adults.

Animals were sexed and classified as adult or juvenile based on tooth-replacement pattern (Kirkpatrick and Sowls, 1962) and tooth wear (Sowls, 1961). Animals lacking complete permanent dentition were considered juveniles. All blood samples consisted of 50 ml of blood drawn from the anterior vena cava using 10-ml B-D Vacutainer assemblies (Becton-Dickinson and Co., Rutherford, New Jersey 07070, USA) fitted with 21-gauge, 3.8-cm needles. Blood samples were placed on ice and fractions were separated by centrifugation within 4 hr of collection. Serum samples were stored at -20°C until analyzed.

Serum samples were analyzed as a batch using a Technicon SMAC biochemical analyzer (Technicon Instruments Corp., Tarrytown, New York 10591, USA) for concentrations of total protein, albumin, serum urea nitrogen, creatinine, total bilirubin, direct bilirubin, cholesterol, glucose, triglycerides, uric acid, alkaline phosphatase (ALP), gamma glutamyl transferase (GGT), aspartate aminotransferase (AST), alanine aminotransferase (ALT), lactate dehydrogenase (LD), calcium, sodium, chloride, inorganic phosphorus, and potassium. Concentration of globulins was estimated by subtracting the concentration of albumin from total protein. The albumin/globulin ratio (A/G) and serum urea nitrogen/creatinine ratio was calculated for each sample. Cortisol was assayed by radio-immunoassay using a kit provided by Radioassay Systems Laboratories, Inc. (Carson, California 90746, USA). Intraassay variability was 2.6%.

Blood data of the 30 collared peccaries collected from the Chaparral Ranch were compared by analysis of variance (Helwig and Council, 1979) with sex and age as the main effects. The effects of capture method on serum biochemistry was assessed for the 14 adult peccaries collected from the Chaparral Wildlife Management Area using analysis of variance with capture method as the main effect. Sexes were pooled in this analysis because the Chaparral Ranch data showed few sexual dimorphisms in serum biochemistry. Seasonal com-

parisons were not made since the two areas of collection differed drastically in habitat quality.

RESULTS AND DISCUSSION

Species comparisons

The serum biochemistry of collared peccaries had many similarities to domestic swine (*Sus scrofa*). Total protein (7.37 g/dl) and albumin (3.10 g/dl) concentrations for the adult peccary were in close agreement with respective values reported for domestic swine (Baetz and Mengeling, 1971). Values for serum urea nitrogen and creatinine were within the normal range of published values for both European wild hogs (*S. scrofa*) (Williamson and Pelton, 1975) and domestic swine (Tumbleson et al., 1969, 1972; Baetz and Mengeling, 1971). Urea nitrogen values in this study were below those reported for collared peccaries in Arizona (range 10.7–22.7 mg/dl) (Zervanos and Hadley, 1973) and four zoological park specimens (range 23.0–37.0 mg/dl) (Wallach et al., 1971).

The peccary appears to have lower concentrations of bilirubin (total and direct) than swine (Tumbleson et al., 1969) (Table 1). Wallach et al. (1971) reported total bilirubin levels of 0.0–0.9 mg/dl for the peccary. Values for serum glucose, cholesterol, and ALP presented by Wallach et al. (1971) were within the observed range of values in this study. However, uric acid levels (2.8–6.5 mg/dl) given by Wallach et al. (1971) appeared abnormally high for the collared peccary. Values in this study averaged only 0.192 ± 0.037 (SE) mg/dl.

Levels of LD in the peccary were extremely high in comparison to domestic swine (Tumbleson et al., 1969) (Table 1). Blood serum from a single feral hog trapped during the present study was found to contain a LD concentration of only 374 I.U./liter compared to the average of 2,617 I.U./liter for adult peccaries. The hog was handled in a manner

TABLE 1. Influence of sex and age on serum chemical values of collared peccaries from southern Texas (July through September 1982).

| Component assay | Adults | | | | Juveniles | | | |
|--|------------------|-------|----------------|-------|-----------------|-------|---------------|-------|
| | Females (n = 17) | | Males (n = 16) | | Females (n = 2) | | Males (n = 4) | |
| | Mean | SE | Mean | SE | Mean | SE | Mean | SE |
| Total protein (g/dl) ^a | 7.28 | 0.19 | 7.49 | 0.15 | 7.15 | 0.75 | 5.85 | 0.40 |
| Albumin (g/dl) | 3.05 | 0.06 | 3.17 | 0.07 | 3.10 | 0.20 | 3.02 | 0.14 |
| Globulin (g/dl) ^a | 4.22 | 0.18 | 4.32 | 0.15 | 4.05 | 0.55 | 2.82 | 0.28 |
| Albumin/globulin (ratio) ^a | 0.74 | 0.04 | 0.75 | 0.03 | 0.78 | 0.06 | 1.09 | 0.07 |
| Serum urea nitrogen (mg/dl) | 8.94 | 0.55 | 8.50 | 0.68 | 12.00 | 5.00 | 5.25 | 0.48 |
| Creatinine (mg/dl) | 1.52 | 0.06 | 1.71 | 0.12 | 1.85 | 0.65 | 1.08 | 0.11 |
| Serum urea nitrogen/creatinine (ratio) | 5.93 | 0.35 | 5.09 | 0.37 | 6.32 | 0.48 | 4.96 | 0.47 |
| Total bilirubin (mg/dl) | 0.106 | 0.005 | 0.106 | 0.006 | 0.100 | 0.000 | 0.100 | 0.000 |
| Direct bilirubin (mg/dl) | 0.023 | 0.106 | 0.044 | 0.013 | 0.0 | 0.0 | 0.0 | 0.0 |
| Cholesterol (mg/dl) ^a | 113 | 7 | 80 | 4 | 96 | 21 | 112 | 15 |
| Glucose (mg/dl) ^a | 160 | 12 | 174 | 12 | 243 | 33 | 198 | 15 |
| Triglycerides (mg/dl) ^a | 54 | 9 | 41 | 6 | 104 | 76 | 214 | 128 |
| Uric acid (mg/dl) | 0.182 | 0.059 | 0.212 | 0.058 | 0.350 | 0.250 | 0.075 | 0.025 |
| Alkaline phosphatase (I.U./liter) ^a | 51 | 5 | 54 | 5 | 106 | 44 | 143 | 10 |
| Gamma glutamyl transferase (I.U./liter) | 9.88 | 0.83 | 8.81 | 1.46 | 8.50 | 3.50 | 8.75 | 1.25 |
| Aspartate aminotransferase (I.U./liter) ^b | 53 | 13 | 31 | 3 | 96 | 40 | 29 | 4 |
| Alanine aminotransferase (I.U./liter) ^a | 33 | 5 | 25 | 3 | 55 | 15 | 46 | 18 |
| Lactate dehydrogenase (I.U./liter) | 2,970 | 397 | 2,242 | 107 | 2,620 | 480 | 1,765 | 134 |
| Calcium (mg/dl) ^a | 9.47 | 0.16 | 9.71 | 0.15 | 10.45 | 0.85 | 10.42 | 0.24 |
| Sodium (mmol/liter) | 151.8 | 1.9 | 150.5 | 1.7 | 154.0 | 6.0 | 145.5 | 3.8 |
| Chloride (mmol/liter) | 115.6 | 1.8 | 115.4 | 1.5 | 118.0 | 3.0 | 112.5 | 3.0 |
| Phosphorus (mg/dl) | 4.51 | 0.34 | 4.39 | 0.34 | 6.25 | 1.35 | 5.10 | 0.52 |
| Potassium (mmol/dl) | 3.72 | 0.09 | 3.67 | 0.10 | 4.15 | 0.75 | 3.90 | 0.28 |

^a Sex effect ($P < 0.05$).^b Sex effect ($P < 0.01$).^c Age effect ($P < 0.05$).^d Age effect ($P < 0.01$).^e Age effect ($P < 0.001$).

identical to that used with peccaries, and the hog's blood serum was assayed along with those of the peccaries. The peccary may have naturally higher levels of LD but this difference also may be related to the physiological stress of capture. According to Henry (1979, p. 365) a 2–40-fold elevation in LD levels can occur in those patients with severe shock. Preliminary analysis of data gathered from a group of captive peccaries has shown LD concentrations to decrease over time with repeated sampling, reaching values as low as 600 I.U./liter; thus, suggesting some degree of adaptation to handling stress occurred.

Levels of sodium and chloride were high among trapped peccaries in this study (Table 1), but appeared to be within the reported range for domestic swine (Tumbleson et al., 1972). Zervanos and Hadley (1973) demonstrated that sodium and chloride levels are elevated significantly when peccaries are maintained under dehydrated conditions. Slight dehydration might have occurred among trapped peccaries. Calcium and potassium levels in this study were similar to values reported by Zervanos and Hadley (1973) for collared peccaries on a prickly pear cactus diet (*Opuntia* spp.). Serum phosphorus levels were below those reported for domestic swine (Tumbleson et al., 1969). These lower levels of phosphorus may be attributable to diet quality. Varner et al. (1976) have reported summer phosphorus content of southern Texas prickly pear to be only 0.08% compared to an average of 0.20% phosphorus for eight forb species. Prickly pear is the primary food source of the collared peccary, often comprising 80 to 90% of the diet (Eddy, 1961; Everitt et al., 1981).

Sexual dimorphisms

There is general lack of sexual dimorphism in serum biochemical attributes in wild and domestic swine (Tumbleson et

al., 1969; Williamson and Pelton, 1975). A similar observation was made among collared peccaries. Only cholesterol and AST concentrations varied significantly between males and females trapped from the Chaparrosa Ranch (Table 1). Cholesterol and AST levels were highest among females. Female swine are also known to exhibit higher levels of cholesterol than males (Tumbleson et al., 1969). The serum urea nitrogen/creatinine ratio and LD concentration showed a slight, but nonsignificant ($P < 0.06$) difference between male and female peccaries.

Age differences

Age differences in serum biochemical attributes among the animals trapped from the Chaparrosa Ranch were more evident than sexual dimorphisms. Juvenile peccaries had significantly lower concentrations of total protein and globulins, but had a higher A/G ratio than did adults (Table 1). There was a significant interaction between sex and age for these three protein measures, indicating that juvenile males differed significantly from adult male and female peccaries. Tumbleson et al. (1969) reported that serum total protein concentrations increased from 4–9 mo of age to a level of 7.5 g/dl in swine. Jakobsen and Moustgaard (1950) reported that serum globulin levels of growing swine resemble those of the adult at between 5 and 6 mo of age.

Age effects also were noted for concentrations of glucose, triglycerides, ALP, ALT, and calcium (Table 1). Juvenile peccaries had significantly higher levels of these constituents than did adult peccaries. Tumbleson et al. (1969) reported that levels of serum glucose do not change with age in swine. The higher ALP levels probably are associated with osteoblastic activity in the growing juveniles (Coles, 1974). Likewise, the higher calcium levels probably resulted, in part, from skeletal and associated development in juveniles.

TABLE 2. Effect of two immobilization techniques on serum biochemistry of adult collared peccaries collected from a single herd from southern Texas in March 1983.

| Component assay | Capture method | | | |
|---|-----------------|-------|-----------------|-------|
| | Gunshot (n = 5) | | Drugged (n = 9) | |
| | Mean | SE | Mean | SE |
| Total protein (g/dl) | 8.26 | 0.32 | 8.77 | 0.26 |
| Ablumin (g/dl) | 3.38 | 0.21 | 3.53 | 0.14 |
| Globulin (g/dl) | 4.88 | 0.37 | 5.23 | 0.29 |
| Albumin/globulin (ratio) | 0.72 | 0.08 | 0.69 | 0.05 |
| Serum urea nitrogen (mg/dl) ^a | 6.80 | 1.32 | 18.33 | 2.44 |
| Creatinine (mg/dl) | 1.24 | 0.12 | 1.54 | 0.12 |
| Serum urea nitrogen/ creatinine (ratio) ^a | 5.30 | 0.65 | 11.69 | 1.26 |
| Total bilirubin (mg/dl) | 0.120 | 0.020 | 0.100 | 0.000 |
| Direct bilirubin (mg/dl) | 0.020 | 0.020 | 0.011 | 0.011 |
| Cholesterol (mg/dl) | 112 | 7 | 143 | 13 |
| Glucose (mg/dl) ^b | 144 | 18 | 197 | 11 |
| Triglycerides (mg/dl) | 39 | 13 | 25 | 3 |
| Uric acid (mg/dl) ^b | 0.500 | 0.104 | 0.244 | 0.044 |
| Alkaline phosphatase (I.U./liter) | 60 | 6 | 45 | 6 |
| Gamma glutamyl transferase (I.U./liter) | 4.20 | 1.24 | 5.44 | 0.82 |
| Aspartate aminotransferase (I.U./liter) | 31 | 6 | 51 | 22 |
| Alanine aminotransferase (I.U./liter) | 30 | 11 | 37 | 6 |
| Lactate dehydrogenase (I.U./liter) | 1,415 | 195 | 1,726 | 178 |
| Calcium (mg/dl) ^d | 12.30 | 0.35 | 10.44 | 0.22 |
| Sodium (mmol/liter) ^a | 151.8 | 2.7 | 157.2 | 1.7 |
| Chloride (mmol/liter) ^c | 110.0 | 0.8 | 120.7 | 2.0 |
| Phosphorus (mg/dl) | 6.42 | 0.80 | 5.85 | 0.60 |
| Potassium (mmol/liter) ^d | 8.16 | 0.24 | 3.52 | 0.16 |
| Cortisol (mg/dl) | 5.53 | 1.51 | 10.02 | 1.85 |

^a $P < 0.10$.^b $P < 0.05$.^c $P < 0.01$.^d $P < 0.001$.

Capture method effects

Method of capture was found to influence biochemical profiles of peccaries (Table 2). Concentration of serum urea nitrogen and the urea nitrogen/creatinine ratio were significantly higher among the trapped peccaries in comparison to those who were shot. Azotemia has been reported in stressed reindeer (*Rangifer tarandus*) (Hyvärinen et al., 1976; Rehbinde and Edgvist, 1981). The higher urea levels observed among trapped peccaries may have been due to a combination of state of hydration and circulating glucocorticoids which have a protein catabolic

effect on muscle. However, although cortisol levels among trapped peccaries averaged twice as high as shot animals, this difference was not significant. The lack of any significant differences in serum enzyme levels (i.e., AST) between trapped and shot peccaries would suggest that muscular trauma contributed little to the observed azotemia. In fact, the hyperkalemic serum of gunshot animals (Table 2) suggested that tissue trauma effects were considerably more pronounced with this capture method.

Serum levels of glucose were significantly higher, while levels of uric acid

were lower, among trapped peccaries. Glucose has been reported to increase initially with handling stress in bighorn sheep (*Ovis canadensis*) (Franzmann, 1972), white-tailed deer (*Odocoileus virginianus*) (Seal et al., 1972), and woodland caribou (*R. tarandus*) (Karns and Crichton, 1978). Concentrations of calcium were lower, and concentrations of chloride higher, among trapped peccaries than among gunshot animals. Sodium levels tended to be higher among trapped animals, but this difference was not significant. The higher chloride levels may have been due in part to slight dehydration. Elevated sodium levels have been demonstrated with handling stress on cattle (Gartner et al., 1965), reindeer (Hyvärinen et al., 1976), and woodland caribou (Karns and Crichton, 1978). Seal et al. (1972) did not find differences that could be attributed to handling stress in white-tailed deer calcium, potassium, or sodium concentrations.

The two areas which were sampled in this study varied considerably in habitat composition and rainfall received. Because rainfall has such a profound influence on forage nutritive quality in southern Texas, we did not attempt an analysis of seasonal influences on serum biochemistry. However, seasonal information should provide a better understanding of metabolic dynamics, hence, baseline blood values for this species in the wild.

ACKNOWLEDGMENTS

We are grateful for the field assistance provided by Dr. Pat Reardon and the rest of the staff at the Chaparrosa Ranch, La Pryor, Texas, and to Ernie Davis and the staff of the Chaparral Wildlife Management Area, Artesia Wells, Texas. We also thank Dr. Larry Varner for providing laboratory equipment and space. The field assistance given by P. J. Ettestad, P. W. Oldenberg, and E. C. Helgren also is appreciated. Financial support was provided by the National Rifle Association of America and the Caesar Kleberg Research Program in Wildlife Ecology, Department of Wildlife and Fisheries Sciences, Texas A&M University.

LITERATURE CITED

- BAETZ, A. L., AND W. L. MENGELING. 1971. Blood constituent changes in fasted swine. *Am. J. Vet. Res.* 32: 1491-1499.
- BAHNAK, B. R., J. C. HOLLAND, L. J. VERME, AND J. J. OZOGA. 1979. Seasonal and nutritional effects on serum nitrogen constituents in white-tailed deer. *J. Wildl. Manage.* 43: 454-460.
- BANDY, P. J., W. D. KITTS, A. J. WOOD, AND I. MCCOWAN. 1957. The effect of age and the plane of nutrition on the blood chemistry of the Columbian black-tailed deer (*Odocoileus hemionus columbianus*). *Can. J. Zool.* 35: 283-289.
- BISSETTE, J. A. 1982. Ecology and social behavior of the collared peccary in Big Bend National Park, Texas. U.S. Nat. Park Ser. Sci. Monogr. Ser. No. 16, 85 pp.
- BUCKLAND, D. E. 1974. Blood urea nitrogen levels of white-tailed deer as an index of condition and nutritional intake. M.S. Thesis. Virginia Polytechnic Institute and State Univ., Blacksburg, Virginia, 48 pp.
- COLES, E. H. 1974. Veterinary Clinical Pathology, 2nd ed. W. B. Saunders Co., Philadelphia, Pennsylvania, 615 pp.
- EDDY, J. A. 1961. Foods and feeding patterns of the collared peccary in southwest Arizona. *J. Wildl. Manage.* 25: 248-257.
- EVERITT, J. H., C. L. GONZALEZ, M. A. ALANIZ, AND G. V. LATIGO. 1981. Food habits of the collared peccary on south Texas rangelands. *J. Range Manage.* 34: 141-144.
- FRANZMANN, A. W. 1972. Environmental sources of variation of bighorn sheep physiological values. *J. Wildl. Manage.* 36: 924-932.
- GARTNER, R. J. W., J. W. RYLEY, AND A. W. BEATTIE. 1965. The influence of degree of excitation on certain blood constituents in beef cattle. *Aust. J. Exp. Biol. Med. Sci.* 43: 713-724.
- HELGREN, J. T., AND K. A. COUNCIL. 1979. SAS User's Guide. SAS Inst. Inc., Cary, North Carolina, 494 pp.
- HENRY, J. B. 1979. Clinical Diagnosis and Management, 16th ed. W. B. Saunders Co., Philadelphia, Pennsylvania, 1,170 pp.
- HYVÄRINEN, H., T. HELLE, M. NIEMINEN, Q. VÄYRYNEN, AND R. VÄYRYNEN. 1976. Some effects of handling reindeer during gatherings on the composition of their blood. *Anim. Prod.* 22: 105-114.
- JAKOBSEN, P. E., AND J. MOUSTGAARD. 1950. Investigation of the serum proteins in pigs from birth to maturity. *Nord. Veterinaarmed.* 2: 812-822.
- KARNS, P. D., AND V. F. J. CRICHTON. 1978. Effects of handling and physical restraint on blood parameters in woodland caribou. *J. Wildl. Manage.* 42: 904-908.

- KIRKPATRICK, R. D., AND L. K. SOWLS. 1962. Age determination of the collared peccary by the tooth-replacement pattern. *J. Wildl. Manage.* 26: 214-217.
- LERESCHE, R. E., U. S. SEAL, P. D. KARNS, AND A. W. FRANZMANN. 1974. A review of blood chemistry of moose and other Cervidae with emphasis on nutritional assessment. *Nat. Can. (Que.)* 101: 263-290.
- LOCHMILLER, R. L., AND W. E. GRANT. 1983. A sodium bicarbonate-acid powered blow-gun syringe for remote injection of wildlife. *J. Wildl. Dis.* 19: 48-51.
- LOW, W. A. 1970. The influence of aridity on reproduction of the collared peccary (*Dicotyles tajacu* (Linn.)) in Texas. Ph.D. Dissert. Univ. of British Columbia, Vancouver, British Columbia, 170 pp.
- NEAL, B. J. 1959. Techniques of trapping and tagging the collared peccary. *J. Wildl. Manage.* 23: 11-16.
- REHBINDER, C., AND L. EDGVIST. 1981. Influence of stress on some blood constituents in reindeer (*Rangifer tarandus* L.). *Acta Vet. Scand.* 22: 480-492.
- SEAL, U. S., L. J. VERME, AND J. J. OZOGA. 1978. Dietary protein and energy effects on deer fawn metabolic patterns. *J. Wildl. Manage.* 42: 776-790.
- , J. J. OZOGA, A. W. ERICKSON, AND L. J. VERME. 1972. Effects of immobilization on blood analysis of white-tailed deer. *J. Wildl. Manage.* 36: 1034-1040.
- SOWLS, L. K. 1961. Hunter-checking stations for collecting data on the collared peccary (*Pecari tajacu*). *Trans. N. Am. Wildl. Nat. Resour. Conf.* 26: 497-505.
- TUMBLESON, M. E., T. M. BADGER, P. C. BAKER, AND D. P. HUTCHESON. 1972. Systematic oscillations of serum biochemic and hematologic parameters in Sinclair (S-1) miniature swine. *J. Anim. Sci.* 35: 48-50.
- , C. C. MIDDLETON, O. W. TINSLEY, AND D. P. HUTCHESON. 1969. Serum biochemic and hematologic parameters of Hormel miniature swine from four to nine months of age. *Lab. Anim. Care* 19: 345-351.
- VARNER, L. W., L. H. BLANKENSHIP, AND G. W. LYNCH. 1976. Seasonal changes in nutritive value of deer food plants in south Texas. *Proc. Annu. Conf. Southeast. Assoc. Game Fish Comm.* 31: 99-106.
- WALLACH, J. D., W. C. RUSSELL, AND K. HERMAN. 1971. The normal hematology and serum chemistry of the collared peccary. *J. Zoo Anim. Med.* 2: 32.
- WARREN, R. J., R. L. KIRKPATRICK, A. OELSCHLAGER, P. F. SCANLON, K. E. WEBB, AND J. B. WHELAN. 1982. Energy, protein, and seasonal influences on white-tailed deer fawn nutritional indices. *J. Wildl. Manage.* 46: 302-312.
- , ———, ———, ———, AND F. G. GWAZDAUSKAS. 1981. Dietary and seasonal influences on nutritional indices of adult male white-tailed deer. *J. Wildl. Manage.* 45: 926-936.
- WILLIAMSON, M. J., AND M. R. PELTON. 1975. Some biochemical parameters of serum of European wild hogs. *Proc. Annu. Conf. Southeast. Assoc. Game Fish Comm.* 29: 672-679.
- ZERVANOS, S. M., AND N. F. HADLEY. 1973. Adaptational biology and energy relationships of the collared peccary (*Tayassu tajacu*). *Ecology* 54: 759-774.