

Skrjablingylus chitwoodorum Hill, 1939 (Nematoda: Metastrongyloidea) in Striped Skunks from Northcentral Minnesota

Authors: Fuller, Todd K., and Kuehn, David W.

Source: Journal of Wildlife Diseases, 20(4) : 348-350

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-20.4.348>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ments overlapped (Fig. 1). The maximum length of *P. boughtoni* (range = 380–500 μm) fell below the range of larvae of *Protostrongylus* spp. from bighorn sheep (527–673 μm), but overlapped with that of *O. macrotis* (488–611 μm). Also, maximum length of *O. macrotis* overlapped with the range of *Protostrongylus* spp. from bighorns. Therefore, where the three definitive hosts are sympatric, only infective larvae in the extremes of the total length range for all species can be identified with confidence. Larvae of *P. boughtoni* can be distinguished from those of *Protostrongylus* spp. from bighorns if there is no possibility of *Orthostrongylus* larvae occurring in a sample.

Although total length is not the most consistent morphologic basis on which to separate species of helminths, such a

method is attractive in its simplicity and adaptability to field studies. Third-stage larvae of some species in the confamilial genus *Parelaphostrongylus* differ in tail structure (Ballantyne and Samuel, 1984, J. Parasitol. In press). Specific structural differences may be found among *Protostrongylus* and *Orthostrongylus* infective larvae, but rapid examination of fine morphological details is prevented by the nearly opaque first-stage cuticle. Removal of this cuticle without damaging the larva inside, though not impossible, is extremely difficult and time-consuming.

We wish to thank Brent Gray and Judith Samson for assistance in exposing and maintaining snails. This study was supported by the Alberta Fish and Wildlife Division, and the Natural Sciences and Engineering Research Council of Canada.

Journal of Wildlife Diseases, 20(4), 1984, pp. 348–350
© Wildlife Disease Association 1984

***Skrjabinylus chitwoodorum* Hill, 1939 (Nematoda: Metastrongyloidea) in Striped Skunks from Northcentral Minnesota**

Todd K. Fuller and David W. Kuehn, Forest Wildlife Populations and Research Group, Minnesota Department of Natural Resources, Grand Rapids, Minnesota 55744, USA

The nematode *Skrjabinylus chitwoodorum* occurs in the frontal sinuses of striped skunks (*Mephitis mephitis* Schreber) and causes progressive deformation of the frontal region of the cranium. The frequency of lesions and degree of deformation were related to relative age and geographic distribution of skunks (Kirkland and Kirkland, 1983, Can. J. Zool. 61: 2913–2920). There are, however, no published accounts relating skull lesions to actual ages of striped skunks, nor is there documentation of how numbers of worms relate to age and degree of deformation.

This note presents age-specific documentation of prevalence of lesions and numbers of adult *S. chitwoodorum* in the frontal sinuses of striped skunks from a forested region in northcentral Minnesota.

Skunks were collected in the northeastern portion of Itasca County, Minnesota (47°52'N, 93°22'W). In Grand Rapids, 50 km southwest of the study area, mean January temperature is –14 C, and in July the mean is 19 C (U.S. Dept. Commerce, unpubl. data). Snow cover is usually present from December through early April, and during the winters of 1970–1971 through 1982, mean January–March snow depth was 46 cm. Total annual precipitation averages 29.9 cm.

Received for publication 25 April 1984.

TABLE 1. Age-specific cranial damage to striped skunks in northcentral Minnesota caused by *Skrjabinigylus chitwoodorum*.

Age (yr)	n	Cranial damage class				% With class 2 or 3 damage
		0	1	2	3	
0.5 ^a	21	19	2	0	0	0.0
1.0–1.5 ^{a,b}	24	2	18	2	2	16.7
2.0–2.5 ^b	26	1	9	12	4	61.5
≥3.0	17	0	4	7	6	76.5

^a Distribution within cranial damage classes (0 vs. 1–3) significantly different ($\chi^2 = 30.36$, 1 df, $P < 0.001$).

^b Distribution within cranial damage classes significantly different ($\chi^2 = 11.08$, 3 df, $P < 0.02$).

Vegetation in the study area is mostly boreal coniferous/deciduous forest (Maycock and Curtis, 1960, Ecol. Monogr. 31: 1–35). Uplands are dominated by aspen (*Populus* spp.), balsam fir (*Abies balsamea* L.), paper birch (*Betula papyrifera* Marshall), and jackpine (*Pinus banksiana* Lambert); black spruce (*Picea mariana* Miller), tamarack (*Larix laricina* DuRoi) and black ash (*Fraxinus nigra* Marshall) are common in lowland areas.

Skunks were captured incidentally in traps set for wolves (*Canis lupus* L.) during spring (April–May) and fall (August–September) 1981–1983. Body weights and measurements were recorded, and skulls, reproductive tracts and stomachs were collected. Skulls were boiled so that they could be measured and canine teeth extracted. Cranial damage was assigned to one of four categories, with class 0 indicating no noticeable damage and class 3 indicating gross deformation of the frontal region accompanied by perforations or erosion of the frontal bones (Kirkland and Kirkland, 1983, op. cit.). Since most skulls were boiled before cranial damage caused by *S. chitwoodorum* was recognized, an additional sample of skulls was examined before boiling. These skulls were dissected and nematodes from frontal sinuses were counted and measured. Canine teeth were radiographed and skunks <10 mo old

TABLE 2. Numbers of *Skrjabinigylus chitwoodorum* in frontal sinuses of six male striped skunks collected in northcentral Minnesota.

Age (yr)	Cranial damage class	Total no. of worms	% Male worms
0.5	1	25	48
2.0	2	97	45
2.0	2	111	50
3.0	2	116	59
3.5	1	61	57
3.5	3	381	64

identified (Fredrickson, 1983, Wildl. Soc. Bull. 11: 297–299). Older individuals were aged by counting cementum annuli (Casey and Webster, 1975, Can. J. Zool. 53: 223–226).

A sample of *S. chitwoodorum* was measured to confirm species identification (Lankester, 1983, Can. J. Zool. 61: 2168–2178). Total length of female worms averaged (\pm SD) 36.4 ± 5.1 mm (range = 28–48 mm, $n = 27$) vs. 19.4 ± 3.4 mm (range = 13–26, $n = 23$) for males. Mean spicule length was 763 ± 50 μ m (range = 690–815, $n = 7$). Both male and female worms averaged larger than those described originally by Hill (1939, J. Parasitol. 25: 475–478) in Oklahoma, were similar to worms from skunks in Illinois (Levine et al., 1962, Ill. Acad. Sci. Trans. 55: 3–5) and New York (Goble, 1942, J. Mammal. 23: 96–97), and smaller than worms found in Ontario (Lankester, 1983, op. cit.). Representative specimens have been deposited in the U.S. National Parasite Collection (Beltsville, Maryland 20705, USA) (Accession No. 78245).

A total of 88 skulls was examined (46 male, 42 female), seven of which (all males) were dissected to collect whole worms. There were no statistical differences ($P > 0.16$) between the sexes in age-specific cranial damage, but damage did increase with age (Table 1). Only 10% of 21 skunks 0.5 yr old had noticeable cranial damage due to *S. chitwoodorum*, but all

17 skunks ≥ 3.0 yr old had damage, and most was moderate to severe.

The total number of worms extracted from skulls of six male skunks was correlated significantly with cranial damage class ($r^2 = 0.80$, 4 df, $P < 0.05$) (Table 2). Older skunks tended to have a higher proportion of male worms (weighted $r^2 = 0.64$, 4 df, $P = 0.06$).

Increased numbers of worms and more severe cranial damage were associated with older skunks, probably resulting from the progressive accumulation of worms over time (Kirkland and Kirkland, 1983, op. cit.). Kirkland and Kirkland (1983, op. cit.) found an average of 70–75% of skulls of adult skunks from across North America exhibited lesions (vs. 25–30% of non-adults), and that rates of damage were highest in regions of high precipitation.

These phenomena may be related to the mode of transmission of *Skrjabinogylus* spp., since terrestrial gastropods are an obligate intermediate host whose abundance and distribution varies directly with environmental moisture.

This study was supported by the Forest Wildlife Populations and Research Group, Minnesota Department of Natural Resources and the U.S. Fish and Wildlife Service, Office of Endangered Species. We are grateful to R. K. Markl for field assistance, P. L. Coy for determining skunk ages, M. J. Pybus and M. W. Lankester for providing definitive identification of the nematodes, and D. Heisey, M. K. Lankester, M. J. Pybus, G. L. Kirkland and P. D. Karns for reviewing the manuscript.

Journal of Wildlife Diseases, 20(4), 1984, pp. 350–351
© Wildlife Disease Association 1984

***Renicola lari* Timon-David, 1933 from the Osprey, *Pandion haliaetus* (L.), from Alberta, Canada**

Murray J. Kennedy, Alberta Agriculture, Veterinary Services Branch, 6909 - 116 Street, Edmonton, Alberta T6H 4P2, Canada; and **Paul F. Frelser**, Alberta Agriculture, Veterinary Services Branch, Airdrie, Alberta T0M 0B0, Canada

Adult Digenea belonging to the genus *Renicola* parasitize the kidneys of birds frequenting marine or brackish waters (Martin, 1982, Proc. Helminthol. Soc. Wash. 49: 19–21). Life cycles have been described by Stunkard (1964, Biol. Bull. 126: 467–489), Werding (1969, Mar. Biol. 3: 306–333), Martin (1971, Trans. Am. Microsc. Soc. 9: 188–194), and Prevot and Bartoli (1978, Ann. Parasitol. (Paris) 53: 561–575).

On 24 July 1983, a female osprey, in poor physical condition, was found in the Chain Lakes area in Alberta, Canada

(50°15'N, 114°13'W) by personnel of the Fish and Wildlife Division and taken to the Calgary Zoo for hospitalization. The bird died of avian cholera (*Pasteurella multocida*) on 7 September, and was submitted to the Veterinary Laboratory, Animal Health Division, Airdrie, Alberta for examination. The kidneys were removed at necropsy and preserved in 10% buffered formalin. Portions of kidney tissue were prepared for histological examination and the remainder dissected. Histological examination of kidney tissue revealed multiple cross-sections of trematodes (Fig. 1). The trematodes were normally present in pairs within dilated kidney tubules lined by cuboidal cells. In

Received for publication 13 March 1984.