

## **Helminths of the Beluga Whale (*Delphinapterus leucas*) from the Mackenzie River Delta, Northwest Territories**

Authors: Wazura, K. W., Strong, J. T., Glenn, C. L., and Bush, Albert O.

Source: Journal of Wildlife Diseases, 22(3) : 440-442

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-22.3.440>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Helminths of the Beluga Whale (*Delphinapterus leucas*) from the Mackenzie River Delta, Northwest Territories

K. W. Wazura,<sup>1</sup> J. T. Strong,<sup>2</sup> C. L. Glenn,<sup>1,3</sup> and Albert O. Bush,<sup>1,3</sup> <sup>1</sup>Department of Zoology, Brandon University, Brandon, Manitoba R7A 6A9, Canada; and <sup>2</sup>Marine Mammals Management Division, Department of Fisheries and Oceans, Freshwater Institute, 501 University Crescent, Winnipeg, Manitoba R3T 2N6, Canada

There are scattered reports on the helminth fauna of the beluga whale from Arctic waters of North America. Lyster (1940, Can. J. Res. 18D: 395-409) reported *Anisakis simplex*, *Anisakis* sp. and *Corynosoma strumosum* from beluga collected in the Gulf of St. Lawrence. Brandly and Rausch (1950, Arctic 3: 105-107) reported the presence of the nematode *Trichinella* sp. from beluga collected in Alaskan waters. Doan and Douglas (1953, Bull. Fish. Res. Board Can. 98: 1-27) reported *Anisakis simplex* and *Stenurus arcticus* (= *Pharurus pallasii*) from whales collected at Churchill, Manitoba. Brodie (1971, J. Fish. Res. Board Can. 28: 1309-1318) reported *Crassicauda* sp. from the middle ear of beluga collected in Cumberland Sound. The identification of this nematode in the middle ear is questionable; *Crassicauda* is normally a parasite of the renal system. These specimens were likely *Pharurus pallasii* (Arnold and Gas-kin, 1975, Can. J. Zool. 53: 713-735). In an annotated list of parasites from sea mammals in North America, Margolis and Dailey (1972, NOAA Tech. Rep. NMFS SSRF-647, 23 pp.) added the following species to those reported in beluga: *Hadwenius seymouri*, *Pharurus oserskiae* (= *P. pallasii*), *Corynosoma semerme*, *C. similis* and *C. wegneri*. Arnold and Gas-kin (1975, op. cit.) reported *Stenurus arc-tomarinus* in beluga collected from the Mackenzie River Delta and Churchill,

Manitoba; they also reported *Pharurus pallasii* in whales from the same two lo-calities plus New Brunswick. Kenyon and Kenyon (1977, J. Wildl. Dis. 13: 338-340) reported on the prevalence of *Pharurus pallasii* in beluga collected in the Chur-chill River, Manitoba. Burns and Seaman (1985, Final Report, Outer Continental Shelf Environmental Assessment Pro-gram, Alaska Dept. of Fish and Game, Fairbanks, 129 pp.) reported *Otopho-caenurus oserskoi* (= *P. pallasii*) in beluga from Alaska. In this note, we present data on the helminth fauna of 10 beluga col-lected in the Mackenzie River Delta, Northwest Territories, Canada.

Between 7 and 25 July 1984 we exam-ined 10 carcasses of whales which had been killed by Inuvialuit hunters in the Kugmallit Bay region of the Mackenzie Delta. Most necropsies were completed within 24 hr following host death and all necropsies were conducted under field conditions. The following procedures were employed.

The stomach was tied off at the anterior and posterior ends and each of the three compartments was examined macroscop-ically for helminths. A sample of dia-phragm (4 × 8 cm) was removed and pre-served in AFA. The anterior and posterior ends of the intestine were tied, the intes-tine was then removed from the carcass and divided into 20 approximately equal sections. The anterior 7 to 8 cm of each section (posterior 7 to 8 cm for rectum) was removed and preserved in AFA for detailed analysis in the laboratory. The re-maining portion of each intestinal section

Received for publication 28 October 1985.  
<sup>3</sup> To whom reprint requests should be sent.

TABLE 1. Relative intensity and location of helminths from 10 beluga from the Mackenzie Delta.

| Host characteristics |        |             | Species of helminth                 |                                   |                                   |                                     |                                      |
|----------------------|--------|-------------|-------------------------------------|-----------------------------------|-----------------------------------|-------------------------------------|--------------------------------------|
| No.                  | Sex    | Length (cm) | <i>Contracaecum</i><br>sp. (larvae) | <i>Anisakis</i><br><i>simplex</i> | <i>Pharurus</i><br><i>pallasi</i> | <i>Hadwenius</i><br><i>seymouri</i> | <i>Leucasiella</i><br><i>arctica</i> |
| 1                    | Male   | 424         | L <sup>a</sup> 1, 2 <sup>b</sup>    |                                   | H 3                               | M 2                                 |                                      |
| 2                    | Female | 310         | M 1                                 | L 1                               |                                   | L 2                                 |                                      |
| 3                    | Male   | 410         | H 1                                 |                                   | M 3                               |                                     |                                      |
| 4                    | Male   | 418         | L 1                                 |                                   | H 3                               |                                     |                                      |
| 5                    | Male   | 431         | H 1                                 |                                   | M 3                               |                                     |                                      |
| 6                    | Female | 371         | H 1                                 |                                   | M 3                               |                                     |                                      |
| 7                    | Male   | 415         | L 1                                 |                                   | M 3                               | L 2                                 | L 4                                  |
| 8                    | Male   | 443         | M 1, 2                              |                                   | H 3                               | L 2                                 |                                      |
| 9                    | Male   | 349         | L 1, 2                              |                                   | M 3                               |                                     |                                      |
| 10                   | Male   | 401         | M 1                                 | L 1                               |                                   |                                     |                                      |

<sup>a</sup> Relative intensity categories were: L = less than 20 worms; M = 20–80 worms; H = greater than 80 worms.

<sup>b</sup> Site in host: 1 = stomach; 2 = intestine; 3 = ears; 4 = rectum.

was examined macroscopically at necropsy. Solid organs (e.g., heart, lung, kidney, liver) and their associated vessels and ducts were sliced and examined using a 20-mesh sieve and river water. Particular attention was paid to the tympanic membrane and auditory ossicles of the middle ear and to the blowhole and sinuses.

In the laboratory, the samples of diaphragm were examined microscopically, but no digestions were performed. Each of the subsamples from each of the intestinal sections was opened, the mucosal surfaces scraped and the contents and scrapings were diluted with water. This mixture was then examined microscopically with a dissecting microscope at 68×.

Because most of each necropsy was conducted in the field, and time limitations were strict, it was impossible to make accurate counts of the helminths encountered. Therefore the data obtained are largely qualitative, however to allow some quantitative assessment of relative intensity, the helminths were recorded as being present in low (<20 individuals), moderate (20–80 individuals) and high (>80 individuals) numbers.

Nematodes were fixed in acetic acid, stored in a mixture of 70% ethanol and 10% glycerin and examined as temporary wet mounts in lactophenol. Trematodes

were fixed in AFA and stained in Ehrlich's or Delafield's hematoxylin. Representative specimens of helminths found in this analysis have been deposited in the U.S. National Parasite Collection, Beltsville, Maryland 20705, USA (Accession Nos. 78969–78973), the National Museum of Canada Invertebrate Collection, National Museum of Natural Sciences, Invertebrate Zoology Division, Ottawa, Ontario K1A 0M8, Canada (Accession Nos. NMCP1985-0171–0175) and the University of Alberta Parasitology Collection, Department of Zoology, University of Alberta, Edmonton, Alberta T6G 2E9, Canada (Accession Nos. UAPC 11103–11107).

Five species of helminths (two trematodes, three nematodes) were found (Table 1). No beluga was helminth-free, a maximum of four species was found in one individual. Most organs were helminth-free. Intestinal sections 7 through 20 were free of helminths.

*Hadwenius seymouri* Price, 1932 was found in four beluga, in low to moderate numbers, and always within the first six sections of the intestine (ca. 6 meters). *Leucasiella arctica* Delyamure and Kleinenberg, 1958 was rare. One host harbored three individuals in the rectum.

*Pharurus pallasi* (van Beneden, 1870) Arnold and Gaskin, 1975 was recovered

from the ears of eight hosts. Three of the hosts had >80 worms, the others had between 20 and 80. *Anisakis simplex* (Rudolphi, 1809) Baylis, 1920 was rare. One host was infected with one female, a second host was infected with four males. All specimens were located in the first compartment of the stomach. Fourth stage larvae of *Contracaecum* sp. were the most prevalent and abundant helminth encountered. All hosts were infected and the infections ranged from low to high numbers. *Contracaecum* sp. occurred in all three of the stomach compartments and were found in the small intestine in the first six sections.

With the exception of *Contracaecum*, all species have been reported previously in beluga. This is the first report of *Leucasiella arctica* and *Contracaecum* from Nearctic waters and the genus *Contracaecum* is reported from beluga for the first time.

Although the data presented are not strictly quantitative, beluga from the Kugmallit Bay region of the MacKenzie Delta did not appear to be seriously parasitized by helminths. In fact, most organs (including much of the small intestine) were virtually helminth-free.

The authors thank the Inuvialuit of East Whitefish Station and Ms. Patt Weaver of the Marine Mammals Management Division, Fisheries and Oceans Division at the Freshwater Institute for their assistance in whale necropsies. An anonymous referee enlightened us considerably on the literature of beluga helminths. We also thank Dr. Robin Overstreet for confirming the identification of *Contracaecum*. This project was funded in part by a grant from the Research Fund of Brandon University (CLG) and an N.S.E.R.C. Operating Grant (AOB). K. W. Wazura participated as a student assistant in a Department of Fisheries and Oceans program.

*Journal of Wildlife Diseases*, 22(3), 1986, pp. 442-445

## Methods of Urine Collection for Male White-tailed Deer

**Terry J. Kreeger and Glenn D. Del Giudice**, Department of Fisheries and Wildlife, University of Minnesota, St. Paul, Minnesota 55108, USA; **Ulysses S. Seal**, Research Service, Veteran's Administration Medical Center, Minneapolis, Minnesota 55417, USA and Departments of Biochemistry and Fisheries and Wildlife, University of Minnesota, St. Paul, Minnesota 55108, USA; and **Patrick D. Korns**, Forest Wildlife Populations and Research Group, Minnesota Department of Natural Resources, Grand Rapids, Minnesota 55744, USA

Wildlife biologists are increasing their use of physiological and biochemical indices to assess individual or collective condition. Hematological and chemical analyses have been used widely in many species to study disease (Davis et al., 1981, *Infectious Diseases of Wild Mammals*, 2nd Ed., Iowa State University Press, Ames, Iowa, 446 pp.), reproduction (Plotka et al., 1977, *Biol. Reprod.* 16: 340-343), nutri-

tion (Seal et al., 1978, *J. Wildl. Manage.* 44: 776-790), and stress (Rehbinder and Edquist, 1981, *Acta Vet. Scand.* 22: 480-492). Blood parameters, however, can be influenced by drug immobilization and handling (Seal et al., 1972, *J. Wildl. Manage.* 36: 1034-1040; Mautz et al., 1980, *J. Wildl. Manage.* 44: 343-351). Urine may be less affected by factors invoking a stress response (Warren and Whelan, 1981, *J. Wildl. Dis.* 17: 479-483).

Urinalysis is used extensively for diagnosis of disease in humans (Harrison et al.,

Received for publication 6 August 1985.