

Isolation of *Actinobacillus suis* from a Canada Goose (*Branta canadensis*)

Authors: Maddux, R. L, Chengappa, M. M., and McLaughlin, B. G.

Source: Journal of Wildlife Diseases, 23(3) : 483-484

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-23.3.483>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Isolation of *Actinobacillus suis* from a Canada Goose (*Branta canadensis*)

R. L. Maddux,^{1,3} M. M. Chengappa,¹ and B. G. McLaughlin,² ¹ Department of Microbiology and ² Department of Pathology, Murray State University Breathitt Veterinary Center, P.O. Box 2000, Hopkinsville, Kentucky 42240, USA.

³ Author to whom reprint requests should be addressed.

ABSTRACT: Blindness from conjunctivitis caused by *Actinobacillus suis* was found in a Canada goose (*Branta canadensis*) from a wildlife refuge in Kentucky. Lesions were not observed elsewhere and other body organs were grossly normal. *Actinobacillus* spp. in birds is rare and this is apparently the first report of conjunctivitis resulting from this infection in waterfowl.

Key words: *Branta canadensis*, Canada goose, *Actinobacillus suis*, conjunctivitis, bacteriology, case report.

On 20 December 1985 a live, adult female Canada goose (*Branta canadensis*) was submitted to the Murray State University Breathitt Veterinary Center, Hopkinsville, Kentucky. The goose had been captured by hand on the Sauerheber Wildlife Refuge near Henderson, Kentucky. The bird had matted eyes and appeared blind at the time of capture. There were approximately 13,000 geese and 12,000 ducks on the refuge at the time. According to the wildlife manager, no other birds appeared ill.

Prior to euthanasia the bird was bright and alert, although thin and moderately dehydrated. The upper and lower eyelids of both eyes were matted closed by a tenacious grey exudate. The conjunctivae were hyperemic and the corneas were clear. Lesions were not observed in the frontal sinuses or the lower respiratory tract. All other body systems were grossly normal at necropsy.

Selected specimens were submitted for bacterial culture and processed according to standard methods (Carter, 1984). Sections of liver, lung, intestine and ocular swabs were inoculated onto trypticase soy

agar containing 5% sheep blood and incubated in 6-8% CO₂ at 37 C. Bacteria were not isolated from the liver. *Escherichia coli* was isolated from the lung, ocular swabs and intestine. The ocular and lung cultures produced small grey colonies of gram-negative rod-shaped bacteria exhibiting β -hemolysis. The organism was oxidase and catalase positive. On the basis of biochemical tests (Krieg and Holt, 1984; Lennette et al., 1985), the organism was identified as *Actinobacillus suis*. The isolate did not display the sticky and adherent properties characteristic of *Actinobacillus* spp. Pathogenicity to mice was demonstrated by intraperitoneal injection with 0.25 ml of a 6 hr broth culture. Death of the mice occurred within 12 hr and the isolate was recovered from the heart blood.

Microscopically, the cutaneous surfaces of the eyelids were hyperkeratotic and acanthotic. Accumulations of necrotic inflammatory cells mixed with cellular debris were on the skin surface and within the superficial keratinized layers. The conjunctival surfaces were thickened and hyperplastic. Numerous heterophils were migrating through the epidermis toward the surface. The dermal tissues contained numerous dense aggregations of mixed inflammatory cells. Heterophils and large mononuclear cells were predominant. The lower respiratory tract was normal.

Actinobacillus suis commonly causes arthritis, pneumonia, pericarditis, nephritis and subcutaneous abscesses in young swine. It is occasionally isolated from the upper respiratory tract of horses (Carter, 1984). The recovery of *Actinobacillus* spp.

from birds is rare. An *Actinobacillus lignieresii*-like organism has been reported to have caused salpingitis in Peking ducks by Bisgaard (1975). Hacking and Sileo (1977) reported a hemolytic *Actinobacillus* sp. from six waterfowl, three with periocular serous exudation and two with airsacculitis and bronchopneumonia. To our knowledge, the present case documents the first report of *A. suis* from a goose with conjunctivitis. Since *Actinobacillus* spp. infections can spread rapidly in young mammals, it was feared the recovery of *A. suis* from this bird could have epizootic significance. However, the geese left the refuge shortly thereafter and no further cases were observed.

The authors wish to thank Sherry Greer for technical assistance.

LITERATURE CITED

- BISGAARD, M. 1975. Characterization of atypical *Actinobacillus lignieresii* isolated from ducks with salpingitis and peritonitis. *Nordisk Veterinary-Medicine* 27: 378-383.
- CARTER, G. R. 1984. *Diagnostic procedures in veterinary bacteriology and mycology*, 4th ed. Thomas Books, Springfield, Illinois, 515 pp.
- HACKING, M. A., AND L. SILEO. 1977. Isolation of a hemolytic *Actinobacillus* sp. from waterfowl. *Journal of Wildlife Diseases* 13: 69-73.
- KRIEG, N. R., AND J. G. HOLT. 1984. *Bergey's manual of systemic bacteriology*, Vol. 1. Williams and Wilkins, Baltimore, Maryland, 964 pp.
- LENNETTE, E. H., A. BALLOWS, W. J. HAUSLER, JR., AND H. J. SHADOMY. 1985. *Manual of clinical microbiology*, 4th ed. American Society for Microbiology, Washington, D.C., 1149 pp.

Received for publication 18 September 1986.