



Gastrointestinal Helminths of Fish-Eating Birds from Chiloe Island, Chile

Authors: Torres, Patricio, Ruíz, Eileen, Gesche, Walter, and Montefusco, Aldo

Source: Journal of Wildlife Diseases, 27(1) : 178-179

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-27.1.178>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Gastrointestinal Helminths of Fish-Eating Birds from Chiloe Island, Chile

Patricio Torres, Eileen Ruiz, Walter Gesche, and Aldo Montefusco, Institute of Parasitology, Universidad Austral de Chile, Valdivia, Chile

ABSTRACT: Of four species of fish-eating birds from Chiloe Island (Chile), three were infected with eight species of helminths. *Tetrabothrius* sp. was found in *Larus scoresbii*. *Tetrabothrius cylindraceus*, *Profilicollis antarcticus*, *Anomotaenia dominicanus*, *Stephanoprora denticulata*, *Capillaria* sp. and *P. antarcticus* were found in *Larus dominicanus*. *Contracaecum rudolphii* and *Corynosoma* sp. infected *Phalacrocorax olivaceus*. With the exception of *S. denticulata*, *C. rudolphii* and *Capillaria* sp., the above helminths are reported for the first time from Chile.

Key words: Helminth parasites, piscivorous birds, *Larus* spp., *Phalacrocorax olivaceus*.

Helminthological data on fish-eating birds of Chile are scarce (Torres et al., 1982, 1983a). Thus, in April 1989, 10 fish-eating birds were examined for gastrointestinal helminths as part of a research project in southern Chile at Lake Huillinco, Lake Natri, and the coastal zone of Yaldad, all on Chiloe Island (Table 1; between 42°30' and 43°15'S).

The birds were killed by shotgun; the gastrointestinal tract was removed, kept at 4 C, and then within 4 hr examined for helminths (Torres et al., 1974). Helminths were preserved following techniques in Pritchard and Kruze (1982); cestodes and trematodes were dyed with Semichon's Acetic Carmin and Malzacher's stain.

With the exception of *Stephanoprora denticulata*, *Contracaecum rudolphii* and *Capillaria* sp., these helminths are reported for the first time in Chile (Table 1). In addition, *Profilicollis antarcticus* is recorded for the first time in *Larus dominicanus*; this acanthocephalan was described from *Chionis alba* (Gmelin) of the Shetland Islands (Antarctic) (Zdzitowiecki, 1985). *Chionis alba* lives in the Magallanes Strait and coasts of Tierra del Fuego up to Antarctica and migrates occasionally to Chiloe (Araya and Millie, 1986).

Anomotaenia dominicanus and *Tetrabothrius cylindraceus* have been recorded

in *L. dominicanus* on the Falkland Islands (Odening, 1982) and in the Antarctic (Zdzitowiecki and Szelenbaum-Cielecka, 1984).

Contracaecum rudolphii was earlier reported in South America and Argentina (Szidat and Nani, 1951) and in Chile (Torres et al., 1982, 1983a) in *Phalacrocorax olivaceus* and *Larus* spp. The wide geographic distribution of *C. rudolphii* (Barus et al., 1978) and *S. denticulata* (Torres et al., 1983b) is related to the migration of some of their numerous definitive hosts; as in the case of *C. rudolphii* wide distribution may be attributed to the wide range of fishes that act as intermediate hosts.

Larvae of *Contracaecum multipapillatum* Von Drasche, 1882 have been reported in fish (*Mugil cephalus*) in the north of Chile (Fernández, 1987), but the definitive host has not been determined in this area.

In different fresh-water ecosystems of the south of Chile the infection by larvae of *Contracaecum* sp. in introduced (Torres and Cubillos, 1987) and autochthonous fishes (Torres et al., 1988) has been established. The predation, consumption of viscera, or spread of these fish by the fishermen is possibly contributing to the infection of the fish-eating birds which act as definitive hosts.

Stephanoprora denticulata was reported previously in *Podiceps major* and *Pelecanus thagus* in the Valdivia River estuary (Torres et al., 1982).

The material collected was deposited in the collection of the Institute of Parasitology, Universidad Austral de Chile, Valdivia, Chile (IPUAT N° 0038-0048).

This study was supported by Grant N° 0069/1989 (FONDECYT) and Project N° S-89-17, Dirección de Investigación, Universidad Austral de Chile. We thank R. Schlatter (Institute of Zoology, Universidad Austral de Chile) for his help in the identification of birds, and the Centro de

TABLE 1. Occurrence of helminth parasites in fish-eating birds from Chiloé Island, Chile.

Birds	Location	n ^a /n ^b	Infected birds (intensity)							
			TEC ^c	TES	AND	STD	COR	CAS	PRA	COS
1 ^d	YA ^e	2/1	0	1 (1)	0	0	0	0	0	0
2	YA	1/1	1 (1)	0	0	0	0	0	1 (2)	0
2	LH	2/2	1 (1)	0	1 (2)	1 (2)	0	1 (1)	1 (9)	0
3	LH	1/0	0	0	0	0	0	0	0	0
4	LH	2/2	0	0	0	0	2 (27-83)	0	0	1 (3)
4	LN	2/1	0	0	0	0	1 (4)	0	0	0

^a Number of birds sampled.

^b Total of infected birds.

^c TEC, *Tetrabothrius cylindraceus*; TES, *Tetrabothrius* sp.; AND, *Anomotaenia dominicanus*; STD, *Stephanoprora denticulata*; COR, *Contracaecum rudolphii*; CAS, *Capillaria* sp.; PRA, *Profilicollis antarcticus*; COS, *Corynosoma* sp.

^d 1, *Larus scoresbii*; 2, *Larus dominicanus*; 3, *Ceryle torquatus*; 4, *Phalacrocorax olivaceus*.

^e YA, Coastal zone of Yaldad (43°07'S, 73°44'W); LH, Lake Huillincó (42°38'S, 74°00'W); LN, Lake Natri (42°47'S, 73°50'W).

Investigaciones Marinas (Universidad Austral) for allowing the use of their laboratory in Yaldad.

LITERATURE CITED

- ARAYA, B., AND G. MILLIE. 1986. Guía de Campo de las aves de Chile. Editorial Universitaria, Santiago, Chile, 389 pp.
- BARUS, V., T. P. SERGEEVA, M. D. SONIN, AND K. M. RYZHIKOV. 1978. Helminths of fish-eating birds of the Palaearctic Region I. Nematoda. W. Junk Publishers, Prague, Czechoslovakia, 318 pp.
- FERNÁNDEZ, J. 1987. Los parásitos de la lisa *Mugil cephalus* L., en Chile: Sistemática y aspectos poblacionales (Perciformes: Mugilidae). Gayana 51: 3-58.
- ODENING, K. 1982. Cestoden aus Flugvögeln der Sudshetlands (Antarktis) und der Falklandinseln (Malwinen). Angewandte Parasitologie 23: 202-223.
- PRITCHARD, M. H., AND G. O. KRUIZE. 1982. The collection and preservation of animal parasites. University of Nebraska Press, Lincoln, Nebraska, 141 pp.
- SZIDAT, L., AND A. NANI. 1951. Diplostomiasis cerebri del pejerrey. Revista del Instituto Nacional de Investigación de las Ciencias Naturales 1: 324-384.
- TORRES, P., R. FRANJOLA, L. YAÑEZ, V. DIAZ, E. GONZALEZ, AND M. MONTECINOS. 1974. Estudio preliminar sobre helmintos y artrópodos de *Gallus gallus domesticus* en la provincia de Valdivia, Chile. Boletín Chileno de Parasitología 29: 115-117.
- , L. FIGUEROA, A. SALDIVIA, AND J. BARRIENTOS. 1982. Gastrointestinal helminths of fish-eating birds from the Valdivia River, Chile. The Journal of Parasitology 68: 1157.
- , V. SIERPE, AND R. SCHLATTER. 1983a. Occurrence of *Contracaecum rudolphii* in new hosts in Chile. Zeitschrift für Parasitenkunde 69: 397-399.
- , L. FIGUEROA, AND A. SALDIVIA. 1983b. *Stephanoprora denticulata* (Trematoda, Echinostomatidae) en gaviotas del sur de Chile. Boletín Chileno de Parasitología 38: 33-34.
- , AND V. CUBILLOS. 1987. Infección por larvas de *Contracaecum* (Nematoda, Anisakidae) en salmónidos introducidos en Chile. Journal of Veterinary Medicine B 34: 177-182.
- , J. ARENAS, A. NEIRA, X. CABEZAS, C. COVARRUBIAS, C. JARA, C. GALLARDO, AND M. CAMPOS. 1988. Nematodos anisakidos en peces autóctonos de la cuenca del río Valdivia, Chile. Boletín Chileno de Parasitología 43: 37-41.
- ZDZITOWIECKI, K., AND D. SZELENBAUM-CIELECKA. 1984. *Anomotaenia dominicana* (Railliet et Henry, 1912) (Cestoda, Dilepididae) from the Dominican gull *Larus dominicanus* Licht. of the Antarctic. Acta Parasitologica Polonica 29: 49-58.
- . 1985. Acanthocephalans of birds from South Shetlands (Antarctic). Acta Parasitologica Polonica 30: 11-24.

Received for publication 7 February 1990.