

A REVIEW OF DISEASES OF PARASITES OF THE KAFUE LECHWE (KOBUS LECHE KAFUENSIS)

Author: Stafford, K. J.

Source: Journal of Wildlife Diseases, 27(4): 661-667

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-27.4.661

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A REVIEW OF DISEASES OF PARASITES OF THE KAFUE LECHWE (KOBUS LECHE KAFUENSIS)

K. J. Stafford

Department of Clinical Studies. Samora Machel School of Veterinary Medicine, UNZA, P.O. Box 32379, Lusaka, Zambia

ABSTRACT: The diseases of the Kafue lechwe (Kobus leche kafuensis) are reviewed in this paper. Kafue lechwe are an important natural resource for Zambia. Bovine tuberculosis is widespread within the lechwe population and they are host to many parasites, especially the warble Strobiloestrous vanzyli.

Key words: Kobus leche kafuensis, bovine tuberculosis, Mycobacterium bovis, endoparasites, Strobiloestrous vanzyli, ectoparasites, review.

INTRODUCTION

Kafue lechwe (Kobus leche kafuensis) are found in the natural state only within the Kafue Flats area of Zambia. The 65,000 lechwe (Howard et al., 1988) are a "natural resource of national importance" for Zambia (Vesey-Fitzgerald, 1965). They are utilized as a source of meat, hides and trophies. They are important as a tourist attraction and in maintaining the fertility of the Kafue river flats and fisheries (Mitchell and Uvs, 1961; Howard, 1977). Lechwe have been domesticated (Bainbridge, 1970) and are being stocked onto game farms in Zambia. In the past lechwe were cropped for meat by traditional kills (Ansell, 1955) and controlled culling, utilizing an abbatoir, was carried out for a short period in the late 1960's (Bainbridge, 1970; Pullan, 1983).

This review is to provide veterinarians and biologists with information on the diseases and parasites of Kafue lechwe.

INFECTIOUS DISEASES

Tuberculous lesions of bone were found in a Kafue lechwe (Leroux, 1958) and pulmonary tuberculosis caused by *Mycobacterium bovis* has been diagnosed in lechwe (Gallagher et al., 1972; Macadam et al., 1974; Clancey, 1977). The prevalence of tuberculosis in lechwe has varied (Table 1) but recent work would indicate that about 16% of adult male animals are affected (Stafford et al., 1990). Lechwe are semi-aquatic animals living in large groups

which become concentrated during the rains and early dry season (November to June) when their range is flooded (Gallagher et al., 1972). Crowding is important in the transmission of tuberculosis (Thoen and Hines, 1981) and Sayer and Van Lavieren (1975) speculated that the disease spread when the lechwe were concentrated at the beginning of the dry season and were suffering from "nutritional stress" (Gallagher et al., 1972). There are few predators in the area inhabited by the Kafue lechwe. Gallagher et al. (1972) estimated that tuberculosis killed 20% of the lechwe population each year and may be the major factor limiting population growth.

Lechwe are in contact with cattle when they graze together on the fresh grass which grows as the flood recedes. Cattle may have introduced tuberculosis into the lechwe population (Dillmann, 1976; Rottcher, 1976); $\leq 25\%$ of cattle from the Kafue Flats showed tubercular lesions on meat inspection (Anonymous, 1965) and $\leq 40\%$ may be affected (Rottcher, 1976). Infected lechwe may introduce the disease to cattle and other game species if they are moved onto cattle-game farms. When lechwe are killed for meat there is a potential public health problem as $\leq 80\%$ of carcasses may be tuberculous (Macadam et al., 1974; Dillmann, 1976; Krauss et al., 1986). The control of tuberculosis amongst free-ranging lechwe on the Kafue Flats would be extremely difficult. Lechwe being moved

Number exam- ined	Num- ber af- fected	Percent affected	Reference
125	45	36	Gallagher et al., 1972
300	90	30	Dillmann, 1976
86	29	34	Gallagher et al., 1972
7	7	100-	Clancey, 1977
63	33	52	Clancey, 1977
141	46	33	Krauss et al., 1986
41	33	80 -	Krauss et al., 1986
147	5	3 [⊾]	Krauss et al., 1986
92	15	16	Stafford et al., 1990

TABLE 1. Tuberculosis in the Kafue lechwe.

* Cachectic animals with suspected tuberculosis.

^b Dead animals collected from Kafue Flats.

onto game farms could be tested for tuberculosis; however the efficacy of testing lechwe for tuberculosis either by skin testing (Macadam, 1973) or serological testing is unknown. Tuberculosis was not a problem among Black lechwe (*Kobus leche smithemani*); they inhabit an environ-

TABLE 2. Serological Surveys of Kafue lechwe.

ment similar to the Kafue lechwe but do not come in contact with cattle (Grimsdell and Bell, 1975).

Serological surveys have been conducted to determine the disease status of lechwe (Table 2). Antibodies against brucella organisms, the Leptospira interogans serovars autumnalis autumnalis and ballum ballum, Pseudomonas pseudomallei, Francisella tularensis, Chlamydia psittaci, Coxiella brunetii, infectious bovine rhinotracheitis/infectious pustular vaginitis virus, parainfluenza 111 virus and reoviruses 1,11,111 were found in <10% of lechwe (Table 2), Antibodies against bluetongue virus were found in 50% of a small number of lechwe. The significance of these findings to the health of lechwe and domestic animals in contact with them are unclear (Krauss et al., 1986) but may become more obvious as lechwe are moved onto cattle-game farms. Foot and Mouth

Infectious agent	Number of samples	Results	Reference
Leptospirosis*	45	3 positive*	K. Stafford (unpubl. data)
Brucellosis ^b	200	positive	Krauss et al., 1986
Brucellosis	55	2 positive	K. Stafford (unpubl. data)
Mycobacterium paratuberculosis	200°	all negative	Krauss et al., 1986
Anaplasma marginale	31	all negative	Dillmann, 1976
Anaplasma centrale	31	all negative	Dillmann, 1976
Pseudomonas pseudomallei	200°	positive	Krauss et al. 1986
Francisella tularemia	200°	positive	Krauss et al. 1986
Chlamydia psittaci	200°	positive	Krauss et al. 1986
Coxiella burnetti	200°	positive	Krauss et al. 1986
Infectious Bovine Rhinotracheitis	10 ^c	positive	Krauss et al. 1986
Parainfluenza 111	10 ^e	positive	Krauss et al. 1986
Reovirus 1, 11, 111	5°	positive	Krauss et al. 1986
Bluetongue	10	5 positive	Krauss et al. 1986
Babesia bigemina	31	all negative	Dillmann, 1976
Theileria parva	31	all negative	Dillmann, 1976
Theileria mutans	31	all negative	Dillmann, 1976

• The microagglutination test was used to test for antibodies against the following serovars: Leptospira interogans serovar, canicola canicola, icterohaemorrhagiae copenhageni, pomona pomona, grippotyphosa grippotyphosa, tarrasovi tarrasovi, australis australis, australis bratislava, sejroe hardjo, hebdomadis hebdomadis, mini mini, bataviae bataviae, javanica javanica, pyrogenes pyrogenes, autumnalis autumnalis, ballum ballum, celledoni celledoni, cynopteri cynopoteri and shermani shermani. Three were positive for autumnalis autumnalis autumnalis and one for Ballum ballum.

^b Serum agglutination test.

^d Rose Bengal test and slide agglutination test.

* Positive to both tests (d).

'The exact number of positive sera was not stated.

^c The exact number tested was not stated.

Oesophagostomum lechwei	(Leroux, 1955)
Bunostomum cobi	(Leroux, 1958)
Bunostomum sp.	(Leroux, 1932)
Trichostrongylus colubriformis a	(Leroux, 1930)
Haemonchus bedfordi	(Leroux, 1955)
Haemonchus contortus	(Leroux, 1930)
Cooperia curticei	(Leroux, 1932)
Dictyocaulus filaria	(Fitzsimmons, 1961)
Dictyocaulus viviparus	(Robinette and Child, 1964)
Thelazia rhodesi	(Robinette and Child, 1964)
Thelazia sp.	(Leroux, 1955)
Onchocerca sp.	(Leroux, 1947, 1955)
Setaria bicoronata	(Yeh, 1959)
Setaria boulengeri	(Yeh, 1959)
Setaria hornbyi	(Yeh, 1959)
Setaria sp.	(Leroux, 1955)
Trichuris sp.	(Ratcliffe, 1930)

TABLE 3. Nematode parasites of Kafue lechwe.

TABLE 4. Trematode and cestode parasites of Kafue lechwe.

• Round (1968) questioned this finding.

disease has not been diagnosed clinically in Kafue lechwe and the causative virus was not isolated from 50 animals sampled by Dillmann (1976). Lechwe are probably succeptible to the disease but their importance as a source of the disease for domestic animals is unknown (Howard, 1977). Heartwater disease caused by Cowdria ruminantium was diagnosed in lechwe with heavy loads of Amblyomma variegatum ticks held on a game farm (G. S. Pandey, pers. comm.). Abscesses were found in the testes of three lechwe (Robinette and Child, 1964) but the lesions were not examined bacteriologically.

HELMINTH PARASITES

Kafue lechwe are host to many parasites (Tables 3, 4, 5; Round, 1968). The effect of these parasites on lechwe is unknown but Robinette and Child (1964) doubted if internal parasites were primary cause of mortalities. Strongyle nematode eggs were found in 74 of 76 faecal samples taken from recently shot adult lechwe (K. Stafford, unpubl. data). The number of eggs per gram were <100 in 68 of these samples. Coccidial oocysts were found in many

Fasciola gigantica	(Gallagher et al., 1972;
	Wright et al., 1979a)
Fasciola sp.	(Leroux, 1957)
Cotylophoron cotylophoron	(Leroux, 1932)
Cotylophoron sp.	(Robinette and Child, 1964)
Carmyerius sp.*	(Wright et al., 1979b)
Gastrothylax crumenifer	(Leroux, 1932)
Paramphistomum cervi ^b	(Leroux, 1932)
Gigantocotyle explanatum	(Leroux, 1932)
Gigantocotyle formosanum	(Round, 1968)
Gigantocotyle lerouxi	(Yeh, 1957)
Choerocotyloides onotragi	(Prudhoe et al., 1964)
Homalogaster paloniae	(Leroux, 1955)
Stephanopharynx com-	
pactus	(Wright et al., 1979b)
Calicophoron calicophoron	(Wright et al., 1979a)
Schistosoma leiperi	(Leroux, 1955)
Schistosoma margrebowiei	(Leroux, 1933, 1955)
Schistosoma matthei	(Leroux, 1958)
Schistosoma spindalis ^e	(Ortlepp, 1961)
Taenia spp.	(Leroux, 1955, 1957;
	Robinette and
	Child, 1964)

* May be C. mancupatus or C. spasiosus (Wright et al., 1979b). ^b May be P. microbothrium or P. phillerouxi (Wright et al., 1979b; Round, 1968).

^c Round (1968) questioned this finding.

of these faecal samples. Coccidia have not previously been reported in lechwe. Lungworms of three different species were found in about 55% of culled lechwe (Dillmann, 1976) but only two species have been definitely identified (Table 3). Thelazia rhodesi found in small numbers in a

Ectoparasites of Kafue lechwe. TABLE 5.

Strobiloestrous vanzyli	(Zumpt, 1961; Howard, 1975,
Damalinia sp.	1980, 1985) (Robinette and Child, 1964)
Amblyomma variegatum	(Colbo, 1973)
Rhipicephalus evertsi	(Colbo, 1973)
Rhipicephalus sanguineus	(Colbo, 1973)
Rhipicephalus appendic- ulatis	(Robinette and Child, 1964)
Muscoid headflies	(Robinette and Child, 1964)

few lechwe (Robinette and Child, 1964) may be the filarial worms referred to by Sayer and Van Lavieren (1975) as the cause of blindness in lechwe. Onchocerca sp. probably O. gibsoni were found in fibrous lesions in periarticular regions especially near the stifle joint (Leroux, 1947, 1955; Robinette and Child, 1964). Approximately 35% of the lechwe in Lochinvar National Park were affected by onchocerciasis which caused lameness in some animals (Dillmann, 1976). Setaria sp. were "abundant" in lechwe (Sayer and Van Lavieren, 1975; Robinette and Chile, 1964; Dillmann, 1976) and three species have been identified (Table 3). Fasciola gigantica were identified in adult lechwe (Wright et al., 1979a). Female lechwe had a higher prevalence of liver fluke (22%) than males (6%) possibly due to the female habit of grazing more in the marsh along the water's edge (Gallagher et al., 1972). However Fasciola eggs were found in 68 of 76 (89%) faecal samples taken from male carcasses and examined by K. Stafford (unpubl. data). Amphistome flukes were regularly seen in lechwe (Table 4). Schistosoma margrebowiei have been found in young lechwe (Leroux, 1933, 1955) and S. leiperi in older animals (Wright et al., 1979a; Howard et al., 1982). Lechwe <1yr of age had a mean infection of 24 S. margrebowiei and no S. leiperi while adult animals >4 yr of age had a mean infection of 273 S. leiperi and no S. margrebowiei. One adult lechwe had over 1,000 S. leiperi (Howard et al., 1982). S. matthei found in lechwe by Leroux (1958) was not identified by Howard et al. (1982) in lechwe sharing pasture with cattle, but was seen in waterbuck (Kobus defassa) living remote from cattle.

ECTOPARASITES

The oestrid larvae of *Strobiloestrous* vanzyli, the southern lechwe fly, were found in the subcutis of the back and hindquarters of lechwe (Zumpt, 1961; Sayer and Van Lavieren, 1975; Howard, 1980). In a sample of 526 lechwe of all ages, 95% of those <1-yr-old were affected with warbles ($\bar{x} = 149$ warbles/animal) while only 20% of older animals were affected (\bar{x} = 8.6 warbles/host) (Howard, 1985). The warbles are on the lechwe from November until September when the prepupae emerge, drop on the ground and pupate in the soil. Adult flies emerge within 1 mo (Howard, 1985). S trobiloestrous vanzyli may not be distinct from S. erikssoni (Howard, 1980). Leroux (1942) reported finding similar warbles on lechwe and cattle and concluded that they were from the same species of fly. The large numbers of larvae seen on young lechwe led Mitchell and Uys (1961) to conclude that the warbles caused "heavy" mortalities but evidence to support this conclusion is lacking. A small unidentified muscoid fly was seen to be a nuisance to lechwe during the warmer months of October and November. Lechwe sought relief from the fly by rubbing their heads in grass or standing in a tight circle (Robinette and Child, 1964). Lice (Damalinia sp.) were common on lechwe (Robinette and Child, 1964). Various species of ticks have been found on lechwe (Table 5) (Ganagarajah, 1976). Rhipicephalus appendiculatis was found in small numbers (<5) only on the perianal region of 20 of 75 lechwe examined, the remaining 55 animals had no ticks (Robinette and Child, 1964).

MISCELLANEOUS PATHOLOGICAL CONDITIONS

A fibrosarcoma and other malignant but unidentified neoplasms were found in lechwe by Robinette and Child (1964).

DISCUSSION

The population of Kafue lechwe has dropped from 94,075 in 1970 to 65,000 in 1988 (Howard et al., 1988). This reduction in numbers has been due to a loss in available habitat, poaching and perhaps disease and malnutrition.

Tuberculosis and warbles are the major disease problems of lechwe. Tuberculosis was seen in most cachectic animals examined (Table 1). However it was only seen in 3% of 147 autopsies conducted on lechwe found dead (Krauss et al., 1986), the majority had died as a result of poaching injury. The significance of the effect of warbles and other internal parasites on the health of lechwe is unknown. The lameness seen in many lechwe may be due to onchocerciasis and/or poaching injuries.

To maintain the present population of lechwe attempts may have to be made to control tuberculosis. Increasing the available habitat and reducing contact between cattle and lechwe have been suggested as means of reducing mortalities due to tuberculosis and malnutrition (Rottcher, 1976). A suggestion by Dillmann (1976) that the culling of 5,000 lechwe/yr might reduce tuberculosis has not been confirmed. However the present level of tuberculosis is much lower than that seen in the seventies and this might be explained in part by the fact that lechwe numbers droppped to 41,000 in 1982 (Howard et al., 1988). Research should be directed towards examining the accuracy and feasibility of tuberculosis testing of lechwe in the wild, in transit and on game farms. The monitoring of lechwe held on game farms stocking cattle free from tuberculosis may demonstrate whether the disease can be maintained in a lechwe population isolated from tubercular cattle.

The limited number of serological surveys conducted suggests the need to evaluate further the prevalence of antibodies to bluetongue, Rift Valley Fever, Bovine Herpes virus and other diseases within the lechwe population. Lechwe are being moved onto farms which stock domestic ruminants. It is important the the transmission of disease from lechwe to domestic ruminants and vice versa, if it occurs, be monitored. Will the warbles of Strobiloestrous vanzyli affect cattle (Leroux, 1942) and Schistosoma matthei infest lechwe on cattle-game farms? Disease problems such as heartwater and heavy tick infections not reported from wild lechwe may become problems on game farms. The cropping scheme proposed and supported by the World Wildlife Fund and initiated in 1989 is an opportunity to study further the disease conditions seen in adult lechwe. As the lechwe will be cropped for meat, postmortem examination will have to be carried out to ensure the quality of the meat being sold. Blood obtained from the cropped animals could be used for serological studies. The cropping and farming of lechwe may be necessary to ensure the survival of the species in Zambia. Both these enterprises demand that lechwe are healthy, fit for human consumption and are not a source of disease for other wild and domestic stock held on game farms.

LITERATURE CITED

- ANONYMOUS. 1965. Annual Report of the Department of Veterinary and Tsetse Control Services. Ministry of Agriculture, Lusaka, Zambia, 32 pp.
- ANSELL, W. F. 1955. The declining red lechwe. The Oryx 3: 15-18.
- BAINBRIDGE, W. A. 1970. Report of the Chief Game Officer, Game and Fisheries Annual Report for the Year 1968. Ministry of Natural Resources and Tourism, Lusaka, Zambia, 72 pp.
- CLANCEY, J. K. 1977. The incidence of tuberculosis in lechwe (marsh antelope). Tubercle 58: 151-156.
- COLBO, M. H. 1973. Ticks of Zambian wild animals: A preliminary checklist. The Puku 7: 97-105.
- DILLMANN, J. S. 1976. Compiled final report. Veterinary Wildlife Research Section, Department of Veterinary Services, Ministry of Agriculture, Lusaka, Zambia, 395 pp.
- FITZSIMMONS, W. M. 1961. A possible dwarf form of *Dictyocaulus filaria* from a waterbuck, *Kobus leche* in northern Rhodesia. Veterinary Record 73: 1111–1112.
- GALLAGHER, J., I. MACADAM, J. SAYER, AND J. P. VAN LAVIEREN. 1972. Pulmonary tuberculosis in free living lechwe antelope in Zambia. Tropical Animal Health and Production 4: 204-213.
- GANAGARAJAH, M. 1976. A preliminary check list and host list of ticks of Zambia. Published by National Council for Scientific Research, Lusaka, Zambia, 16 pp.
- GRIMSDELL, J. J., AND R. H. BELL. 1975. Black lechwe research project, final report. Published by National Council for Scientific Research, Lusaka, Zambia, 175 pp.
- HOWARD, G. 1975. Differentially delayed development in the lechwe warble, *Strobiloestrous vanzyli* (Diptera:Oestridae). Parasitology 71(2): 10.

666 JOURNAL OF WILDLIFE DISEASES, VOL. 27, NO. 4, OCTOBER 1991

- —. 1977. National Parks of the Kafue Basin. In Development and ecology in the lower Kafue Basin in the 1970s, G. Williams and G. Howard (eds.). The University of Zambia, Lusaka, Zambia, pp. 47–56.
- . 1980. Immature stages and affinities of the southern Lechwe warblefly, *Strobiloestrous van*zylt Zumpt (Diptera:Oestridae). Journal of Natural History 14: 669–683.
- 1985. Warble infestations of lechwe antelope. Proceedings of the Fifth International Conference on Wildlife Diseases, Uppsala, Sweden, 23 pp.
- C. MALAMBO, AND R. C. JEFFERY. 1988.
 Kafue lechwe population census May/June, 1988.
 A report to the Director, National Parks and Wildlife Service, Lusaka, Zambia, 8 pp.
- C. A. WRIGHT, AND V. R. SOUTHGATE. 1982.
 Schistosome infections of lechwe and waterbuck in Zambia—A preliminary report. *In* Wildlife Diseases of the Pacific Basin and other countries, M. E. Fowler (ed.). Wildlife Disease Association, Ames, Iowa, pp. 136–138.
- KRAUSS, H., D. ROTTCHER, D. WEISS, K. DANNER, AND O. J. HUBSCHLE. 1986. Wildlife as a potential source of infection in domestic animals— Studies on game in Zambia. Animal Research and Development 24: 41–58.
- LEROUX, P. L. 1930. Helminths of domestic stock in the Republic of South Africa. Journal of South African Veterinary Medical Association 1(4): 43– 65.
 - 1932. List of helminths collected from mammals and birds in the Mazabuka area of northern Rhodesia. Annual Report (1931) of the Department of Animal Health, Northern Rhodesia, Appendix 3: 31–34.
 - —. 1933. A preliminary note on Bilharzia margrebowiei, a new parasite of ruminants and possibly man in northern Rhodesia. Journal of Helminthology 11: 57-62.
 - ——. 1942. Report of the Associate Veterinary Research Officer. Annual report of the Department of Animal Health, Northern Rhodesia, 32 pp.
 - 1947. Onchocerca in the tendons of insertion of the M. rectus femoris of a Red Lechwe (Onotragus leche) a species of Kob, Kafue River, Mazebuka, northern Rhodesia. Transcripts of the Royal Society of Tropical Medicine and Hygeine 41: 9.
 - —. 1955. Game conservation and severe helminthiasis of stock in national game reserves. Transcripts of the Royal Society of Tropical Medicine and Hygeine 49: 293.
 - -----. 1957. Report to the government of the Federation of Rhodesia and Nyasaland on the control of parasite diseases in livestock. FAO Reprint number 696, Rome, Italy, 47 pp.
 - ----. 1958. Bone tuberculosis in a Red Lechwe

(Adenota lechwe) in northern Rhodesia. Transcripts of the Royal Society of Tropical Medicine and Hygeine 52: 15.

- MACADAM, I. 1973. Handling lechwe for vaccination and tuberculosis testing. Veterinary Record 92: 461.
- ------, J. GALLAGHER, AND J. MCKAY. 1974. Experimental tuberculosis in lechwe antelope in Zambia. Tropical Animal Health and Production 6: 107–109.
- MITCHELL, B. L., AND J. M. UYS. 1961. The problem of the lechwe (*Kobus leche*) on the Kafue Flats. The Oryx 6: 171-183.
- ORTLEPP, R. J. 1961. South African helminths. Onderstepoort Journal of Veterinary Science 9: 311– 336.
- PRUDHOE, S., L. YEW, AND L. F. KHALI. 1964. A new amphistome from ruminants in northern Rhodesia. Journal of Helminthology 38: 57-62.
- PULLAN, R. A. 1983. The use of wildlife in the development of Zambia. In Natural resources in tropical countries, Ooi Jin Bee (ed.). Singapore University Press, Singapore, Malaysia, pp. 267– 325.
- RATCLIFFE, H. C. 1930. Animal parasites. In Report of the Zoological Society of Philadelphia, H. Fox (ed.). Philadelphia, Pennsylvania, pp. 57-60.
- ROBINETTE, W. L., AND G. F. CHILD. 1964. Notes on the biology of the Lechwe (*Kobus leche*). The Puku 2: 84–117.
- ROTTCHER, D. 1976. Research on wildlife diseases in Zambia. Proceedings of the fourth regional conference on wildlife for East and Central Africa, Ministry of Lords, Natural Resources and Tourism, Lusaka, Zambia, pp. 147–157.
- ROUND, M. C. 1968. Checklist of the helminth parasites of African mammals. Technical Communication, number 38, Commonwealth Bureau of Helminthology, The Commonwealth Agricultural Bureau, England, 25 pp.
- SAYER, J. A., AND L. P. VAN LAVIEREN. 1975. The ecology of the Kafue lechwe population of Zambia before the operation of hydroelectric dams on the Kafue River. East African Wildlife Journal 13: 9–37.
- STAFFORD, K., A. MWEENE, J. BAER, AND G. PANDEY. 1990. Tuberculosis in Kafue lechwe (Kobus leche kafuensis). Sixth International Conference on Wildlife Diseases. Berlin, German Democratic Republic, 76 pp.
- THOEN, C. O., AND E. M. HINES. 1981. Tuberculosis. In Infectious diseases of wild animals, J. W. Davis, L. H. Karstad and D. O. Trainer (eds.). Iowa State University Press, Ames, Iowa, pp. 263– 274.
- VESEY FITZGERALD, L. D. 1965. Lechwe pastures. The Puku 3: 143-147.
- WRIGHT, C., V. SOUTHGATE, AND G. HOWARD. 1979a. Observations on the life cycle of Schistosoma margrebowiei and its possible interac-

_

tions with *S. letpert* in Zambia. Journal of Natural History 13: 499–506.

- , ____, AND _____. 1979b. A note on the life cycles of some amphistome flukes in Zambia. Journal of Helminthology 53: 251–252.
- YEH, L. S. 1957. On a new paramphistomid trematode *Gigantocotyle lerouxi* from the stomach of a Red Lechwe (*Onotragus leche*) from northern Rhodesia. Parasitology 47: 432-434.
- ——. 1959. On the identity of the filarial worms. Transcripts of the Royal Society of Tropical Medicine and Hygeine 52: 297.
- ZUMPT, F. 1961. The enigma of Strobiloestrous brauer (Diptera: Oestridae). Proceedings of the Royal Entomological Society, London 30: 93-102.

Received for publication 6 December 1989.