

Haematozoa of Fishes in Humboldt Bay, California

Authors: Hill, Jeffrey P., and Hendrickson, Gary L.

Source: Journal of Wildlife Diseases, 27(4): 701-705

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-27.4.701

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Haematozoa of Fishes in Humboldt Bay, California

Jeffrey P. Hill^{1,2} and Gary L. Hendrickson,¹ ¹ Department of Fisheries, Humboldt State University, Arcata, California 95521, USA. ² Present address: Department of Biological Sciences, Campus Box 8007, Idaho State University, Pocatello, Idaho 83209, USA

ABSTRACT: Five hundred seven fish representing 45 species from Humboldt Bay, California (USA) were examined for blood parasites. Four fish (<1%) from two species were infected. Haemogregarina leptocotti sp. n. is described from one of 33 staghorn sculpin (Leptocottus armatus). Haemogregarina roelofsi sp. n. is described from three of 15 black rockfish (Sebastes melanops). Gametocytes of H. leptocotti sp. n. averaged 6.1 \times 2.1 μ m with a 2.7 \times 1.7 μ m oval nucleus; those of H. roelofsi sp. n. averaged 5.5 \times 2.7 µm with a 2.5 \times 2.2 µm rectangular nucleus. Neither species of parasite had distinct chromatin granules, a polar cap, or more than one gametocyte in an infected cell. Haematozoa are relatively rare in fishes of the northeastern Pacific Ocean.

Key words: Haemogregarina roelofsi sp. n., Haemogregarina leptocotti sp. n., haematozoa, marine fish, parasite, survey.

Little is known about the occurrence and distribution of haematozoa in marine fish off of the western coast of North America. Laird (1961) examined 10 padded sculpins (Artedius fenestralis) collected off of Vancouver Island, British Columbia (Canada). Two fish were infected with Haemogregarina bigemina. Burreson and Pratt (1972) surveyed 498 fish of 11 species from Oregon coastal waters. They found 67 of 392 (17%) English sole (Parophrys vetulus) infected with Trypanosoma pacifica (Burreson and Pratt, 1972). Burreson (1979) described Cryptobia (Trypanoplasma) beckeri from cabezon (Scorpaenichthys marmoratus) collected from Oregon coastal waters. The present study describes two new species of the genus Haemogregarina from marine fish from Humboldt Bay, Humboldt County, California (USA).

Fish were collected from Humboldt Bay monthly from September 1983 through August 1984. Collections were made with a semi-balloon otter trawl, with 19-mm square nylon mesh in the body and a 6-mm mesh liner in the cod end. Trawls (15 min to 1 hr) were made at low tide when fish were concentrated in bay channels.

Fish were kept alive in seawater and transported back to the laboratory (Humboldt State University, Arcata, California 95521, USA). They were killed by a sharp blow to the head and the tail was severed. At least one drop of free flowing blood was discarded before making blood smears. Blood smears were air dried, methanol fixed, and stained with Wright-giemsa stain for 10 min and then moved to a phosphate buffer for 10 min. Slides then were rinsed in tap water and air dried.

Each of two slides from each fish was examined under oil immersion $(1,000 \times)$ for 20 min. Haematozoa were identified and the total number of parasites observed was recorded. Relative intensity of infection of infected hosts was rated using the method of Becker and Katz (1966): 0, no parasites observed; +1, one to five parasites observed; +2, six to 20 parasites observed; +3, five to 10 parasites observed in the first minute; +4, one or more parasites observed in each of the first five fields.

Blood from 507 fish representing 45 species was examined (Table 1). Only four fish (<1%) were infected. One of 33 (3%) staghorn sculpin (Leptocottus armatus) was infected with Haemogregarina leptocotti sp. n. The infected fish was 20.8 cm in standard length, and had a +2 infection. Three of 13 (23%) black rock fish (Sebastes melanops) were infected with Haemogregarina roelofsi sp. n. Infected rock fish ranged from 17 to 23 cm in standard length and all fish had +1 infections. These parasites are described herein, based on morphology of gametocytes in host erythrocytes.

702 JOURNAL OF WILDLIFE DISEASES, VOL. 27, NO. 4, OCTOBER 1991

Fish	Number sampled	Fish	Number sampled
Carcharhinidae		Hexagrammidae	
Mustelus henlei	17	Hexagrammos decagrammus	11
Triakis semifasciata	4	Hexagrammos lagocephalus	2
Myliobatididae		Ophiodon elongatus	5
Muliobatis californica	40	Cottidae	
Clupeidae		Artedius notospilotus	5
Alosa sapidissima	6	Enophrys bison	12
Clupea harengus	28	Hemilepidotus hemilepidotus	10
Dorosoma petenense	2	Leptocottus armatus	33
Engraulidae		Oligocottus snyderi	1
Engraulis mordax	7	Scorpaenichthys marmoratus	1
Osmeridae		Embiotocidae	
Hypomesus pretiosus	3	Amphistichus rhodoterus	12
Spirinchus starksi	9	Cymatogaster aggregata	19
Spirinchus thaleichthys	14	Rhacochilus vacca	4
Gadidae		Embiotoca lateralis	1
Microgadus proximus	30	Hyperprosopon argenteum	22
Atherinidae		Hyperprosopon ellipticum	3
Atherinops affinis	2	Phanerodon furcatus	35
Atherinopsis californiensis	4	Pholidae	
Syngnathidae		Pholis ornata	8
Sungnathus leptorhynchus	24	Ammodytidae	
Gasterosteidae		Ammodytes hexapterus	1
Aulorhynchus flavidus	15	Gobiidae	
Gasterosteus aculeatus	31	Lepidogobius lepidus	10
Scorpaenidae		Bothidae	
Sebastes sp.	1	Citharichthys stigmaeus	2
Sebastes auriculatus	13	Paralichthys californicus	3
Sebastes flavidus	1	Pleuronectidae	
Sebastes melanops ^b	13	Parophrys vetulus	21
Sebastes mystinus	5	Platichthys stellatus	15
Sebastes rastrelliger	1	Cynoglossidae	
		Symphurus atricauda	1

TABLE 1. Fish from Humboldt Bay, California (USA) examined for haematozoa.

• One fish infected with Haemogregarina leptocotti. ^b Three fish infected with Haemogregarina roelofsi.



FIGURES 1, 2. 1. Haemogregarina leptocotti sp. n. in an erythrocyte of Leptocottus armatus. Wrightgiemsa stain. 2. Haemogregarina roelofsi sp. n. in an erythrocyte of Sebastes melanops. Wright-giemsa stain.

Haemogregarina leptocotti sp. n. (Figure 1)

Description: Measurements are in micrometers and based on 20 specimens (ranges given followed by means in parentheses). One gametocyte per infected host cell. Gametocytes 5.3-7.8 (6.1) long by 1.2-2.9 (2.1) wide, attenuated towards the end closest to their nucleus. Gametocytes reniform, depth of cell curvature 0.0-1.0 (0.4). Nucleus not centered in cell, oval, 2.4-3.4 (2.7) along long axis of parasite cell by 1.0-1.9(1.7) along short axis of parasite cell. Longer cytoplasmic extremity from the nucleus 1.7-2.7 (2.2) long. Shorter cytoplasmic extremity from nucleus 1.0-2.4 (1.5) long. Nucleus stained deep purple and lacked distinct chromatin granules. Cytoplasm stained pale blue. No basophilic granulation or polar cap observed. Parasitized erythrocytes [7.3-10.7 (8.9) long by 5.1-8.3 (6.6) wide not significantly longer (t = 0.25, P > 0.50) or wider (t = 0.43, P > 0.50) than non-parasitized cells [7.3– 10.6 (8.9) long by 4.8-8.2 (6.5) wide]. No difference in nucleus length (t = 0.43, P> 0.50) or width (t = 0.28, P > 0.50) between parasitized [3.8-4.9 (4.3) by 3.2-4.3 (3.6)] and non-parasitized [3.9-4.8 (4.2) by 3.4-4.4 (3.6)] erythrocytes.

Type host: Leptocottus armatus (Cottidae).

Location in host: Erythrocytes.

Type locality: Humboldt Bay, California, USA (40° 48'N, 124°10'W).

Type specimens: Syntype deposited in U.S. National Museum, Beltsville, Maryland 20705, as USNM 81941.

Etymology: This species is named after the genus of its type host.

Remarks: Two other haemogregarine gametocytes are similar to those of *H. lep*tocotti. Haemogregarina coelorhynchi (Laird, 1952) has been reported from Coelorhynchus australis (Macrouridae) and Physiculus bachus (Moridae) collected off the coast of New Zealand (Laird, 1952). Haemogregarina coelorhynchi is 5.0-5.9 μ m long × 2.5-3.1 μ m wide and is distinguished by its polar cap. Haemogregarina mavori (Laird and Bullock, 1969) has only been reported from Macrozoarces americanus (Zoarcidae) collected from coastal waters of eastern Canada (Laird and Bullock, 1969). Haemogregarina mavori is $5.5-8.6 \ \mu m \ long \times 2.2-3.5 \ \mu m \ wide and$ is distinguished by a nucleus which contains 8 to 18 discrete chromatin granules.

Four haemogregarines have been described from other marine cottids: H. bigemina, H. cotti, H. cotti scorpi and H. myoxocephali. Haemogregarina bigemi*na* from padded sculpin is the only marine haemogregarine described from the western coast of North America. Gametocytes of *H*. bigemina are much larger (10.1–14.1 μ m long × 1.0–1.9 μ m wide) (Laird, 1961) than those of H. leptocotti. Henry (1910) did not adequately describe H. cotti scorpii and it may be synonymous with H. cotti. Haemogregarina cotti has been reported from Cottus bubalis and C. scorpius collected from European coastal waters. It is also much larger (12 μ m long × 2.5 µm wide) than H. leptocotti (Laird and Bullock, 1969). Haemogregarina myoxocephali has only been described from Myoxocephalus octodecemspinosus collected in coastal waters of eastern Canada and the United States. Haemogregarina myoxocephali is 4.9–9.6 μ m long × 1.6– 3.7 μ m wide and is distinguished from the parasite in this study by the occurrence of 8 to 14 discrete chromatin granules in its nucleus (Laird and Bullock, 1969).

Haemogregarina roelofsi sp. n. (Figure 2)

Description: Measurements are in micrometers and based on 16 specimens (ranges given followed by means in parentheses). One gametocyte per infected host cell. Most gametocytes oval 4.3-6.4 (5.5) long by 1.9-3.7 (2.7) wide. Some gametocytes are slightly reniform, depth of cell curvature 0.0-1.0 (0.4). Nucleus usually near center of cell, somewhat trapezoidal, 1.6-3.4 (2.2) along long axis of parasite cell by 1.4-3.7 (2.5) along short axis of parasite cell. Longer cytoplasmic ex-

tremity from the nucleus 1.4–2.5 (1.9) long. Shorter cytoplasmic extremity from nucleus 0.9-2.0 (1.5) long. Nucleus stained red violet and lacked distinct chromatin granules. Cytoplasm did not stain. No basophilic granulation or polar cap observed. Parasitized erythrocytes [8.9–11.7 (10.1) long by 5.8-8.3 (7.2) wide not significantly longer (t = 1.39, 0.10 < P < 0.20) or wider (t = 1.12, 0.30 < P < 0.40) than nonparasitized cells [8.7-10.5 (9.7) long by 5.8-7.8 (6.9) wide]. No difference in nucleus length (t = 0.78, 0.40 < P < 0.50) or width (t = 1.02, 0.30 < P < 0.40) between parasitized [3.3-5.1 (4.2) by 2.4-3.4 (2.9)] and non-parasitized [3.6-4.9 (4.4) by 2.4-3.9 (3.1)] erythrocytes.

Type host: Sebastes melanops (Scorpaenidae).

Location in host: Erythrocytes.

Type locality: Humboldt Bay, California, USA (40°48'N, 124°10'W).

Type specimens: Syntype deposited in U.S. National Museum, Beltsville, Maryland 20705, as USNM 81940.

Etymology: Named in honor of Dr. Terry Roelofs, Professor of Fisheries, Humboldt State University, for his contributions to fisheries biology and education.

Remarks: Two other haemogregarine gametocytes are similar to those of H. roelofsi. Haemogregarina coelorhynchi (Laird, 1952) has been reported from Coelorhynchus australis (Macrouridae) and Physiculus bachus (Moridae) collected off the coast of New Zealand (Laird, 1952). Haemogregarina coelorhynchi is 5.0-5.9 μ m long \times 2.5–3.1 μ m wide and is distinguished by its polar cap. Haemogregarina mavori (Laird and Bullock, 1969) has only been reported from Macrozoarces americanus (Zoarcidae) collected from coastal waters of eastern Canada (Laird and Bullock, 1969). Haemogregarina mavori is 5.5-8.6 μ m long × 2.2-3.5 μ m wide and is distinguished by a nucleus which contains 8 to 18 discrete chromatin granules.

One haemogregarine has been previously described from marine scorpaenids. Neumann (1909) briefly described *H. scor*- paenae from Scorpaena scrofa collected off the coast of Italy. Neumann (1909) encountered only a few parasites and suggested that his description of the parasite may be inadequate. Gametocytes occurred singly in red blood cells and were oval, 6 μ m long × 1.2 μ m wide. Gametocytes not in blood cells occurred in groups of four and were 17 μ m long × 1.5 μ m wide. Neumann's (1909) description of H. scorpaenae is inadequate and it is difficult to make comparisons. However, H. roelofsi is much wider than H. scorpaenae and occurs in a different host and in a substantially different geographic location.

Burreson and Pratt (1972) surveyed 67 staghorn sculpins from Oregon (USA) and detected no haemogregarines, but found 17% of the English sole (*Parophrys vetulus*) to be infected with a trypanosome. We surveyed 21 English sole and detected no haematozoa. Burreson (1979) reported a new cryptobiid from cabezon collected off of Oregon. We examined one cabezon and detected no haematozoa.

The prevalence of haematozoan infections in a given species of marine fish from the western coast of North America has been low: 3 to 23% (present study), 17% (Burreson and Pratt, 1972), and 20% (Laird, 1961). The percentage of species infected also has been low: 4% (present study) and 9% (Burreson and Pratt, 1972). Haematozoan surveys of marine fish from eastern North America often have found parasites to be more prevalent. In fact, Laird and Bullock (1969) and Khan (1986) found more than 50% of the individuals of some species infected. In addition, eastern surveys have regularly found over 30% of the fish species examined infected with haematozoa (Laird and Bullock, 1969; Daily, 1978; Khan, 1986).

The apparent difference in prevalence of haematozoa between eastern and western North America may be an artifact and a result of the small number of surveys conducted on the western coast. However, Khan (1986) suggested that the abundance of leech vectors controls the incidence and prevalence of haematozoan infections. Leeches known to transmit haematozoa occur along the Pacific coast (Burreson, 1977, 1979), but little is known about their distribution or abundance.

ACKNOWLEDGMENTS

Humboldt State University provided the vessel and equipment that made this work possible. We are also grateful to William Matter for review of the manuscript.

LITERATURE CITED

- BECKER, C. D., AND M. KATZ. 1966. Host relationships of *Cryptobia salmositica* (Protozoa: Mastigophora) in a western Washington hatchery stream. Transactions of the American Fisheries Society 95: 196–202.
- BURRESON, E. M. 1977. Oceanobdella pallida n. sp. (Hirudinea: Piscicolidae) from the English sole, Parophrys vetulus, in Oregon. Transactions of the American Microscopical Society 96: 526–530.
- . 1979. Structure and life cycle of *Trypanoplasma beckeri* sp. n. (Kinetoplastida), a parasite of the cabezon, *Scorpaenichthys marmoratus*, in Oregon coastal waters. The Journal of Protozoology 26: 343–347.

- , AND I. PRATT. 1972. Trypanosoma pacifica sp. n. from the English sole Parophrys vetulus Girard from Oregon. The Journal of Protozoology 19: 555-556.
- DAILY, D. D. 1978. Marine fish hematozoa from Maine. The Journal of Parasitology 64: 361-362.
- HENRY, H. 1910. On the haemoprotozoa of British sea-fish (a preliminary note). Journal of Pathology and Bacteriology 14: 463-465.
- KHAN, R. A. 1986. Haematozoa of marine fish from Ungava Bay and adjacent northwestern Atlantic Ocean. Canadian Journal of Zoology 64: 153– 157.
- LAIRD, M. 1952. New haemogregarines from New Zealand fishes. Transactions of the Royal Society of New Zealand 79: 589-600.
- ——. 1961. Trichodinids and other parasitic protozoa from the intertidal zone at Nanaimo, Vancouver Island. Canadian Journal of Zoology 39: 833–844.
- ------, AND W. L. BULLOCK. 1969. Marine fish haematozoa from New Brunswick and New England. Journal of the Fisheries Research Board of Canada 26: 1075-1102.
- NEUMANN, R. O. 1909. Studien über protozoische Parasiten im Blut von Meeresfischen. Zeitschrift für Hygiene und Infektionskrankheiten 64: 1– 112.

Received for publication 17 September 1990.