

Capturing and Tagging Free-ranging Bighorn Sheep

Authors: Jorgenson, Jon T., Samson, Judith, and Festa-Bianchet, Marco

Source: Journal of Wildlife Diseases, 27(4) : 733-734

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-27.4.733>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

REPLY TO LETTER TO THE EDITOR . . .

Capturing and Tagging Free-ranging Bighorn Sheep

Any technique to capture wild animals should strive to reduce stress, accidental injury, and mortality as much as possible. Consideration of the method to be used must also take into account such factors as manpower availability and time and budget constraints. In Jorgenson et al. (1990), we reported a technique that allowed a single researcher (even a "nonveterinarian wildlife worker") to capture and tag a large number of free-ranging bighorn sheep (*Ovis canadensis*). This method was inexpensive, safe, and resulted in a low mortality rate. No accidental deaths occurred once the antagonist, idazoxan, was available for trial. Our long-term research program (e.g., Festa-Bianchet, 1989) would not have been possible without this technique.

Kock (1991) stated that our bighorn sheep were not "really wild" and that our ability to approach within easy darting range was an unusual situation not likely to be duplicated anywhere else. We agree that the bighorns in our study were habituated to humans, and consider that a major asset of our study populations. This is certainly not, however, a unique situation. Our study involved three different populations and we have used the same ground darting technique to capture bighorns from six other populations. We also know of many other herds in Canada and the northwest United States where the technique is either used or feasible. Ground darting could be used for any species provided one could get within range and maintain contact with the animal following dart impact. The habituation of study animals is an old and well established technique for in-depth study of wildlife behavior and ecology. Examples of the use of habituated ungulates include studies on red deer (*Cervus elaphus*) (Clutton-Brock et al., 1982), Nilgiri thar (*Hemitragus hylocrius*) (Rice, 1988), moose (*Alces alces*) (Miquelle, 1990), chamois (*Rupicapra*

pyrenaica) (Locati and Lovari, 1990), kudu (*Tragelaphus strepsiceros*) (Owen-Smith, 1990), our studies on bighorn sheep and mountain goats, and many others.

Kock (1991) criticized comparisons made between our study and the study by Kock et al. (1987). The comparison made in Jorgenson et al. (1990) was between ground darting and helicopter darting to immobilize bighorn sheep. It would have been invalid for us to make comparisons between the various drugs used. Our intent was to question the conclusion in Kock et al. (1987) that drugs should not be used to immobilize bighorn sheep by making a distinction between delivery systems. While not stated as such in Kock et al. (1987), it is obvious from Kock (1991) that he agrees with us that helicopters should not be used to dart sheep.

Kock et al. (1987) did not distinguish between problems caused by the drugs and those associated with the helicopter chase. We acknowledge that it would be difficult to determine the exact contribution of the helicopter to stress and accidental deaths, but a comparison of studies that have used helicopters to dart sheep (Jessup et al., 1985a; Kock et al., 1987) and ground darting studies (Festa-Bianchet and Jorgenson, 1985; Jorgenson et al., 1990) suggests that the helicopter chase contributes significantly to accidental mortality.

Kock offers his opinion that xylazine is only useful for the sedation of captive or confined wildlife species and should not be recommended as a sole immobilizing agent. We take note of his opinion and have used xylazine to immobilize 45 mountain goats, first captured in box traps, and reversed the immobilization with idazoxan, without any drug-induced deaths. Our success in capturing almost 400 (includes all sheep captured using xylazine or a combination of xylazine and ketamine) free-ranging bighorn sheep, however, appears to argue against Kock's opinion.

It was stressed in Jorgenson et al. (1990) that the effectiveness of using xylazine was dependent on administering the drug in a manner that would elicit the least excitement from target animals. If such could not be achieved, then alternate drugs or methods would have to be considered. It was pointed out that higher doses would be required if sheep were to be darted from a helicopter. We certainly did not recommend using xylazine to immobilize sheep, as Kock (1991) implies, if helicopter darting was to be the mode of delivery.

Under the circumstances described in Jorgenson et al. (1990), xylazine proved to be a very effective immobilizing agent. Negative side effects were offset by the use of idazoxan. Kock (1991) suggested that other drugs such as opioids should be used in place of xylazine. Opioids such as etorphine hydrochloride and carfentanil citrate have been used in bighorn sheep but not completely without adverse side effects (Jessup et al., 1985b; Jessup and Clark, 1986). Induction times were comparable to those reported in Jorgenson et al. (1990) when only xylazine was used. Helicopters were used to dart sheep in those studies and some of the side effects may not have occurred had drugs been administered by a less stressful method. There would have been no advantage to using narcotics in our situation, especially considering the high cost and extreme hazard to humans. We captured 10 sheep with etorphine and found no advantages over the use of xylazine.

Finally, we are intrigued by Kock's definition of "long-term survival" as 1 wk. Nevertheless, the minimum survival from recovery from immobilization to 10 days post-capture before the availability of alpha-2 antagonists was 98.8% ($N = 169$) at Sheep River and Canmore, compared to 98.4% survival for sheep captured in a corral trap at Sheep River in 1981 to 1984.

LITERATURE CITED

- CLUTTON-BROCK, T. H., F. E. GUINNESS, AND S. D. ALBON. 1982. Red deer. University of Chicago Press, Chicago, Illinois, 378 pp.
- FESTA-BIANCHET, M. 1989. Individual differences, parasites, and the costs of reproduction for bighorn ewes (*Ovis canadensis*). *Journal of Animal Ecology* 58: 785-795.
- , AND J. T. JORGENSEN. 1985. Use of xylazine and ketamine to immobilize bighorn sheep in Alberta. *The Journal of Wildlife Management* 49: 162-165.
- JESSUP, D. A., K. R. JONES, R. MOHR, AND T. KUCERA. 1985a. Yohimbine antagonism to xylazine in free-ranging mule deer and desert bighorn sheep. *Journal of the American Veterinary Medical Association* 187: 1251-1253.
- , W. E. CLARK, K. R. JONES, R. CLARK, AND W. R. LANCE. 1985b. Immobilization of free-ranging desert bighorn sheep, tule elk, and wild horses, using Carfentanil and Xylazine: Reversal with naloxone, diprenorphine, and yohimbine. *Journal of the American Veterinary Medical Association* 187: 1253-1254.
- , AND B. CLARK. 1986. Wildlife restraint handbook, 2nd ed. California Department of Fish and Game. Rancho Cordova, California, 151 pp.
- JORGENSEN, J. T., J. SAMSON, AND M. FESTA-BIANCHET. 1990. Field immobilization of bighorn sheep with xylazine hydrochloride and antagonism with idazoxan. *Journal of Wildlife Diseases* 26: 522-527.
- KOCK, M. D. 1991. Letter to the editor. *Journal of Wildlife Diseases* 27: In press.
- , D. A. JESSUP, R. K. CLARK, C. E. FRANTI, AND R. A. WEAVER. 1987. Capture methods in five subspecies of free-ranging bighorn sheep: An evaluation of drop-net, drive net, chemical immobilization, and the net-gun. *Journal of Wildlife Diseases* 23: 634-640.
- LOCATI, M., AND S. LOVARI. 1990. Sexual differences in aggressive behaviour of the Apennine chamois. *Ethology* 84: 295-306.
- MIQUELLE, D. G. 1990. Why don't bull moose eat during the rut? *Behavioral Ecology and Sociobiology* 27: 145-151.
- OWEN-SMITH, N. 1990. Demography of a large herbivore, the greater kudu *Tragelaphus strepsiceros*, in relation to rainfall. *Journal of Animal Ecology* 59: 893-913.
- RICE, C. G. 1988. Habitat, population dynamics, and conservation of the Nilgiri thar, *Hemitragus hyloctus*. *Biological Conservation* 44: 137-156.

Received for publication 8 April 1991.

Jon T. Jorgenson,¹ Judith Samson,² and Marco Festa-Bianchet,³ ¹ Alberta Fish and Wildlife Division, #200 5920-1A St. S.W., Calgary, Alberta, Canada T2H 0G3; ² Associated Veterinary Clinics, 4204 4th St. N.W., Calgary, Alberta, Canada T2K 1A2; ³ Département de Biologie, Faculté des Sciences, Université de Sherbrooke, Sherbrooke, Québec, Canada J1K 2R1.