

Borrelia sp. in Ticks Recovered from White-tailed Deer in Alabama

Authors: Luckhart, Shirley, Mullen, Gary R., Durden, Lance A., and Wright, James C.

Source: Journal of Wildlife Diseases, 28(3): 449-452

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-28.3.449

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Borrelia sp. in Ticks Recovered from White-tailed Deer in Alabama

Shirley Luckhart,¹³ Gary R. Mullen,¹ Lance A. Durden,¹⁴ and James C. Wright,² ¹ Department of Entomology and Alabama Agricultural Experiment Station, Auburn University, Alabama 36949, USA; ² Department of Pathobiology, College of Veterinary Medicine, Auburn University, Alabama 36849, USA; ³ Present address: Department of Entomology, Rutgers University, New Brunswick, New Jersey 08903, USA. ⁴ Present address: Department of Biology, Georgia Southern University, Statesboro, Georgia 30460, USA

ABSTRACT: Six hundred sixty-five hunter-killed white-tailed deer (Odocoileus virginianus) from 18 counties in Alabama (USA) were examined for ticks. Most of the collections were made at state-operated wildlife management areas. Four species of ticks (n = 4,527) were recovered: the lone star tick Amblyomma americanum (n =482); the Gulf Coast tick A. maculatum (n =11); the winter tick Dermacentor albipictus (n = 1,242); and the black-legged tick Ixodes scapularis (n = 2,792). Fifty-six percent of the ticks (n = 2,555) were examined for *Borrelia* sp. spirochetes using an immunofluorescent, polyclonal antibody assay. Spirochetes were detected in I. scapularis (five females, seven males) from Barbour, Butler, Coosa, and Lee counties and A. americanum (four males, four nymphs) from Hale, Lee, and Wilcox counties. Area-specific prevalences in ticks were as high as 3.3% for I. scapularis and 3.8% for A. americanum.

Key words: Ticks, Amblyomma americanum, Ixodes scapularis, Lyme disease, Borrelia sp., white-tailed deer, Odocoileus virginianus, prevalence, survey.

First recognized in the United States in 1975, Lyme disease (LD) now accounts for the majority of cases of arthropod-borne diseases reported nationwide (Anonymous, 1989). Studies from the northeastern United States indicate that the etiologic agent of Lyme disease, Borrelia burgdorferi, is maintained in a complex multihost cycle involving principally the white-footed mouse (Peromyscus leucopus), the white-tailed deer (Odocoileus virginianus), and the deer tick (Ixodes dammini). The number of human cases of LD in the southeastern United States has grown steadily in recent years, but the vector species from this region remain uncertain. The first human case of LD was reported in Alabama in 1986 (Mullen and Piesman, 1987). About 80 additional confirmed cases have been reported since that time according to the Alabama Department of Public Health, with 13 of these cases reported in the literature (Woernle, 1989). Since the appearance of LD in Alabama, statewide efforts have been undertaken to collect and identify ticks parasitizing deer and to examine them for *Borrelia* spp. spirochetes.

During the 1985-86, 1988-89, and 1989-90 hunting seasons (mid-November through January), 665 deer were examined at State-operated hunter check stations and on private properties in 18 counties throughout Alabama (USA; Fig. 1). Most animals were examined at the following Wildlife Management Areas (WMA): Black Warrior WMA (Lawrence County), Skyline WMA (Jackson County), Kinterbish WMA (Sumter County), T. R. Miller WMA (Escambia County), Choccolocco WMA (Cleburne County), Oakmulgee WMA (Hale County), Barbour County WMA, Butler County WMA, Coosa County WMA, and Hollins WMA (Clay County). Animals were examined for ticks by brushing back the fur and visually, as well as tactilely, searching the ears, chin, chest, neck, axillae, escutcheon, back, and inguinal regions. Ticks were removed with forceps, transferred to vials containing slightly moistened cotton, transported to the laboratory, and maintained at 4 C until they were identified. Voucher tick specimens have been deposited with the United States National Tick Collection (Georgia Southern University, Statesboro, Georgia 30460, USA) under accession number RML 119,900.

Live ticks were examined for *Borrelia* sp. using direct and indirect immunofluorescent assays with polyclonal antisera.



FIGURE 1. Map of Alabama depicting counties in which ticks were collected from white-tailed deer and those counties in which ticks infected with Borrelia sp. were recovered.

fluorescent

direct

using

sp.

Q

examined

Ŀ.

The rationale for using polyclonal, rather than monoclonal, antisera was that we do not known how reliably the southern strains of B. burgdorferi respond to available monoclonal antibodies (e.g., H5332). The Borrelia use of polyclonal antibody tests thus reduces the prospect of overlooking Borrelia sp. infections even when they are present. Portions of midgut and salivary gland tissue were removed from each tick and teased apart on a glass slide. Dissections were air-dried and fixed with cold acetone. ticks Direct assays were performed as described by Piesman et al. (1986). Prepared slides Deer and control slides of cultured spirochetes were overlaid with fluorescein-labelled, **TABLE** high-titered rabbit anti-B. burgdorferi polyclonal conjugate (Guilford strain), incubated, washed, and examined using fluorescence microscopy. Indirect assays also

				Ticks exa	mined		
Date	Site	Number of deer examined	Amblyomma americanum N, M, F	Ixodes scapularis N, M, F	Dermacentor albipictus N, M, F	Amblyomma maculatum N, M, F	Positive ticks
16 Jun 88	Lee County	1	14, 4, 0	0	0	0	1 N Ambluomma americanum
18 Jun 88	Lee County	I	11, 1, 0	0	0	0	3 N Amblyomma americanum
29 Nov 88	Lee County	I	0	0, 6, 5	0	0	1 M Ixodes scapularis
2 Dec 88	Lee County	I	0	0, 4, 5	0, 1, 0	0	1 M Ixodes scapularis
25 Jan 89	Lee County	1	0, 1, 0	0, 4, 5	0	0	1 M Ambluomma americanum
26 May 89	Wilcox County	1	2, 8, 2	0	0	0	1 M Amblyomma americanum
13 Dec 89	Coosa WMA ^b	61	1, 13, 1	0, 39, 82	0	0, 2, 0	2 M, 2 F Ixodes scapularis
16 Dec 89	Barbour WMA	92	0	0, 164, 150	0	0	1 F Ixodes scapularis
12 Jan 90	Barbour WMA	27	0	0, 224, 171	0	0	1 M. 1 F Ixodes scapularis
27 Jan 90	Oakmulgee WMA	39	7, 29, 16	0, 36, 30	0	0	1 F Ixodes scapularis
							2 M Amblyomma americanum
31 Jan 90	Butler WMA	18	5, 8, 2	1, 58, 40	0	0	2 F Ixodes scapularis
• N, nymphs, M, male • WMA, wildlife man	s; F, females. sgement area.						

Date	Site	Number of ticks examined	Number of ticks positive	Prevalence (%)
	Ixodes	scapularis		
13 Dec 89	Coosa WMA [*]	121	4	3.3
16 Dec 89	Barbour WMA	314	1	0.3
12 Jan 90	Barbour WMA	395	2	0.5
27 Jan 90	Oakmulgee WMA	66	1	1.5
31 Jan 90	Butler WMA	99	2	2.0
	Amblyomm	a americanum		
27 Jan 90	Oakmulgee WMA	52	2	3.8

TABLE 2. Area-specific prevalence of ticks infected with Borrelia sp. in Alabama, 1989-90.

• WMA, wildlife management area.

were performed using unlabeled rabbit anti-*B. burgdorferi* antiserum and fluorescein-labeled goat antirabbit antiserum as described (Luckhart et al., 1991).

Four thousand five hundred twenty seven nymphal and adult ticks of four species were recovered from the deer examined. The majority of ticks collected were *Ixodes* scapularis (n = 2,792; 62%), followed by Dermacentor albipictus (n = 1,242; 27%), Amblyomma americanum (n = 482; 11%), and A. maculatum (n = 11; 0.2%). Two thousand five hundred fifty-five ticks (56%) were dissected and examined for spirochetes.

Twenty ticks (0.8%) were found infected with spirochetes which were IFA-positive and morphologically indistinguishable from cultured B. burgdorferi on control slides: five male and seven female I. scapularis from Barbour, Butler, Coosa, and Lee counties, and four nymphal and four male A. americanum from Hale, Lee, and Wilcox counties (Table 1). Five of these infected ticks were collected from deer at the same study site in Lee County where I. scapularis from cotton mice (Peromyscus gossypinus) previously were found to be infected with B. burgdorferi as confirmed using B. burgdorferi-specific murine monoclonal antibody H5332 (Luckhart et al., 1991). Efforts to isolate and culture spirochetes at this site have not been unsuccessful.

Area-specific prevalence was calculated for the December 1989 and January 1990 wildlife management area collections by dividing the number of infected ticks of one species from an area on a specific date by the total number of that species recovered and assayed from that area on the same date. Because only individual deer were examined on the dates indicated in Lee and Wilcox counties, no attempt was made to report prevalence of infection for these collections.

Prevalence of infection at the wildlife management areas ranged from 0.3 to 3.3% for *I. scapularis* and was 3.8% for the single IFA-positive *A. americanum* collection (Table 2). These findings are consistent with observations made in North Carolina by Magnarelli et al. (1986). They reported infection prevalences of 0.5% and 4.8% for *I. scapularis* and *A. americanum*, respectively, recovered from deer in Onslow County in December 1984.

In our previous field studies, *B. burg-dorferi*-infected *I. scapularis* immatures were recovered from rodents in areas of confirmed human cases of Lyme disease in Alabama (Luckhart et al., 1991). This supports the belief that *I. scapularis* plays a role in at least the maintenance cycle of *B. burgdorferi* among wild reservoir hosts. However, there is no conclusive evidence documenting the involvement of this tick in actual transmission of the Lyme disease agent to humans in the southeastern United States. In fact, *I. scapularis* is not commonly collected on humans in Alabama. Among more than 200 cases of human in-

festations with ticks in Alabama in 1990– 91, only eight involved *I. scapularis*; in contrast, 132 of those cases (65%) involved *A. americanum* (G. R. Mullen, unpubl. data). The detection of *B. burgdorferi*-like spirochetes in *A. americanum* therefore continues to raise the prospect that this species plays a role in transmission of *B. burgdorferi* among wild hosts and to domestic animals and humans in the southeastern United States.

We thank personnel from the Alabama Department of Conservation and Natural Resources, Game and Fish Division, and the Alabama Department of Forestry for their assistance in collecting ticks at the hunter check stations. We also acknowledge Joseph Piesman, Centers for Disease Control, Fort Collins, Colorado, for providing the polyclonal antibody for the tick assays, and the following individuals at Auburn University for laboratory and field support: Larry Swango, Sheila Smith, Lawrence J. Hribar, and C. Steven Murphree. Funding for this work was provided by a Student Research Grant from the Alabama Academy of Science, a Sigma Xi Grant-in-Aid of Research Award, and the Alabama Agricultural Experiment Station.

LITERATURE CITED

- ANONYMOUS. 1989. Lyme disease—United States, 1987 and 1988. Morbidity and Mortality Weekly Report 38: 668–672.
- LUCKHART, S., G. R. MULLEN, AND J. C. WRIGHT. 1991. Etiologic agent of Lyme disease, *Borrelia burgdorferi*, detected in ticks (Acari: Ixodidae) collected at a focus in Alabama. Journal of Medical Entomology 28: 652–657.
- MAGNARELLI, L. A., J. F. ANDERSON, C. S. APPERSON, D. FISH, R. C. JOHNSON, AND W. A. CHAPPELL. 1986. Spirochetes in ticks and antibodies to *Borrelia burgdorferi* in white-tailed deer from Connecticut, New York, and North Carolina. Journal of Wildlife Diseases 22: 178–188.
- MULLEN, G. R., AND J. PIESMAN. 1987. Serologically substantiated case of Lyme disease and potential tick vectors in Alabama. Alabama Journal of Medical Science 24: 306–307.
- PIESMAN, J., T. N. MATHER, J. G. DONAHUE, J. LE-VINE, J. D. CAMPBELL, S. J. KARAKASHIAN, AND A. SPIELMAN. 1986. Comparative prevalence of *Babesia microti* and *Borrelia burgdorferi* in four populations of *Ixodes dammini* in eastern Massachusetts. Acta Tropica 43: 263-270.
- WOERNLE, C. H. 1989. Surveillance for Lyme disease in Alabama. Alabama Medicine 58: 19-20.

Received for publication 6 May 1991.