

## **Encephalomyocarditis (EMC) Virus Recovered From Two Cotton Rats and a Raccoon 1**

Authors: GAINER, J. H., and BIGLER, W. I.

Source: Bulletin of the Wildlife Disease Association, 3(2) : 47-49

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-3.2.47>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Encephalomyocarditis (EMC) Virus Recovered From Two Cotton Rats and a Raccoon<sup>1</sup>

J. H. GAINER<sup>2</sup> and W. I. BIGLER

*Animal Disease Diagnostic Laboratory, Florida Department of Agriculture,  
Kissimmee, Florida and the Division of Veterinary Public Health, Florida State  
Board of Health, Jacksonville, Florida*

Received for publication 27 December, 1966

## ABSTRACT

Encephalomyocarditis in wild mammals is rarely reported. This paper documents the isolation of the encephalomyocarditis (EMC) virus in two cotton rats (*Sigmodon hispidus*) and a raccoon (*Procyon lotor*) from central Florida.

## INTRODUCTION

Neutralizing antibodies have been demonstrated in the sera of several rats: *Rattus rattus* and *R. alexandrinus* trapped in the United States<sup>10</sup>, *R. rattus* trapped in Panama<sup>11</sup>, and *R. rattus R. soridus conatus*, and *Melomys lutillus littoralis* in Australia<sup>12</sup>. Experimental studies have shown that certain species of rodents - the cotton rat, the hamster (*Mesocricetus auratus*)<sup>6</sup>, the multimammate rat (*Mastomys natalensis*)<sup>7</sup>, the black rat<sup>1</sup>, and the albino rat (*R. norvegicus*)<sup>8</sup>, are susceptible to infection with EMC virus. It is the understanding of the authors that EMC virus has been isolated from only four wild animals indigenous to the United States. In the first case, the virus was recovered by the Florida State Board of Health Laboratory from the brain of a squirrel submitted for rabies examination<sup>12</sup>. In the second, third, and fourth cases, the virus was isolated respectively from rat feces<sup>3</sup>, from a squirrel brain<sup>2</sup>, and a raccoon spleen<sup>2</sup>.

## MATERIALS AND METHODS

From February 22 to 24, 1966, 14 mammals, 1 wild bird, and 1 slug were live-trapped or collected as a part of an epizootiologic investigation following an outbreak of EMC in swine near Winter Haven, Florida<sup>4</sup>. Sera for neutralization tests and tissues for isolation attempts were taken from 6 cotton rats, 2 rice rats (*Oryzomys palustris*), 2 raccoons, 3 opossums (*Didelphis marsupialis*), and one civet cat (*Spilogale ambervalis*).

The mammals were captured in standard box-type live traps near the hog pen. All animals were anesthetized, bled and euthanized. Serums were stored at -60C for subsequent antibody determinations. Small portions of the heart, spleen, lung, brain and intestines were removed aseptically from each animal. The tissues were placed in separate petri dishes and frozen on dry ice for 1-3 days. They were transported to the Animal Disease Diagnostic Laboratory at Kissimmee, Florida, and stored in a -60C freezer until processed. Due to shortness of time, only the intestines were tested for the presence of virus. They were thawed and ground in sterile mortars and pestles with the aid of alundum. Twenty per cent suspensions of tissue were made with brain liver heart broth\* (BLHB) containing 1000 units of penicillin and 1000 mcg streptomycin per ml and 20-30% normal inactivated rabbit serum (NRS). The suspensions

\* Difco Company, Detroit, Michigan.

<sup>1</sup> A more detailed report of these findings is to be published in the *Cornell Veterinarian*<sup>4</sup>.

<sup>2</sup> Present address - National Cancer Institute, Wiscon Building, Room 4C-01, Bethesda, Maryland 20014.

were centrifuged at 3000 RPM for 30 minutes. The supernatant fluid was stored in 4 ml screwcap vials at -60°C. Weanling mice inoculated intracranially with .03 ml suspension were observed daily for clinical signs. Deaths occurring within 24 hours after inoculation were considered non-specific. All mice dying later than 1 day after inoculation were frozen for subsequent brain harvest. Infected brain material was aseptically removed, triturated in a sterile mortar and pestle and made into 20% suspensions with BLHB and centrifuged. The EMC virus was identified by incubating decimal dilutions of this first-mouse-passage brain suspension with NRS and EMC immune serum, inoculating mice, recording deaths, and comparing LD50's or harmonic mean survival times of controls with immune serum groups.

Serums were inactivated at 56°C. for 30 minutes. They were then tested in mice in a screening neutralization test against approximately 100 LD50's of virus. Mouse deaths were recorded; harmonic mean survival times (HMST's) of test serums were compared with HMST's of NRS as a control.

### RESULTS

Table 1 lists the results of the findings. As indicated, no detectable antibodies were present in the serums tested. Virus was isolated from 2 of 6 cotton rats and 1 of 2 raccoons.

### DISCUSSION

The isolations of EMC virus from the cotton rats is not too surprising con-

sidering the fact that other workers have found rats to: (1) possess neutralizing antibodies in nature and (2) to be susceptible to infection by EMC virus. The source of infection for the raccoon could have been from the ingestion of infected cotton rats. Kilham<sup>9</sup> showed that mongooses are susceptible to infection by the oral route, and speculated that they might become infected by predation upon field rats carrying EMC virus. The role, if any, raccoons may play in the continued transmission of EMC to domestic swine is unknown. But, isolation of the agent from an intestinal suspension of the raccoon indicates the raccoon could easily in the course of its daily activities, serve to maintain and disseminate EMC virus in nature.

The lack of presence of any detectable antibody in the serums of these animals yet the presence of virus in the intestine suggests: (1) that the infection was very recent, possibly from the infected water<sup>4</sup> on the premise, and was before antibodies had had a chance to be formed; (2) that the virus was just passing through the intestines of these wild animals without actual replication taking place. The titer of virus in the intestine of these animals was of a low level, near the fifty per cent end-point for mice<sup>4</sup>.

Table 1. Serum Neutralization Tests and EMC Virus Isolations on Samples Collected February 21-24, 1966\*, Winter Haven, Florida

| Host   | Serum Neutralizations |            |              | Virus Isolation Attempts |              |
|--|-----------------------|------------|--------------|--------------------------|--------------|
|  | No. Taken             | No. Tested | No. Positive | No. Tested               | No. Positive |
| Cotton Rat ( <i>Sigmodon hispidus</i> )          | 6                     | 4**        | 0            | 6                        | 2            |
| Rice Rat ( <i>Oryzomys palustris</i> )           | 2                     | 2          | 0            | 2                        | 0            |
| Opossum ( <i>Didelphis marsupialis</i> )         | 3                     | 3          | 0            | 3                        | 0            |
| Raccoon ( <i>Procyon lotor</i> )                 | 2                     | 2          | 0            | 2                        | 1            |
| Civet Cat ( <i>Spilogale ambervalis</i> )        | 1                     | 1          | 0            | 1                        | 0            |
| Slugs ( <i>Deroceros agriolimos</i> )            | 1                     | NA         | --           | 1                        | 0            |
| Boattailed Grackle ( <i>Cassidix mexicanus</i> ) | 1                     | 1          | 0            | NT                       | --           |
| Totals   | 16                    | 13         | 0            | 15                       | 3            |

NA - not applicable; NT - not tested

\* Portion of Table 3<sup>4</sup>.

\*\* Two were not tested due to insufficient serum.

### ACKNOWLEDGEMENT

The author expresses gratitude to Mr. G. W. Stroup for his technical assistance.

## LITERATURE CITED

1. DICK, G. W. A., 1963. Epidemiological notes on some viruses isolated in Uganda, Trans. Roy. Soc. Trop. Med. and Hyg. 47: 13-43.
  2. GAINER, J. H., 1961. Studies on the Natural and Experimental Infections of Animals in Florida with the Encephalomyocarditis Virus. From the 65th Annual Proceedings United States Livestock Sanitary Assoc. 556-572.
  3. GAINER, J. H., and MURCHISON, T. E., 1961. Encephalomyocarditis Virus Infection of Swine. Veterinary Medicine. 56: 173-175.
  4. GAINER, J. H., SANDEFUR, J. R., and BIGLER, W. J. High Mortality in a Florida Swine Herd Infected with the Encephalomyocarditis virus; An accompanying Epizootologic Survey. In press, Cornell Veterinarian.
  5. JUNGBLUT, C. W. and DALLDORF, G., 1943. Epidemiology and experimental observations on the possible significance of rodents in a suburban epidemic of poliomyelitis. Am. J. Pub. Hlth. 33: 169-172.
  6. JUNGBLUT, C. W. and SANDERS, M., 1940. Studies of a murine strain of poliomyelitis virus in cotton rats and white mice. J. Exper. Med. 72: 407-36.
  7. KILHAM, L., MASON, P., and DAVIES, J. N. P., 1955a. Host-virus relations in encephalomyocarditis (EMC) virus infections. I. Infections of wild rats. A. J. Trop. Med. and Hyg. 5: 647-654.
  8. KILHAM, L., MASON, P., and DAVIES, J. N. P., 1955b. Pathogenesis of fatal encephalomyocarditis (EMC) virus in albino rats. Proc. Soc. Exp. Biol. and Med. 90: 383-387.
  9. KILHAM, L., MASON, P., and DAVIES, J. N. P., 1955c. Host-virus relations in encephalomyocarditis (EMC) virus infections. II Myocarditis in mongooses. Amer. J. Trop. Med. and Hyg. 5: 655-663.
  10. KISSLING, R. E., VANELLA, J. M., and SCHAEFFER, M., 1956. Recent isolations of encephalomyocarditis virus. Proc. Soc. Exper. Biol. and Med. 91: 148-150.
  11. MURANE, T. G., CRAIGHEAD, J. E., MONDRAGON, H., and SHELOKOV, A., 1960. Fatal disease of swine due to encephalomyocarditis virus. Science 131: 498-499.
  12. POPE, J. H., and SCOTT, W., 1960. A survey for antibodies to EMC virus in man and animals. Aust. J. Exp. Biol. Med. Sci. 38: 447-450.
  13. WARREN, J., RUSS, S. B., and JEFFERIES, H., 1949. Neutralizing antibody against viruses of the encephalomyocarditis group in the sera of wild rats. Proc. Soc. Exper. Biol. and Med. 71: 276-378.
-