

Babesia sp. in Colombian bats (Microchiroptera)

Author: Marinkelle, Cornells J.

Source: Journal of Wildlife Diseases, 32(3) : 534

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-32.3.534>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***Babesia* sp. in Colombian bats (Microchiroptera)**

Cornelis J. Marinkelle, Centro de Investigaciones en Microbiología y Parasitología Tropical, Universidad de los Andes, Apartado Aéreo 4976, Santafé de Bogotá, Colombia

ABSTRACT: Two leaf-chinned bats (*Mormoops megalophylla*) collected in 1963 in central Colombia were heavily infected with *Babesia* sp., probably *Babesia vesperuginis*. Both bats had pronounced splenomegaly. This is the first report of a *Babesia* sp. infection of a bat in the Americas.

Key words: *Babesia* sp., leaf-chinned Bat, *Mormoops megalophylla*.

Babesia vesperuginis (Dionisi, 1899) has been reported from two European countries (Goedbloed et al., 1964; Gardner and Molyneux, 1987). In this paper I report for the first time a member of the genus *Babesia* from bats in the Americas.

Between 1961 and 1972 more than 20,000 Colombian bats were captured in 165 localities from all over Colombia and examined for parasites and fungi. The methods and sites of capture of the bats have been described by Grose and Marinkelle (1970) and Marinkelle (1982). Giemsa-stained thick and thin blood films were made of blood obtained by heart puncture. The preparations were examined microscopically for at least 15 min at 600 \times magnification. When bats were found to be positive for hematozoa impression, smears of lungs, hearts, livers, spleens, brains and kidneys were fixed with 100% methanol and stained with Giemsa and microscopically examined. After preparation of the smears, the organs were preserved in 10% formalin solution for histological sections.

The bats as well as their resting places were examined for the presence of ectoparasites. The cave where *Babesia* sp.-infected bats were found was searched extensively for ectoparasites during eight visits between June 1961 and April 1972. This cave is locally known as the Cueva de Macaregua and is located near the village of San Gil in Central Colombia (6°33'N, 73°08'W).

Only two of 168 leaf-chinned bats, (*Mormoops megalophylla*), both collected in November 1963, were found infected with *Babesia* sp. Both bats had a heavy infection with about 18% of the red blood cells infected.

The very young trophozoites were round or oval and measured 1.0 to 1.8 μ m. Small rings appeared to divide by binary fission. Large rings, 2 to 4 μ m in diameter, were numerous and had a central or marginal nucleus; sometimes two nuclei were present in one ring. The large rings had no evidence of division. Pear-shaped parasites of 1.4 to 2.0 μ m were not uncommon (Fig. 1). Some ameboid forms with delicate pseudopodia reached 6 μ m and contained a large irregular nucleus, 2.5 μ m in diameter, or three to four dots of chromatin. Cross-shaped (Maltese Cross) forms were found only twice. All intra-erythrocytic forms had a well-defined white central vacuole; exoerythrocytic parasites were observed infrequently. Pigment was absent and reproduction was without synchronicity, there was no evidence of parasite multiplication stages in the tissues smears. The brain capillaries were blocked by the parasites (Fig. 2).

The formalin-fixed spleens of the two infected bats weighed 360 and 388 mg, respectively; while the mean (\pm SE) weight of the preserved spleens of 10 other non-infected *M. megalophylla* was 44 ± 4.4 mg.

The piroplasms found were considered to be members of the genus *Babesia* because division occurred in the erythrocytes; no pigment was found, and no exoerythrocytic schizonts were found in the impression smears of the organs.

The morphology of the parasite seemed similar or identical to *Babesia vesperuginis*, as described and pictured from British bats




FIGURE 1. *Babesia* sp. (probably *B. vesperuginis*) in blood of *Mormoops megalophylla*. Giemsa. Bar = 5 μ m.




FIGURE 2. *Babesia* sp. (probably *B. vesperuginis*) in brain capillary of *Mormoops megalophylla*. H+E. Bar = 10 μ m.

(Gardner and Molyneux, 1987). Until other evidence becomes available. We speculate this parasite to be *B. vesperuginis*. The observed splenomegaly also was consistent with the observations of Gardner and Molyneux (1987). One blood smear of *Babesia* sp. probably *B. vesperuginis* is deposited in the U.S. National Parasite Collection, (Animal Parasitology Institute, Beltsville, Maryland, USA, voucher number 86005; collection date: 12 May 1966).

In Colombia, seven species of *Ornithodoros* spp. (Argasidae) have been found in association with bats, but hard ticks (Ixodidae) have not been reported from bats in Colombia (Marinkelle and Grose, 1981).

The only argasid ticks found in caves inhabited by *M. megalophylla* in Colombia were *Ornithodoros marinkellei* and *Antricola mexicanus* (Marinkelle and Grose, 1981; Marinkelle, 1982). *Antricola mexicanus* was the only tick found in Macaregua Cave, which was inhabited by thousands of *M. megalophylla*. Many *A. mexicanus* were found in association with *M. megalophylla* and its guano in the cave (Grose and Marinkelle, 1968).

The ticks were identified with the key of Cooley and Kohls (1944) and were deposited in the U.S. National Parasite Collection, voucher number: 84762. *Antricola mexicanus* has not been reported from

other parts of South America and is not known to transmit any pathogenic micro-organism. Argasid ticks previously have been found in the Netherlands on *Babesia* sp.-infected bats, (Goedbloed et al., 1964).

This work was supported by a grant from Colciencias, Colombia.

LITERATURE CITED

- COOLEY, R. A., AND G. M. KOHLS. 1944. The Argasidae of North America, Central America and Cuba. Monograph No. 1, American Midland Naturalist, University of Notre Dame, Notre Dame, Indiana, 152 pp.
- GARDNER, R. A., AND D. H. MOLYNEUX. 1987. *Babesia vesperuginis* natural and experimental infections in British bats (Microchiroptera). Parasitology 95: 461–469.
- GOEDBLOED, E., H. CREMERS-HOYER, AND N. M. PERIE. 1964. Blood parasites of bats in the Netherlands. Annals of Tropical Medicine and Parasitology 58: 257–260.
- GROSE, E. S., AND C. J. MARINKELLE. 1968. *Candida chiropterorum* n. sp. from Colombian bats. Mycopathologia et Mycologia Applicata 36: 225–227.
- , AND ———. 1970. Biospeleology of the Macaregua Cave (Colombia). Mitteilungen aus dem Instituto Aleman de Investigaciones Cientificas-Punta Betin (Colombia) 4: 11–13.
- MARINKELLE, C. J. 1982. Prevalence of *Trypanosoma cruzi*-like infections of Colombian bats. Annals of Tropical Medicine and Parasitology 76: 125–134.
- , AND E. S. GROSE. 1981. A list of ectoparasites of Colombian bats. Revista de Biología Tropical 29: 11–20.

Received for publication 4 January 1995.