

## **Relationship between Gonadal Steroids and Corticosterone during Blood Sampling in Saker Falcons**

Author: Al-Ankari, Abdul-Rahman S.

Source: Journal of Wildlife Diseases, 34(3) : 653-655

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-34.3.653>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Relationship between Gonadal Steroids and Corticosterone during Blood Sampling in Saker Falcons

**Abdul-Rahman S. Al-Ankari** Department of Clinical Studies, College of Veterinary Medicine and Animal Resources, King Faisal University, P.O. Box 3498, Hofuf, Al-Ahsa 31982, Saudi Arabia.

**ABSTRACT:** Blood sampling in manually restrained or ketamine (15 mg/kg given intramuscularly) treated saker falcons (*Falco cherrug*) induced an increased concentration in plasma corticosterone. Elevated plasma progesterone, oestradiol 17 $\beta$ , and testosterone concentrations also were observed in some of these birds. An inverse relationship was demonstrated between levels of corticosterone and progesterone, but not with the levels of other hormones. It is suggested that progesterone measurement should be taken into consideration when studying the influence of stressors in falcons.

**Key words:** Blood sampling, corticosterone, *Falco cherrug*, saker falcons, gonadal steroids, stressors.

Blood collection for endocrinological analysis allows monitoring many physiological, nutritional or disease conditions in birds of prey. Light sedation with ketamine can safely restrain falcons for collection of a blood sample (Samour and D'Aloia, 1996). Although ketamine sedation has many helpful features (Bywater, 1990), there are some problems including its effects on metabolism and the endocrine system (Chiasson et al., 1973; Hellgren et al., 1985; Lumeij, 1993). These authors have reported that poultry anesthetized with ketamine exhibit reduced adrenal weight suggesting some physiological changes in adrenal function. This study was conducted to investigate the relationship of blood concentration of reproductive steroids during ketamine sedation and manual restraint of saker falcons (*Falco cherrug*).

Jugular blood samples (0.5 ml) from 25 clinically normal adult female saker falcons were obtained by venipuncture using 23 gauge needles. The birds were admitted to Salman Falcon Hospital (Al-Areen Wildlife Park, Manama, Bahrain) for routine checks during hunting season (October to

February, 1995). Birds were housed singly in a large fiberglass cages located in a closed room that was maintained with 12L:12D photoperiod. Feeding and cage cleaning were the only direct contacts by humans with the birds. Falcons were sampled while either manually restrained (Group 1, n = 12) by holding the two unfolded wings by an assistant or while under light sedation (Group 2, n = 13) using 15mg/kg of ketamine hydrochloride (Vetalar, Parke Davis Ltd, Pontypool, UK) administered intramuscularly. Blood samples were collected into heparinized tubes, then centrifuged at 3,000 rpm for 10 min and plasma was separated and stored at -20 C until analysis.

Progesterone (P), oestradiol 17 $\beta$  (E<sub>2</sub>), testosterone (T), and corticosterone (C) were measured by radioimmunoassay methods previously validated and described (Harvey et al., 1981; Homeida, 1986; Homeida et al., 1988). The specificities and cross reactions of the antisera used have been reported elsewhere; the respective intra- and interassay coefficients of variation were 7 (n = 10) and 10% for P, 6 (n = 11) and 10% for E<sub>2</sub>, 9 (n = 11) and 10% for T, and 9 (n = 9) and 10% for C. Extraction efficiencies were 87% for P, 89% for E<sub>2</sub>, 75% for T, and 83% for C (Dobson and Dean, 1974; Harvey et al., 1981; Homeida and Al-Afaliq, 1994). Results were corrected for recovery values.

Data are given for each bird in Table 1. Pearson's correlation coefficients were calculated to determine the association between corticosterone and other hormones (Tallarida and Murray, 1981).

Blood sampling increased levels of C (Table 1) in plasma of five ketamine-treated birds (42%) and five manually restrained falcons (38%). Activation of the

TABLE 1. Peripheral plasma concentration of corticosterone (C), progesterone (P), oestradiol 17 $\beta$  (E<sub>2</sub>), and testosterone (T) in manually restrained (Group 1) or ketamine treated (Group 2) saker falcons in Bahrain.

Bird number	Group 1				Bird number	Group 2			
	C (ng/ml)	P (pg/ml)	E <sub>2</sub> (pg/ml)	T (pg/ml)		C (ng/ml)	P (pg/ml)	E <sub>2</sub> (pg/ml)	T (pg/ml)
1	8.0	1,300*	150*	220*	13	23.4*	270	120*	350*
2	22.3*	220	120*	120	14	26.1*	250	85	290*
3	25.1*	260	80	250*	15	6.5	1,250*	120*	160
4	7.5	1,200*	140*	270*	16	7.5	1,200*	130*	320*
5	21.3*	230	140*	350*	17	8.2	1,300*	70	310*
6	9.2	1,100*	130*	320*	18	9.1	1,220*	140*	290*
7	9.5	1,150*	60	360*	19	7.6	1,400*	160*	300*
8	10.1	1,200*	125*	370*	20	7.1	1,100*	120*	325*
9	8.2	1,350*	120*	250*	21	6.8	1,150*	50	115
10	24.6*	270	140*	130	22	20.1*	210	150*	320*
11	7.5	1,150*	130*	315*	23	22.6*	240	120*	170
12	19.3*	350	140*	290*	24	20.5*	260	65	330*
					25	7.2	1,250*	120*	320*

\* Significantly different from other birds in the group at  $P < 0.001$ .

pituitary-adrenal axis of birds in response to stress like bleeding is well established and is generally reflected by increased concentrations of C in plasma of the peripheral circulation of birds (Holmes and Philips, 1976; Harvey et al., 1980). Handling and blood sampling are known to be stressful in birds like hawks and guinea fowl (Mays et al., 1991; Cooper, 1991; Cooper et al., 1996). Ketamine sedation has no effect on C response to venipuncture in falcons. Similar results were observed in goshawks, pigeons, monkeys and deer sedated with ketamine (Wessen et al., 1979; Fuller et al., 1984; Lumeij, 1993).

Elevated concentrations of P, E<sub>2</sub>, and T were observed in both Group 1 and Group 2 falcons (Table 1) suggesting an active ovarian function in some of these birds (Hartelendy et al., 1993). No relation between C and E<sub>2</sub> ( $r = 0.17$ ,  $n = 25$ ,  $P > 0.1$ ) or between C and T ( $r = 0.06$ ,  $n = 25$ ,  $P > 0.1$ ) could be demonstrated. In past studies injections of oestradiol benzoate and testosterone propionate did not alter plasma concentration of C or luteinizing hormone in chickens (Wilson and Cunningham, 1980). An inverse ( $r = -0.45$ ,  $n = 25$ ,  $P < 0.001$ ) relationship was shown to occur between levels of P and C (Table 1) suggesting that P may inhibit the rise in

C concentration. It is shown that injection of P may suppress the pituitary-adrenal system and thus modulate the pattern of C secretion in chicken (Wilson and Cunningham, 1980) due to handling and bleeding. Thus, it appears that P measurement should be taken into consideration when studying the influence of stressors in saker falcons.

I thank A. M. Homeida for reading the manuscript, and A. M. Al-Awadi, A. Ashour, and D. O. Diab for technical assistance.

#### LITERATURE CITED

- BYWATER, R. J. 1990. Sedatives, tranquilizers and narcotics. In *Veterinary applied pharmacology and therapeutics*, G. C. Brander, D. M. Pugh and R. T. Bywater (eds.). Balliere Tindall, London, UK, pp. 289–299.
- CHIASSON, R. B., A. S. EGGE, AND B. LUNCH JR. 1973. The effect of ketamine on the adrenal gland of young white leghorn cockerels. *Poultry Science* 52: 1014–1018.
- COOPER, J. E. 1991. Caged and wild birds. In *Practical animal medicine*, R. C. Anderson and A. T. B. Edney (eds.). Pergamon Press, Oxford, UK, pp. 28–34.
- , R. A. MAX AND G. K. MBASSA. 1996. Health studies on a group of captive helmeted guinea fowl (*Numida meleagris*) in Tanzania. *Avian Pathology* 25: 135–145.
- DOBSON, H., AND P. D. G. DEAN. 1974. Radioimmunoassay of testosterone, oestradiol 17 $\alpha$  and

- oestradiol 17 $\beta$  in bovine plasma during oestrous cycle and last stages of pregnancy. *Journal of Endocrinology* 61: 479–486.
- FULLER, G. B., W. C. HOBSON, F. C. REYES, J. S. D. WINTER, AND C. FARMAN. 1984. Influence of restraint and ketamine anaesthesia on adrenal steroids, progesterone and gonadotropins in rhesus monkeys. *Proceedings of the Society of Experimental and Biological Medicine* 175: 487–490.
- HARTELENDY, F., M. JAMALUDDIN, AND M. MOLNAR. 1993. Signal transduction in avian granulosa cells. In *Avian Endocrinology*, P. J. Sharp (ed.). Burgess Science Press, Bristol, UK, pp. 331–341.
- HARVEY, S., E. BEDRACK, AND A. CHADWICK. 1981. Serum concentrations of prolactin, luteinizing hormone, growth hormone, corticosterone, progesterone, testosterone and oestradiol in relation to broodiness in domestic turkeys (*Meleagris gallopavo*). *Journal of Endocrinology* 89: 187–195.
- , B. J. MERRY, AND J. G. PHILIPS. 1980. Influence of stress on secretion of corticosterone in the duck (*Anas platyrhynchos*). *Journal of Endocrinology* 87: 161–171.
- HELLGREN, E. C., R. L. LOCHMILLER, M. S. AMOSS, AND W. E. GRANT. 1985. Endocrine and metabolic responses of the Collared peccary (*Tayassu tajacu*) to immobilization with ketamin hydrochloride. *Journal of Wildlife Diseases* 21: 417–425.
- HOLMES, W. N., AND J. G. PHILLIPS. 1976. The adrenal cortex of birds. In *General comparative and clinical endocrinology of the adrenal cortex*, I. Chester-Jones and I. W. Henderson (eds.). Academic Press, New York, New York, pp. 293–420.
- HOMEIDA, A. M. 1986. Use of spironolactone to investigate the role of testosterone secretion during luteolysis in the goat. *Journal of Reproduction and Fertility* 76: 153–157.
- , AND A. I. AL-AFALIQ. 1994. Delayed luteolysis and suppression of testosterone secretion after recombinant ovine interferon treatment in goats (*Capra hircus*). *Journal of Reproduction and Fertility* 102: 39–42.
- , M. G. R. KHALIL, AND A. A. M. TAHA. 1988. Plasma concentrations of oestrogen, testosterone and LH-like activity during the oestrous cycle of the camel (*Camelus dromedarius*). *Journal of Reproduction and Fertility* 83: 593–598.
- LUMEIJ, J. T. 1993. Effect of Ketamine-Xylazine anaesthesia on adrenal function and cardiac conduction in goshawks and pigeons. In *Raptor biomedicine*, P. T. Redig, J. E. Cooper, J. D. Remple and D. B. Hunter (eds.). Chiron Publication Ltd., Keighley, UK, pp. 145–149.
- MAYS, N. A., C. M. VLECK, AND J. DAWSON. 1991. Plasma luteinizing hormone, steroid hormones, behavioral role and nest stage in cooperatively breeding harris hawks (*Parabuteo unicinctus*). *Auk* 108: 619–637.
- SAMOUR, J. H., AND MARIE-ANN D'ALOIA. 1996. Normal blood chemistry of the saker falcon (*Falco cherrug*). *Avian Pathology* 25: 175–178.
- TALLARIDA, R. J., AND R. B. MURRAY. 1981. In *Manual of pharmacologic calculations with computer programs*, R. J. Tallarida and R. B. Murray (eds.). Springer-Verlag, New York, New York, 147 pp.
- WESSEN, J. A., P. F. SCANLON, R. L. KIRKPATRICK, AND H. S. MOSBY. 1979. Influence of chemical immobilization and physical restraint and packed cell volume, total protein, glucose and blood urea nitrogen in blood of white-tailed deer. *Canadian Journal of Zoology* 56: 756–767.
- WILSON, S. C., AND F. J. CUNNINGHAM. 1980. Concentrations of corticosterone and luteinizing hormones in plasma during the ovulatory cycle of the domestic hen and after the administration of gonadal steroids. *Journal of Endocrinology* 85: 209–218.

Received for publication 26 June 1997.