



## **PHENYLKETONURIA IN A MULE DEER (*Odocoileus hemionus*) 1**

Author: STUDIER, E. H.

Source: Bulletin of the Wildlife Disease Association, 4(4) : 143

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-4.4.143>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

The examination of Giemsa-stained blood smears from deer suggests that this disease is enzootic in East Texas and may be an important cause of mortality of deer.




Figure 1. *Babesia cervi* (N. sp.) within an erythrocyte.

H. R. EMERSON,  
*Department of Veterinary Pathology*  
*Texas A&M University*  
*College Station, Texas*  
 and W. T. WRIGHT,  
*Texas Parks and Wildlife Department*  
*Silsbee, Texas*  
 August 14, 1968

#### PHENYLKETONURIA IN A MULE DEER (*Odocoileus hemionus*)<sup>1</sup>

Potential space fluids and urine samples were taken from several mule deer collected in Northeast New Mexico during the spring, 1968. A urine sample, collected by manual expression of the bladder of a three year old buck, was tested for phenylpyruvic acid using Phenistix Reagent Strips (Ames Company, Inc., Elkhart, Ind.) and gave a positive test of more than 100 mg/100 ml urine. Urine samples from three other deer were tested with negative results. To our knowledge, this is the first reported case of phenylketonuria in mule deer. The buck suffering from this inherited metabolic disorder showed no obvious behavioral features indicative of the mental retardation associated with phenylketonuria in humans, and it had survived a relatively long time under natural conditions. Its weight, 105 lbs., compares favorably with the 97 and 110 lb. weights of two other adult bucks collected in the same area, indicating that it had developed normally despite its supposed handicap. The generally low weights of these three deer are indicative of the poor browse available in the collection area.

E. H. STUDIER  
*Department of Biology,*  
*New Mexico Highlands University*  
 Las Vegas, New Mexico 87701

W. G. EWING  
*Department of Zoology,*  
*Oregon State University*  
 Corvallis, Oregon 97331  
 July 17, 1968

<sup>1</sup> This study was supported by the Institute of Scientific Research, New Mexico Highlands University.