

Thallium Contamination in Wild Ducks in Japan

Authors: Mochizuki, Mariko, Mori, Makoto, Akinaga, Mayumi, Yugami, Kyoko, Oya, Chika, et al.

Source: Journal of Wildlife Diseases, 41(3) : 664-668

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-41.3.664>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Thallium Contamination in Wild Ducks in Japan

Mariko Mochizuki,^{1,2} Makoto Mori,³ Mayumi Akinaga,¹ Kyoko Yugami,¹ Chika Oya,⁴ Ryo Hondo,¹ and Fukiko Ueda^{1,5} ¹ Department of Veterinary Public Health, Nippon Veterinary and Animal Science University, Tokyo, Japan; ² Current address: Department of Veterinary Nursing, Nippon Veterinary and Animal Science University, Tokyo, Japan; ³ Department of Applied Biological Chemistry, Faculty of Agriculture, Shizuoka University, Shizuoka, Japan; ⁴ Department of Biomedical Science, Graduate School of Agricultural and Life Sciences, University of Tokyo, Tokyo, Japan; ⁵ Corresponding author (email: fueda@nvau.ac.jp)

ABSTRACT: Although thallium (Tl) is toxic to both humans and animals, there is little information on contamination in wildlife. In this study, Tl contents in wild ducks in Japan were determined. Contents of Tl in kidney and liver ranged from 0.42 to 119.61 and 0.10 to 33.94 $\mu\text{g/g}$ dry weight, respectively. Significant correlations between Tl contents in kidney and liver were observed for all dabbling ducks except mallard (*Anas platyrhynchos*); similar correlations were not observed in diving ducks. Variation in Tl content was observed between sampling locations with the highest mean Tl content in the Eurasian wigeon (*Anas penelope*) collected in Ibaraki Prefecture.

Key words: Duck, thallium, wild bird.

Thallium (Tl) exists naturally in the environment (Mulkey and Oehme, 1993; Asami, 2001), and background levels of <0.00001 mg/l have been reported for fresh water and seawater (Mason, 1966). Concentrations in nonpolluted soil of 0.10–0.56 mg/kg dry weight (wt.) (Asami et al., 1996), 0.07–0.91 mg/kg dry wt. (Hoffer et al., 1990), and 0.292–1.172 mg/kg dry wt. (Qi et al., 1992) have been reported. In plants, concentrations do not generally exceed 0.05 mg/kg dry wt. (Wierzbicka et al., 2004) and, in animals, Tl levels are normally less than 1 ppb and 10 ppb in blood and tissues, respectively (Mulkey and Oehme, 1993). In humans, Tl has been detected in both kidney and liver, but concentrations were less than 4.05 and 1.42 ng/g wet wt., respectively (Weinig and Zink, 1967).

Although rodenticides and insecticides containing Tl have been regulated in many countries since the 1960s and 1970s (Asami, 2001), coal-burning power plants, certain cement plants, and mining and smelting operations can represent sources for this element (Pielow, 1979; Prinz et al.,

1979; Sabbioni et al., 1984; Ewers, 1988). Nriagu and Pacyna (1988) estimated that the total emission of Tl from coal combustion and cement production ranged from 3,320 to $6,950 \times 10^3$ kg/yr. This metal and its compounds are also used in various industrial products, such as the production of sulfuric acid, dye and pigments, semi-conductors, and superconducting ceramics (Mulkey and Oehme, 1993; Asami, 2001). Thallium has been suggested as a priority elemental pollutant with regard to human health (Keith and Telliard, 1979), and a significant positive correlation between the Tl content of moss and the incidence of circulatory disease in humans has been reported (Heim et al., 2002).

With the exception of reports of acute poisoning due to rodenticides and insecticides (Cromartie et al., 1975; Clausen and Karlog, 1977), there is little information on Tl contamination in wildlife. In this study, we determined levels of Tl in several species of wild ducks collected in various locations in Japan.

Samples were collected from 58 dabbling ducks representing five species and 15 diving ducks representing three species (Fig. 1). Dabbling ducks included spotbill ducks (*Anas poecilorhyncha*, $n=19$), mallards (*Anas platyrhynchos*, $n=7$), common teal (*Anas crecca*, $n=6$), northern pintails (*Anas acuta*, $n=11$) and Eurasian wigeon (*Anas penelope*, $n=15$). Diving ducks included greater scaup (*Aythya marila*, $n=6$), tufted ducks (*Aythya fuligula*, $n=6$) and pochards (*Aythya ferina*, $n=3$). Ducks were captured between 1993 and 1995 as a part of unrelated projects being conducted by the Japanese Ministry of the Environment. Other birds were supplied




FIGURE 1. Sampling locations for wild ducks.

through the Gyotoku Bird Observatory in Chiba Prefecture.

Samples (approximately 200 mg) of kidney and liver from each duck were dried, weighed, and digested. Thallium contents were determined using an inductively coupled plasma atomic emission spectrometer (ICP-AES, Spectro A.I., Germany) as described in the previous reports (Mochizuki et al., 1999, 2002a, b, c). The detection limit for Tl was 0.007 ppm, and for statis-

tical analyses, nondetectable values were estimated to contain half of the detection limit. Results are presented as mean \pm standard error (SE). Pearson's correlation coefficients were calculated in Excel[®] (Microsoft Corporation, Redmond, Washington, USA).

Thallium contents in kidney and liver for each species are shown in Table 1. Mean contents of Tl in kidney were greater than 4 $\mu\text{g/g}$ dry wt. in all dabbling spe-

TABLE 1. Thallium contents ($\mu\text{g/g}$ dry wt.) in the kidney and liver of each species and the correlation between thallium contents of kidney and that of liver. The correlation with an asterisk shows significant correlation (** $P < 0.01$, * $P < 0.05$). n = number of samples.

Species	n	Kidney		Liver		Correlation
		Mean \pm SE	Range	Mean \pm SE	Range	
Dabbling ducks						
Spotbill duck	19	7.15 \pm 1.48	(0.91–23.73)	4.57 \pm 1.32	(0.60–26.62)	0.601**
Mallard	7	8.11 \pm 1.96	(2.65–15.76)	4.50 \pm 1.42	(1.09–11.86)	0.692
Teal	6	10.68 \pm 6.18	(2.12–40.33)	5.32 \pm 2.91	(0.37–19.12)	0.881**
Pintail	11	4.00 \pm 1.35	(1.01–15.70)	6.52 \pm 3.21	(0.12–33.94)	0.692*
Wigeon	15	20.38 \pm 8.08	(0.42–119.61)	4.27 \pm 1.14	(0.48–14.72)	0.848**
Total	58	10.46 \pm 2.34	(0.42–119.61)	4.93 \pm 0.85	(0.12–33.94)	0.411**
Diving ducks						
Scaup	6	2.10 \pm 0.32	(1.05–2.98)	2.71 \pm 1.15	(0.43–7.85)	0.214
Tufted duck	6	3.04 \pm 0.82	(1.02–6.60)	5.27 \pm 3.83	(0.10–24.33)	0.810
Pochard	3	2.13 \pm 0.33	(1.63–2.75)	2.09 \pm 0.85	(0.42–3.21)	–0.019
Total	15	2.48 \pm 0.36	(1.02–6.60)	3.61 \pm 1.56	(0.10–24.23)	0.745**

TABLE 2. Thallium contents ($\mu\text{g/g}$ dry wt.) in the kidney and liver of each prefecture. n =number of samples.

Prefecture	Dabbling ducks			Diving ducks		
	n	Kidney	Liver	n	Kidney	Liver
Akita	11	8.43 \pm 2.05	5.73 \pm 2.32			
Chiba	12	3.32 \pm 0.78	2.83 \pm 0.70	13	2.28 \pm 0.22	2.04 \pm 0.59
Fukushima	9	4.08 \pm 1.56	6.93 \pm 3.94			
Ibaraki	10	28.25 \pm 11.44	5.26 \pm 1.55			
Ishikawa	5	14.73 \pm 6.74	6.96 \pm 3.13	1	6.60	24.23
Miyagi	9	7.80 \pm 2.15	3.99 \pm 0.74			
Sizuoka	1	8.71	1.93			
Tokyo	1	2.18	1.22	1	1.02	3.32

cies, while those in diving ducks were less than 3 $\mu\text{g/g}$ dry wt. for all species. The total mean value for dabbling duck species (10.5 $\mu\text{g/g}$ dry wt.) was approximately four times higher than the mean for diving duck (2.5 $\mu\text{g/g}$ dry wt.). Although Tl levels in liver also were higher in species of dabbling ducks, mean values were less than 6.5 $\mu\text{g/g}$ dry wt. in all species, and these differences were not statistically significant.

The Tl levels in ducks in this study were lower than those reported from birds of prey (63 ppm wet wt.) that died due to acute poisoning by Tl (Cromartie et al., 1975), and wood mice (kidney; 44.05, liver; 11.34 $\mu\text{g/g}$ dry mass) and magpies (maximum value of kidney; 45 $\mu\text{g/g}$ dry wt.) captured in a polluted area next to a zinc mine (Dmowski et al., 1998). Thallium contents in kidney and liver exceeding 0.5 ppm wet wt. (approximately 2 $\mu\text{g/g}$ dry wt.) are believed to be indicative of poisoning (Clausen and Karlog, 1977). Mean contents of Tl observed for all species of ducks in this study (Table 1) exceeded this proposed threshold. There was no indication, however, that any of the ducks sampled in this study were physically affected.

Mochizuki et al. (1999, 2002a, c) reported spatial variation in contents of cadmium, molybdenum, and vanadium in wild birds, suggesting that birds may represent effective indicators of environmental contamination. This same relationship may exist with Tl contents in ducks, which approximated 10 $\mu\text{g/g}$ dry wt. in all pre-

fectures except Chiba Prefecture (Table 2). In Japan, high Tl contents have been reported in plants (28.9 mg/kg dry wt.) and bottom sediment (79.9 mg/kg dry wt.) associated with the Hosokura mines in Miyagi Prefecture (Asami, 2001). Background concentrations of 4.81 mg/kg dry wt. in sediments collected 20 km from this point source have also been reported. A high Tl content (maximum, 4.13 mg/kg dry wt.) has also been reported for one area near the copper mine in Ibaraki Prefecture and the maximum/background ratio of 10.7 was reported from soil near the mine. Although the authors could not find similar reports in other prefectures, the prefectures of Akita, Fukushima, and Ishikawa have active and idle silver, copper, lead, and zinc mines (Hata, 1997; Asami, 2001) that have been suggested as possible sources of Tl (Ewers, 1988). There are no such mines in the Chiba Prefecture, which may explain the low levels of Tl observed in ducks sampled from this location (Table 2).

With cadmium (Mochizuki et al., 2002a), vanadium (Mochizuki et al., 1999), and lead (Gerhardsson et al., 1995), a correlation between the contents observed in kidney and liver has been reported in wildlife species and humans. Similar correlations for Tl were obtained from all dabbling species except for the mallard and all diving species in the present study (Table 1). Sample sizes for individual species by location were too low for a meaningful comparison of kidney and liver Tl con-




FIGURE 2. Relation between the Tl contents ($\mu\text{g/g}$ dry wt.) in kidney and liver of Eurasian wigeon sampled from Ibaraki Prefecture.

tents. However, a significant correlation was obtained with the results from Eurasian wigeon ($n=9$) from Ibaraki Prefecture ($R=0.856$, $P<0.01$) (Fig. 2). Results of this study suggest that elevated contents of Tl are common in ducks in Japan but the potential clinical significance of these Tl levels is not understood. Additional monitoring of waterfowl may provide a means to better understand potential sources of Tl to wildlife populations.

The authors express their thanks to H. Kajigaya at Nippon Veterinary and Animal Science University, H. Kobayashi of Japanese Ministry of the Environment, and S. Hasuo and T. Sato of the Gyoutoku Bird Observatory for the supply of wild ducks. This study was supported by grant 13660328 in 2001–2003 from the Japanese Ministry of Education.

LITERATURE CITED

- ASAMI, T. 2001. *Nihon dojyo no yugaikinzoikuosen* [Toxic metal pollution in soils in Japanese soil]. AGNE Gijyutu Center, Tokyo, Japan. [in Japanese.]
- , C. MIZUI, T. SHIMADA, AND M. KUBOTA. 1996. Determination of thallium in soils by flame atomic absorption spectrometry. *Fresenius Journal of Analytical Chemistry* 356: 348–51.
- CLAUSEN, B., AND O. KARLOG. 1977. Thallium loading in owls and other birds of prey in Denmark. *Nordic Veterinary Medicine* 29: 227–231.
- CROMARTIE, E., W. L. REICHEL, L. N. LOCKE, A. A. BELISLE, T. E. KAISER, T. G. LAMONT, B. M. MULHERN, R. M. PROUTY, AND D. M. SWINFORD. 1975. Residues of organochlorine pesticides and polychlorinated biphenyls and autopsy data for bald eagles, 1971–1972. *Pesticide Monitoring* 9: 11–14.
- DMOWSKI, K., A. KOZAKIEWICZ, AND M. KOZAKIEWICZ. 1998. Small mammal populations and community under conditions of extremely high thallium contamination in the environment. *Ecotoxicology and Environmental Safety* 41: 2–7.
- EWERS, U. 1988. Environmental exposure to thallium. *The Science of the Total Environment* 71: 285–92.
- GERHARDSSON, L., V. ENGLYST, N. G. LUNDSTROM, G. NORDBERG, S. SANDBERG, AND F. STEINVALL. 1995. Lead in tissues of deceased lead smelter workers. *Journal Trace Element Medical Biology* 9: 136–143.
- HATA, A. 1997. *Kinzokusangyou no gijyutu to kogai* [Technology and public hazards of metal industry]. AGNE Gijyutu Center, Tokyo, Japan. [in Japanese.]

- HEIM, M., O. WAPPELHORST, AND B. MARKERT. 2002. Thallium in terrestrial environments—occurrence and effects. *Ecotoxicology* 11: 369–377.
- HOFFER, G. F., K. AICHBERGER, AND U. S. HOCHMAIR. 1990. Thalliumgehalte landwirtschaftlich genutzter Böden Oberösterreichs. *Die Bödenkultur* 41: 187–193.
- KEITH, L. H., AND W. A. TELLIARD. 1979. Priority pollutants. *Environmental Science Technology* 13: 416–423.
- MASON, B. 1966. *Principles of geochemistry*. 3rd Edition, New York: John Wiley and Sons, Inc. [Translated in Japanese, Iwamami Publishers.]
- MOCHIZUKI, M., F. UEDA, S. SASAKI, AND R. HONDO. 1999. Vanadium contamination and the relation between vanadium and other elements in wild birds. *Environmental Pollution* 106: 249–251.
- , R. HONDO, K. KUMON, R. SASAKI, H. MATSUBA, AND F. UEDA. 2002a. Cadmium contamination in wild birds as an indicator of environmental pollution. *Environmental Monitoring and Assessment* 73: 229–235.
- , ———, AND F. UEDA. 2002b. Simultaneous analysis for multiple heavy metals contaminated biological samples. *Biological Trace Element Research* 87: 211–223.
- , R. SASAKI, Y. YAMASHITA, M. AKINAGA, A. ANAN, S. SASAKI, R. HONDO, AND F. UEDA. 2002c. The distribution of molybdenum in the tissues of wild ducks. *Environmental Monitoring and Assessment* 77: 155–161.
- MULKEY, J. P., AND F. W. OEHME. 1993. A review of thallium toxicity. *Veterinary Human Toxicology* 35: 445–453.
- NRIAGU, J. O., AND M. PACYNA. 1988. Quantitative assessment of worldwide contamination of air, water and soils by trace metals. *Nature* 33: 134–139.
- PIELOW, E. 1979. Umweltschäden durch Thallium in der Umgebung eines Zementwerkes. *Umwelt* 5: 394–396.
- PRINZ, B., G. H. M. KRAUS, AND H. STRATMANN. 1979. Thalliumschäden in der Umgebung der Dyckerhoff Zementwerke AG in Lengerich, Westfalen. *Staub-Reinhalt Luft* 39: 457–462.
- QI, W., Y. CHEN, AND J. CAO. 1992. Indium and thallium background contents in soils of China. *International Journal of Environmental Studies* 40: 311–315.
- SABBIONI, E., L. GOETZ, AND G. BIGNOLI. 1984. Health and environmental implications of trace metals released from coal-fired power plants: An assessment study of the situation in the European Community. *The Science of the Total Environment* 40: 141–154.
- WEINIG, E., AND P. ZINK. 1967. Über die quantitative massenspektrometrische Bestimmung des normalen Thallium-Gehalts in menschlichen Organismus. *Archiv für Tokikologie* 22: 255–274.
- WIERZBICKA, M., G. SZAREK-LUKASZEWSKA, AND K. GRODZINSKA. 2004. Highly toxic thallium in plants from the vicinity of Olkusz (Poland). *Ecotoxicology and Environmental Safety* 59: 84–88.

Received for publication 1 March 2004.