



## **Seroprevalence of Antibodies to *Toxoplasma gondii* in the Pennsylvania Bobcat (*Lynx rufus rufus*)**

Authors: Mucker, Eric M., Dubey, J. P., Lovallo, Matthew J., and Humphreys, Jan G.

Source: Journal of Wildlife Diseases, 42(1) : 188-191

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-42.1.188>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Seroprevalence of Antibodies to *Toxoplasma gondii* in the Pennsylvania Bobcat (*Lynx rufus rufus*)

Eric M. Mucker,<sup>1,4</sup> J. P. Dubey,<sup>2</sup> Matthew J. Lovallo,<sup>3</sup> and Jan G. Humphreys<sup>1</sup> <sup>1</sup> Indiana University of Pennsylvania, Department of Biology, Weyandt Hall, Indiana, Pennsylvania 15705, USA; <sup>2</sup> United States Department of Agriculture, Agricultural Research Service, Animal and Natural Resources Institute, Animal Parasitic Diseases Laboratory, Building 1001, Beltsville, Maryland 20705-2350, USA; <sup>3</sup> Pennsylvania Game Commission, 2001 Elmerton Avenue, Harrisburg, Pennsylvania 17110-9797, USA; <sup>4</sup> Corresponding author (email: Eric.Mucker@det.amedd.army.mil)

**ABSTRACT:** From 2000 to 2002 bobcat blood samples were collected, in association with the Pennsylvania Game Commission, during the recently reactivated bobcat hunting and trapping season. Sex, age, and county/township data were recorded for each animal. Blood was tested for antibodies to *Toxoplasma gondii* using the modified agglutination test. In the 2-yr study, 131 bobcat samples were collected in 14 Pennsylvania counties and 109 (83%) of these had antibodies to *T. gondii* (titer $\geq$ 25). A two-way Chi-Square test (95% confidence interval) yielded no significance differences in antibody prevalence between males (83%) and females (88%) or adults (83%) and juveniles (77%). All 14 counties had at least one bobcat with antibodies to *T. gondii*.

**Key words:** Blood collection strips, bobcat, *Lynx rufus*, modified agglutination test, seroprevalence, *Toxoplasma gondii*.

*Toxoplasma gondii* is an intracellular protozoan that infects animals and humans. The life cycle of *T. gondii* involves asexual reproduction in intermediate hosts and both asexual and sexual reproduction in the definitive host (Felidae). The infectious product of sexual reproduction, the oocyst, is shed in the feces of infected felids during the acute phase of infection that can persist for 2 wk, during which thousands of oocysts can be produced from a single infected animal (Dubey and Beattie, 1988). Intermediate hosts become infected by consuming the oocyst and/or consuming another infected host. Humans can act as an intermediate host, but infection is rarely fatal unless the person is immunocompromised (Dubey and Beattie, 1988).

Excluding feral cats, the bobcat (*Lynx rufus rufus*) is the only wild felid in Pennsylvania. Therefore, the bobcat is

a potentially important component for the propagation and perpetuation of *T. gondii* in Pennsylvania's sylvatic cycle and, consequently, a good indicator of the prevalence of *T. gondii* in the wild. For these reasons, we evaluated the prevalence of antibodies to *T. gondii* in bobcats.

Pennsylvania Game Commission (PGC) personnel collected blood samples from bobcats harvested during 2000–2001 and 2001–2002. Harvesting was restricted to Furbearer Management Zones 2 and 3, which included 20 counties in north-central and northeastern Pennsylvania. The PGC also collected blood samples from road-killed bobcats. Nobuto blood collecting strips Type I (Toyo Roshi Kaisha, Ltd; distributed by Advantec MFS, Inc., Dublin, California, USA) were mailed to field personnel along with an information packet, which included the reasons for conducting the study and directions for properly acquiring, drying, and mailing the Nobuto strips back to the Indiana University of Pennsylvania (IUP) for further processing and analysis.

Each bobcat was examined within 4 days after death by the PGC. Blood was taken immediately by using Nobuto strips if the carcass was fresh. Bobcats were frozen if they could not be examined immediately, then later thawed and blood samples taken. Sex, age (cementum-annulus method of age determination), location, and tag number were recorded for all animals. The strips were refrigerated until they could be further processed.

Nobuto strips were cut into three or four segments at IUP and placed into a 1.5 ml centrifuge tube. One milliliter of

phosphate buffered saline (pH 7.4; Sigma Diagnostics, St. Louis, Missouri, USA) was added to each tube, diluting the blood 1:25. After 2 hr, the supernatant was removed, centrifuged, and stored at 4 C until tested for antibodies to *T. gondii* using the modified agglutination test (MAT) as described (Dubey and Desmonts, 1987). Antibody testing was performed at the Animal Parasitic Diseases Laboratory, US Department of Agriculture, Beltsville, Maryland, USA. Blood samples were further diluted to provide 1:50 and 1:500 dilutions, and a 1:25 dilution was used as the positive threshold dilution as described (Dubey and Beattie, 1988). A two-way Chi-Square test with a 95% confidence interval was used to compare antibody prevalence differences in sex and age groups. A *P* value  $\leq 0.05$  was considered significant.

One hundred thirty-one blood samples were collected in two consecutive years, 2000–2001 and 2001–2002, encompassing two harvesting seasons and 109 (83%) of these samples had antibodies to *T. gondii* at a titer  $\geq 25$ . Animals with antibodies to *T. gondii* were 83% (52/63) males, 88% (53/60) females, 88% (81/92) adults, and 77% (23/30) juveniles, 50% (4/8) unknown sex, and 56% (5/9) unknown age. Although the prevalence of antibodies was higher in females than males and higher in adults than juveniles, these differences were not statistically significant. All 14 counties from which bobcats were sampled had at least one animal with antibodies to *T. gondii* (Table 1).

This is the highest prevalence of antibodies to *T. gondii*, with a sample size of this magnitude, reported for *Lynx rufus*. Four previous studies with  $>50$  bobcat blood samples used serum from whole blood and indirect hemagglutination test (IHA), while this study used blood reconstituted from Nobuto blood collecting strips and MAT. They also used different positive dilution thresholds than this study. Based on Dubey and Beattie (1988), a 1:25 dilution was chosen as the

TABLE 1. Prevalence of antibodies to *Toxoplasma gondii* in *Lynx rufus rufus* in Pennsylvania by county.

County	No. of samples	Percent positive ( $\geq 1:25$ )	No. with titers of:		
			25	50	$\geq 500$
Bradford	10	90	0	6	3
Cameron	4	50	0	1	1
Centre	1	100	0	1	0
Clearfield	10	90	0	7	2
Clinton	10	80	0	8	0
Columbia	1	100	0	0	1
Elk	7	86	0	6	0
Forest	1	100	0	1	0
Luzerne	3	67	1	1	0
Lycoming	19	84	2	8	6
Mckean	8	88	0	7	0
Potter	20	85	2	13	2
Sullivan	9	100	2	5	2
Tioga	17	82	0	11	3
Unknown	11	64	1	4	2
Total	131	83	8	79	22

positive dilution threshold for this study, whereas Oertley and Walls (1980) used a positive threshold of 1:16 and Franti et al. (1976), Riemann et al. (1978), and Kikuchi et al. (2004) used 1:64. Because dilutions of 1:25, 1:50, and 1:500 were used in this study, there can be no direct comparison with their data. However, comparing similar dilutions, the number of bobcats with antibodies to *T. gondii* at 1:25 (83%) is four times that reported by Oertley and Walls (1980) at 1:16 (18%). If the number of animals positive for antibodies to *T. gondii* at the 1:50 dilution (77%) is compared with animals positive at 1:64 in the studies of Franti et al. (1976; 69%), Riemann et al. (1978; 61%), and Kikuchi et al. (2004; 50%), the prevalence in this study is still higher. These differences could result from sensitivity of MAT compared with IHA or discrepancies in titer comparisons, or they could simply indicate different levels of *T. gondii* infection from different geographical locations (Dubey and Beattie, 1988). Although other studies have been conducted, a meaningful comparison would be difficult because of their smaller sample size (Walton and Walls, 1964; Franti et al.,

TABLE 2. Number of harvested deer (*Odocoileus virginianus*) and bear (*Ursus americanus*) per sampled county in 2001 and the number of theoretically *T. gondii* infected gut piles.

County	No. harvested deer <sup>a</sup>	No. harvested bear <sup>a</sup>	No. of theoretically infected		Total theoretically infected gut piles
			Deer (60%) <sup>b</sup>	Bear (79.8%) <sup>c</sup>	
Bradford	11,467	47	6,880	38	6,919
Cameron	1,183	126	710	101	811
Centre	6,566	151	3,940	120	4,061
Clearfield	8,781	130	5,269	104	5,373
Clinton	2,494	267	1,496	213	1,711
Columbia	5,343	36	3,206	29	3,235
Elk	2,861	109	1,717	87	1,805
Forest	4,695	81	2,817	65	2,882
Luzerne	5,671	95	3,403	76	3,479
Lycoming	6,040	242	3,624	193	3,818
Mckean	6,859	107	4,115	85	4,202
Potter	6,567	117	3,940	93	3,735
Sullivan	2,215	55	1,329	44	1,373
Tioga	7,328	133	4,397	106	4,504
Total	78,070	1,696	46,843	1,354	48,208

<sup>a</sup> Numbers of harvested deer/bear are from Pennsylvania Game Commission.

<sup>b</sup> Percent seroprevalence to *Toxoplasma gondii* in deer (*Odocoileus virginianus*) is from Humphreys et al. (1995).

<sup>c</sup> Percent seroprevalence to *Toxoplasma gondii* in bear (*Ursus americanus*) is from Briscoe et al. (1993).

1975; Riemann et al., 1975; Marchiondo et al., 1976; Burrige et al., 1979; Dubey et al., 1987; Heidt et al., 1988; Labelle et al., 2001; Dubey et al., 2004; Riley et al., 2004).

Higher prevalence of antibodies to *T. gondii* in adults than juveniles is similar to other studies (Riemann et al., 1978; Labelle et al., 2001), with the exception of the study Oertly and Walls (1980). Differences such as decreases in the rate of vertical transmission and/or environmental prevalence of the parasite may account for the incongruity between this study and that of Oertly and Walls (1980). There was no statistical difference in prevalence of antibodies to *T. gondii* between males and females, which is in accordance with previous studies (Riemann et al., 1978; Oertly and Walls, 1980; Labelle et al., 2001).

The high prevalence of antibodies to *T. gondii* in the Pennsylvania bobcat is probably a reflection of the abundant source of infected intermediate hosts. With a seroprevalence of 60% and 79.8%, respectively, white-tailed deer,

and to a lesser extent bear, may represent a major reservoir of infection (Briscoe et al., 1993; Humphreys et al., 1995; McLean et al., 2005). In part, this may be due to the copious supply of potentially infected carrion (gut piles) left after harvest (Table 2). Raccoons (48.3%) may represent yet another possible source of infection (Dubey et al., 1992). Further research is needed to determine whether other reservoirs, such as lagomorphs and/or rodents, exist in Pennsylvania.

It can be inferred that in Pennsylvania the bobcat plays a major role in the dissemination of *T. gondii* impacting other wildlife and subsequently humans. The ramifications of *T. gondii* infection (or sequelae to infection) on the Pennsylvania bobcat population are not well understood.

#### LITERATURE CITED

- BRISCOE, N., J. G. HUMPHREYS, AND J. P. DUBEY. 1993. Prevalence of *Toxoplasma gondii* infections in Pennsylvania black bear (*Ursus americanus*). *Journal of Wildlife Diseases* 29: 599-601.
- BURRIDGE, M. J., W. J. BIGLER, D. J. FORRESTER, AND

- J. M. HENNEMANN. 1979. Serologic Survey for *Toxoplasma gondii* in wild animals in Florida. *Journal of the American Veterinary Medical Association* 175: 964–967.
- DUBEY, J. P., AND C. P. BEATTIE. 1988. Toxoplasmosis of animals and man. CRC Press, Inc., Boca Raton, Florida, 220 pp.
- , AND G. DESMONTS. 1987. Serological responses of equids fed *Toxoplasma gondii* oocysts. *Equine Veterinary Journal* 19: 337–339.
- , W. J. QUINN, AND W. WEINANDY. 1987. Fatal neonatal toxoplasmosis in a bobcat (*Lynx rufus*). *Journal of Wildlife Diseases* 23: 324–327.
- , A. N. HAMIR, C. A. HANLON, AND C. E. RUPPRECHT. 1992. Prevalence of *Toxoplasma gondii* infection in raccoons. *Journal of the American Veterinary Medical Association* 200: 534–536.
- , D. H. GRAHAM, R. W. DE YOUNG, E. DAHL, M. L. EBERHARD, E. K. NACE, K. WON, H. BISHOP, G. PUNKOSDY, C. SREEKUMAR, M. C. B. VIANNA, S. K. SHEN, O. C. H. KWOK, J. A. SUMNERS, S. DEMARAIS, J. G. HUMPHREYS, AND T. LEHMANN. 2004. Molecular and biologic characteristics of *Toxoplasma gondii* isolates from wildlife in the United States. *Journal of Parasitology* 90: 67–71.
- FRANTI, C. E., G. E. CONNOLLY, H. P. RIEMANN, D. E. BEHYMER, R. RUPPANNER, C. M. WILLADSEN, AND W. LONGHURST. 1975. A survey for *Toxoplasma gondii* antibodies in deer and other wildlife on a sheep range. *Journal of the American Veterinary Association* 167: 565–568.
- , H. P. RIEMANN, D. E. BEHYMER, D. SUTHER, J. A. HOWARTH, AND R. RUPPANNER. 1976. Prevalence of *Toxoplasma gondii* antibodies in wild and domestic animals in northern California. *Journal of the American Veterinary Association* 169: 901–906.
- HEIDT, G. A., R. A. RUCKER, M. L. KENNEDY, AND M. E. BAEYENS. 1988. Hematology, intestinal parasites, and selected disease antibodies from a population of bobcats (*Felis rufus*) in Central Arkansas. *Journal of Wildlife Diseases* 24: 180–183.
- HUMPHREYS, J. G., R. L. STEWART, AND J. P. DUBEY. 1995. Prevalence of *Toxoplasma gondii* antibodies in sera of hunter-killed white-tailed deer in Pennsylvania. *American Journal of Veterinary Research* 56: 172–173.
- KIKUCHI, Y., B. B. CHOMEL, R. W. KASTEN, J. S. MARTENSON, P. K. SWIFT, AND S. J. O'BRIEN. 2004. Seroprevalence of *Toxoplasma gondii* in American free-ranging or captive pumas (*Felis concolor*) and bobcats (*Lynx rufus*). *Veterinary Parasitology* 120: 1–9.
- LABELLE, P., J. P. DUBEY, I. MIKAEELIAN, N. BLANCHETTE, R. LAFOND, S. ST-ONGE, AND D. MARTINEAU. 2001. Seroprevalence of antibodies to *Toxoplasma gondii* in Lynx (*Lynx Canadensis*) and bobcats (*Lynx rufus*) from Quebec, Canada. *Journal of Parasitology* 87: 1194–1196.
- MARCHONDO, A. A., D. W. DUSZYNSKI, AND G. MAVEIN. 1976. Prevalence of antibodies to *Toxoplasma gondii* in wild and domestic animals of New Mexico, Arizona and Colorado. *Journal of Wildlife Diseases* 12: 226–231.
- MCLEAN, M. L., T. S. MCCAY, AND M. J. LOVALLO. 2005. Influence of age, sex, and time of year on diet of the bobcat (*Lynx rufus*) in Pennsylvania. *American Midland Naturalist* 153: 450–453.
- OERTLEY, K. D., AND K. W. WALLS. 1980. Prevalence of antibodies to *Toxoplasma gondii* among bobcats of West Virginia and Georgia. *Journal of the American Veterinary Medical Association* 177: 852–853.
- RIEMANN, H. P., J. A. HOWARTH, R. RUPPANNER, C. E. FRANTI, AND D. E. BEHYMER. 1975. *Toxoplasma* antibodies among bobcats and other carnivores of northern California. *Journal of Wildlife Diseases* 11: 272–276.
- , R. A. THOMPSON, D. E. BEHYMER, R. RUPPANNER, AND C. E. FRANTI. 1978. Toxoplasmosis and Q-fever antibodies among wild carnivores in California. *Journal of Wildlife Management* 42: 198–202.
- RILEY, P. D., J. FOLEY, AND B. CHOMEL. 2004. Exposure to feline and canine pathogens in bobcats and gray foxes in urban and rural zones of a national park in California. *Journal of Wildlife Diseases* 40: 11–22.
- WALTON, B. C., AND K. W. WALLS. 1964. Prevalence of toxoplasmosis in wild animals from Fort Stewart, Georgia as indicated by serological tests and mouse inoculation. *American Journal of Tropical Medicine and Hygiene* 13: 530–533.

Received for publication 10 November 2004.