

**Bartonella spp. in deer keds, *Lipoptena mazamae* (Diptera: Hippoboscidae), from Georgia and South Carolina, USA**

Authors: Reeves, Will K., Nelder, Mark P., Cobb, Kristin D., and Dasch, Gregory A.

Source: Journal of Wildlife Diseases, 42(2) : 391-396

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-42.2.391>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## ***Bartonella* spp. in deer keds, *Lipoptena mazamae* (Diptera: Hippoboscidae), from Georgia and South Carolina, USA**

Will K. Reeves,<sup>1,3</sup> Mark P. Nelder,<sup>2</sup> Kristin D. Cobb,<sup>2</sup> and Gregory A. Dasch<sup>1</sup> <sup>1</sup> Centers for Disease Control and Prevention, 1600 Clifton Rd. NE, Mailstop G-13, Atlanta, Georgia 30333, USA; <sup>2</sup> Clemson University, Department of Entomology, Soils, and Plant Sciences, 114 Long Hall, Clemson, South Carolina 29634, USA; <sup>3</sup> Corresponding author (email: wreeves@alumni.clemson.edu)

**ABSTRACT:** Deer keds, *Lipoptena mazamae* (Diptera: Hippoboscidae), were collected from white-tailed deer (*Odocoileus virginianus*) and humans in Georgia and South Carolina, USA (1 October 2001–6 January 2005) and screened for the presence of DNA from *Bartonella* spp. Forty deer keds were screened for *Bartonella* spp. by polymerase chain reaction using primers specific to the riboflavin synthase gene (*ribC*) of *Bartonella*. *Bartonella* species closely related to *Bartonella schoenbuchensis* and to the etiologic agent of cat-scratch disease (*Bartonella henselae*) were detected in 10 keds and one ked, respectively.

**Key words:** Deer ked dermatitis, ectoparasite, *Odocoileus virginianus*, vector, zoonoses.

*Lipoptena mazamae* Rondani (Diptera: Hippoboscidae) is a blood-feeding ectoparasite of white-tailed deer (*Odocoileus virginianus*) in North America and brocket deer (*Mazama* spp.) in Central and South America (Samuel and Trainer, 1972). In North America, *Lipoptena mazamae* is host specific to the white-tailed deer, but has been reported as an incidental ectoparasite of domestic cattle, pumas (*Puma concolor*), and humans in the United States (Drummond, 1966; Forrester et al., 1996). *Lipoptena mazamae* has been implicated in the transmission of anaplasmosis of cattle (Drummond, 1966) and *Trypanosoma cervi* in cervids (Strickland et al., 1981); however, the precise role of deer keds in the transmission of either agent has not been investigated.

*Bartonella* are gram-negative bacteria that infect the erythrocytes of vertebrates and are putatively transmitted by arthropods or by blood to blood contact (Chomel et al., 1996; Chang et al., 2001; Comer et al., 2001). At least nine species of *Barto-*

*nella* are associated with bartonellosis in humans (Ciervo and Ciceroni, 2004; Dehio et al., 2004). Eight of these are zoonotic pathogens; however, *Bartonella bacilliformis* (agent of Carrion's disease and Oroya fever) has no recognized non-human reservoirs (Karem et al., 2000; O'Rourke et al., 2005). The animal reservoirs, vectors, pathogenicity, and natural history are unknown for most species of *Bartonella*.

The transmission of *Bartonella* spp. to ruminants and *Bartonella schoenbuchensis* to humans and deer by hippoboscidae has been suggested (Dehio et al., 2004; Halos et al., 2004). The role of *Lipoptena cervi* (Linnaeus) in the transmission cycle of *B. schoenbuchensis* is unknown; but this European ked is established in the north-eastern United States (Maa, 1969). White-tailed deer in the southeastern United States are infested with *L. mazamae* (Maa, 1969), which might harbor *B. schoenbuchensis* or similar species of *Bartonella*. Host-seeking deer keds are attracted to and will bite humans (Rantanen et al., 1982). In order to determine if *Bartonella* spp. are present in *L. mazamae*, we collected deer keds from Georgia and South Carolina and tested them for known pathogenic *Bartonella* spp. by polymerase chain reaction (PCR) targeting the riboflavin synthase gene (*ribC*; Johnson et al., 2003).

Adult *L. mazamae* were collected from white-tailed deer carcasses killed by motor vehicles in Georgia and South Carolina (Nelder and Reeves, 2005); two additional keds were collected from humans in Georgia (Table 1). The two keds collected from humans still retained their wings; all

TABLE 1. Collection data and *Bartonella* spp. detected in *Lipoptena mazamae* from Georgia and South Carolina, USA (2003–05).

State/county	Collection site	Host	Date collected	No. tested	<i>Bartonella</i> spp. identified (No. of PCR-positive keds)
Georgia/Rockdale	Panola Mountain State Park	<i>Homo sapiens</i> Linnaeus	25 September 2004	2	None
South Carolina / Anderson	Fants Grove	<i>Odocoileus virginianus</i> (Zimmermann)	1 June 2003	9	None
South Carolina / Clarendon	Highway 251 near Sumter County line	<i>O. virginianus</i>	6 January 2005	3	<i>Bartonella</i> sp. near <i>B. schoenbuchensis</i> (1)
South Carolina / Fairfield	Fairfield	<i>O. virginianus</i>	5 January 2005	4	<i>Bartonella</i> sp. near <i>B. schoenbuchensis</i> (1)
South Carolina / Greenville	Greenville	<i>O. virginianus</i>	1–2 October 2001	1	<i>Bartonella</i> sp. near <i>B. schoenbuchensis</i> (1)
South Carolina / Pickens	Clemson, US 76	<i>O. virginianus</i>	13 June 2004	9	<i>Bartonella</i> sp. near <i>B. schoenbuchensis</i> (3)
SC/Richland	Harbison State Forest, Columbia	<i>O. virginianus</i>	14 November 2004	9	<i>Bartonella</i> sp. near <i>B. schoenbuchensis</i> (3)
South Carolina / Williamsburg	Kingston, Highway 251	<i>O. virginianus</i>	9 September 2004	3	<i>Bartonella henselae</i> (1) <i>Bartonella</i> sp. near <i>B. schoenbuchensis</i> (1) <sup>a</sup>

<sup>a</sup> Positive results for *Bartonella henselae* and *Bartonella* sp. from the South Carolina/Williamsburg samples were not detected from the same keds.

others were wingless. All specimens were fixed and stored in 95% ethanol and identified using taxonomic keys by Maa (1969).

Forty *L. mazamae* were tested by PCR for *Bartonella* spp. Individual whole flies were frozen in liquid nitrogen and crushed with a sterile Teflon pestle. Teflon pestles were cleaned in 10% sodium hypochlorite for 3 hr, rinsed in distilled water, and autoclaved before each use. Total DNA was extracted from the pulverized keds with an IsoQuick Nucleic Acid Extraction Kit (ORCA Research Inc., Bothell, Washington, USA) and resuspended in nuclease-free water. For PCR, the BARTON-1 (5'-TAACCGATATTGGTTGTGTTGAAG-3') and BARTON-2 (5'-TAAAGCTAGAAAGTCTGGCAACATAACG-3') primers were used to amplify a fragment of the riboflavin synthase gene (*ribC*; Johnson et al., 2003). These primers were selected because they were designed to amplify DNA from pathogenic *Bartonella*

spp. Each PCR reaction contained 12.5 µl of Taq PCR Master Mix Kit (Qiagen, Valencia, California, USA), 7.5 µl of nuclease-free water, 1.25 µl of each primer, and 2.5 µl of DNA template in water. The PCR products were separated by 2% agarose gel electrophoresis and visualized under ultraviolet light with ethidium bromide. Distilled water was used as a negative control.

The PCR products were purified with a QIAquick PCR Purification Kit (Qiagen). Duplicate sequencing reactions were performed with a BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, California, USA) using PCR primers, and excess dye was removed with a DyeEx 2.0 column (Qiagen). Sequences were determined using an ABI 3100 capillary sequencer (Applied Biosystems). Primer sequences were removed and sequences were assembled with Seqmerge (Accelrys, San Diego, California, USA). Assembled sequences

were compared to those in GenBank using the BLAST 2.0 program (NCBI, Bethesda, Maryland, USA). Identification of *Bartonella* spp. was based on a sequence similarity to known taxa. We considered 100% similarity in sequence data to represent the same taxa.

Voucher specimens of *L. mazamae* were deposited in the Clemson University Arthropod Collection and the novel sequence for the *ribC* gene from the *Bartonella* sp. closely related to *B. schoenbuchensis* was deposited in GenBank under accession number DQ125469.

DNA from a novel *Bartonella* sp. closely related to *B. schoenbuchensis* was detected in 10 of 40 deer keds from South Carolina (Table 1). The 533-base pair (bp) sequence from the *ribC* gene of this new *Bartonella* sp. was 96% similar to *Bartonella schoenbuchensis* (GenBank accession number AY116628) from Europe. We detected the same bacterial agent in keds from all geographical regions sampled including the Piedmont and Upper and Lower Coastal Plains. Our data imply that a novel *Bartonella* sp. closely related to *B. schoenbuchensis* is widely distributed in South Carolina and might be transmitted by *L. mazamae*. Hippoboscid ectoparasites of deer have been implicated as vectors of *B. schoenbuchensis* in Europe (Dehio et al., 2004), but hippoboscid ectoparasites of white-tailed deer in North America have not been tested. *Bartonella schoenbuchensis* infects deer and may be the causative agent of deer ked dermatitis in humans (Dehio et al., 2004). In severe cases, deer ked dermatitis causes itchy lesions that can persist for up to a year (Rantanen et al., 1982; Dehio et al., 2004).

Genetic heterogeneity within strains of *Bartonella* spp. closely related to *B. schoenbuchensis* in North America are partially known (e.g., Chang et al., 2000) and cervids in the Nearctic region might harbor a variant strain of *B. schoenbuchensis* or a closely related species. The full extent of genetic variability within *B. schoenbuchensis* or related species cannot

be fully determined until these agents are cultured and their taxonomy better defined. Heterogeneity among strains of *B. schoenbuchensis* exists in Europe (Dehio et al., 2001). Culturing of the *Bartonella* spp. from both deer and keds would allow for analyses and comparison between strains or species of *Bartonella*.

The sequence for the 535-bp *ribC* amplicon was 100% identical to the *ribC* sequence for *Bartonella henselae* Houston strain-1 (GenBank# AJ132928). *Bartonella henselae* infects over 20,000 humans in the United States annually (Kaplan et al., 2002). Humans are infected by *B. henselae* when scratched or bitten by a bacteremic domestic cat, but this bacterium is possibly transmitted enzootically by the cat flea, *Ctenocephalides felis* (Bouche) (Chomel et al., 1996). DNA from *B. henselae* has been detected in argasid and ixodid ticks and stable flies (reported as *Stomoxys* spp. but presumably *Stomoxys calcitrans* Linnaeus), but the role of ticks and stable flies in the transmission cycle of this bacterium is unexplored (Sanogo et al., 2003; Chung et al., 2004; Loftis et al., 2005). Deer keds were previously not known to harbor *B. henselae* in North America. Presumably, the deer ked either acquired the bacterium by feeding on an infected deer or via a transovarial route. Cat fleas will rarely infest cervids (Szabo et al., 2000) and transmission of *B. henselae* by infected cat fleas to deer would seem unlikely.

The presence of DNA from *Bartonella* in *L. mazamae* does not demonstrate infection of the ectoparasite or vector competence; the DNA from *Bartonella* could have originated in an undigested blood meal or have been transmitted by transovarial routes. We did not detect *Bartonella* spp. in all flies, including the winged forms, which indicates that *L. mazamae* acquire these agents and that these bacteria are not ubiquitous obligate symbiotes.

Wild and domestic ruminants harbor *Bartonella* spp. (Chang et al., 2000;

Breitschwerdt et al., 2001) and infection rates can be high; more than 90% of mule deer (*Odocoileus hemionus*) in the western United States are infected with unnamed *Bartonella* spp. (Chang et al., 2000). The vectors of these *Bartonella* are unknown but could include ticks, lice, or hippoboscids. The relatively high prevalence of *Bartonella* infections in wild cervids suggests that the vectors of these agents may be widespread, common, or highly efficient vectors. The possibility of pathogen transfer between wildlife and humans or domestic animals exists, and might have been the case when a cattle rancher was infected with *Bartonella vinsonii* (Welch et al., 1999).

Hippoboscids harbor symbiotic species of *Bartonella* (e.g., Candidatus "*Bartonella melophagi*" formerly "*Wolbachia melophagi*") and other bacteria that are not pathogenic or transmitted to vertebrates (Bequaert, 1953; Reeves, 2005; Small, 2005). The PCR primers used in our study were designed to amplify DNA from species of *Bartonella* that infect vertebrates (Johnson et al., 2003) and did not amplify DNA from symbiotic *Bartonella* spp. of hippoboscids. If the symbiotic *Bartonella* spp. of hippoboscids were amplified by our PCR primers, all extracts would produce amplicons. A larger sample size is needed to determine if winged hippoboscids harbor potentially pathogenic *Bartonella* spp.

There are substantial populations of white-tailed deer in the southeastern United States that might serve as reservoirs for a *Bartonella* sp. closely related to *B. schoenbuchensis*. The vector potential of *L. mazamae* and other hematophagous arthropods feeding on white-tailed deer are of potential medical or veterinary concern. Samuel and Trainer (1972) noted that infestation of white-tailed deer by *L. mazamae* increases with deer age and that infestations are more prevalent during warmer months of the year. Future studies should address the transmission of this *Bartonella* sp. closely related to *B. schoen-*

*buchensis* by keds; potential relationships between *Bartonella*-bacteremia and the age of deer; the potential for associated pathology; and further detection and identification of *Bartonella* spp. in white-tailed deer ectoparasites. The assumption that deer are infected with these agents but remain healthy can only be accepted if clinical trials are conducted. For example, *B. henselae* was presumed to be non-pathogenic to cats but recent discoveries by Kordick et al. (1999) have implicated chronic infections by this bacterium as the cause of several previously idiopathic diseases in cats.

We thank Russell Hubright and Laurie Reid (South Carolina Forestry Commission) for collecting keds and the American Society for Microbiology for partial funding of this project. We also thank P. H. Adler (Clemson University) for providing laboratory equipment and work space, J. A. Korecki for loaning material from the Clemson University Arthropod Collection, and A. D. Loftis for help with the initial PCR setup. The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the funding agency.

#### LITERATURE CITED

- BEQUAERT, J. C. 1953. The Hippoboscidae or louse flies (Diptera) of mammals and birds. Part I. Structure, physiology and natural history. *Entomologica Americana* 32: 1–209.
- BREITSCHWERDT, E. B., S. SONTAKKE, A. CANNEDY, S. I. HANCOCK, AND J. M. BRADLEY. 2001. Infection with *Bartonella weissii* and detection of *Nanobacterium* antigens in a North Carolina beef herd. *Journal of Clinical Microbiology* 39: 879–882.
- CHANG, C. C., B. B. CHOMEL, R. W. KASTEN, R. HELLER, K. M. KOCAN, H. UENO, K. YAMAMOTO, V. C. BLEICH, B. M. PIERCE, B. J. GONZALES, P. K. SWIFT, W. M. BOYCE, S. S. JANG, H. BOULOUIS, AND Y. PIEMONT. 2000. *Bartonella* spp. isolated from wild and domestic ruminants in North America. *Emerging Infectious Diseases* 6: 306–311.
- , V. ROMANO, AND N. TIETZE. 2001. Molecular evidence of *Bartonella* spp. in questing adult *Ixodes pacificus* ticks in California. *Journal of Clinical Microbiology* 39: 1221–1226.

- CHOMEL, B. B., R. W. KASTEN, K. FLOYD-HAWKINS, B. CHI, K. YAMAMOTO, J. ROBERTS-WILSON, A. N. GURFIELD, R. C. ABBOTT, N. C. PEDERSEN, AND J. E. KOEHLER. 1996. Experimental transmission of *Bartonella henselae* by the cat flea. *Journal of Clinical Microbiology* 34: 1952–1965.
- CHUNG, C. Y., R. W. KASTEN, S. M. PAFF, B. A. VAN HORN, M. VAYSSIER-TAUSSAT, H.-J. BOULOUIS, AND B. B. CHOMEL. 2004. *Bartonella* spp. DNA associated with biting flies from California. *Emerging Infectious Diseases* 10: 1311–1313.
- CIERVO, A., AND L. CICERONI. 2004. Rapid detection and differentiation of *Bartonella* spp. by a single-run real time PCR. *Molecular and Cellular Probes* 18: 307–312.
- COMER, J. A., C. D. PADDOCK, AND J. E. CHILDS. 2001. Urban zoonoses caused by *Bartonella*, *Coxiella*, *Ehrlichia*, and *Rickettsia* species. *Vector-Borne and Zoonotic Diseases* 1: 91–118.
- DEHIO, C., C. LANZ, R. POHL, P. BEHRENS, D. BERMOND, Y. PIEMONT, K. PELZ, AND A. SANDER. 2001. *Bartonella shoenbuchii* sp. nov., isolated from the blood of wild roe deer. *International Journal of Systematics and Evolutionary Microbiology* 51: 1557–1565.
- , U. SAUDER, AND R. HIESTAND. 2004. Isolation of *Bartonella schoenbuchensis* from *Lipoptena cervi*, a blood-sucking arthropod causing deer ked dermatitis. *Journal of Clinical Microbiology* 42: 5320–5323.
- DRUMMOND, R. O. 1966. *Lipoptena mazamae* Rondani (Diptera: Hippoboscidae), a louse fly of deer, on cattle in southwestern Texas. *Journal of Parasitology* 52: 825.
- FORRESTER, D. J., G. S. MCLAUGHLIN, S. R. TELFORD, JR., G. W. FOSTER, AND J. W. MCGOWN. 1996. Ectoparasites (Acari, Mallophaga, Anoplura, Diptera) of white-tailed deer, *Odocoileus virginianus*, from southern Florida. *Journal of Medical Entomology* 33: 96–101.
- HALOS, L., T. JAMAL, R. MAILLARD, B. GIRARD, J. GUILLOT, B. CHOMEL, M. VAYSSIER-TAUSSAT, AND H. J. BOULOUIS. 2004. Role of Hippoboscidae flies as potential vectors of *Bartonella* spp. infecting wild and domestic ruminants. *Applied and Environmental Microbiology* 70: 6302–6305.
- JOHNSON, G., M. AYERS, S. C. MCCLURE, S. E. RICHARDSON, AND R. TELLIER. 2003. Detection and identification of *Bartonella* species pathogenic for humans by PCR amplification targeting the riboflavin synthase gene (*ribC*). *Journal of Clinical Microbiology* 41: 1069–1072.
- KAPLAN, S., J. RAWLINGS, C. PADDOCK, J. CHILDS, R. REGNERY, AND M. REYNOLDS. 2002. Cat-scratch disease in children—Texas, September 2000–August 2001. *Morbidity and Mortality Weekly Report* 51: 2647–2649.
- KAREM, K. L., C. D. PADDOCK, AND R. L. REGNERY. 2000. *Bartonella henselae*, *B. quintana*, and *B. bacilliformis*: Historical pathogens of emerging significance. *Microbes and Infection* 2: 1193–1205.
- KORDICK, D. L., T. T. BROWN, K. SHIN, AND E. B. BREITSCHWERDT. 1999. Clinical and pathologic evaluation of chronic *Bartonella henselae* or *Bartonella clarridgeiae* infections in cats. *Journal of Clinical Microbiology* 37: 1536–1547.
- LOFTIS, A. D., J. S. GILL, M. E. SCHRIFTER, M. L. LEVIN, M. E. EREMEEVA, M. J. R. GILCHRIST, AND G. A. DASCH. 2005. Detection of *Rickettsia*, *Borrelia*, and *Bartonella* in *Carios kelleyi* (Acari: Argasidae). *Journal of Medical Entomology* 42: 473–480.
- MAA, T. C. 1969. Studies in Hippoboscidae (Diptera) Part 2. *Pacific Insects Monograph* 20: 1–312.
- NELDER, M. N., AND W. K. REEVES. 2005. Ectoparasites of road-killed vertebrates in northwestern South Carolina, USA. *Veterinary Parasitology* 129: 313–322.
- O'ROURKE, L. G., C. PITULLE, B. C. HEGARTY, S. KRAYCIRIK, K. A. KILLARY, P. GROSENSTEIN, J. W. BROWN, AND E. B. BREITSCHWERDT. 2005. *Bartonella quintana* in cynomolgus monkey (*Macaca fascicularis*). *Emerging Infectious Diseases* 11: 1931–1934.
- RANTANEN, T., T. REUNALA, P. VUOJOLAHTI, AND W. HACKMAN. 1982. Persistent pruritic papules from deer ked bites. *Acta Dermatovenere (Stockholm)* 62: 307–311.
- REEVES, W. K. 2005. Molecular genetic evidence for a novel bacterial endosymbiont of *Icosta americana* (Diptera: Hippoboscidae). *Entomological News* 116: 263–265.
- SAMUEL, W. M., AND D. O. TRAINER. 1972. *Lipoptena mazamae* Rondani, 1878 (Diptera: Hippoboscidae) on white-tailed deer in southern Texas. *Journal of Medical Entomology* 9: 104–106.
- SANOGO, Y. O., Z. ZEAITER, G. CARUSO, F. MEROLA, S. SHPYNOV, P. BROUQUI, AND D. RAOULT. 2003. *Bartonella henselae* in *Ixodes ricinus* ticks (Acari: Ixodida) removed from humans, Belluno province, Italy. *Emerging Infectious Diseases* 9: 329–332.
- SMALL, R. W. 2005. A review of *Melophagus ovinus* (L.), the sheep ked. *Veterinary Parasitology* 130: 141–155.
- STRICKLAND, R., R. R. GERRISH, AND J. R. SMITH. 1981. Arthropods. In *Diseases and parasites of white-tailed deer*, W. R. Davidson, F. A. Hayes, V. F. Nettles and F. F. Kellogg (eds.). Tall Timbers Research Station, Tallahassee, Florida. pp. 363–389.
- SZABO, M. P. J., E. R. MATUSHIMA, M. DE CAMPOS PEREIRA, K. WERTHER, AND J. M. B. DUARTE. 2000. Cat flea (*Ctenocephalides felis*) infestation in quarantined marsh deer (*Blastocerus dichotomus*) populations. *Journal of Zoo and Wildlife Medicine* 31: 576–577.
- WELCH, D. F., K. C. CARROLL, E. K. HOFMEISTER, D.

H. PERSING, D. A. ROBINSON, A. G. STEIGERWALT,  
AND D. J. BRENNER. 1999. Isolation of a new  
subspecies, *Bartonella vinsonii* subsp. *arupensis*,  
from a cattle rancher: Identity with isolates  
found in conjunction with *Borrelia burgdorferi*

and *Babesia microti* among naturally infected  
mice. *Journal of Clinical Microbiology* 37: 2598–  
2601.

*Received for publication 8 August 2005.*