

Hematologic and Biochemical Reference Intervals of Free-Living Mediterranean Pond Turtles (*Mauremys leprosa*)

Judit Hidalgo-Vila,^{1,3} Carmen Díaz-Paniagua,¹ Natividad Pérez-Santigosa,¹ Agustín Plaza,² Inés Camacho,² and Fernando Recio² ¹Estación Biológica de Doñana, CSIC. Avda. María Luisa s/n, 41013, Sevilla, Spain; ²Clinical Chemistry Department, Hospital Universitario de Valme. Bellavista, 41014 Sevilla, Spain; ³Corresponding author (email: judit@ebd.csic.es)

ABSTRACT: Reference intervals of hematologic and biochemical blood profiles were obtained from 56 male and 58 female Mediterranean pond turtles (*Mauremys leprosa*) captured from the wild in different periods of their annual cycle. Mean (or median in nonnormal distributions) values of leukocyte differential were 53.8% and 58.5% heterophils, 35.3% and 32.6% eosinophils, 6.3% and 5.8% lymphocytes, 4.3% and 2% monocytes, and 0% and 0% basophils in males and females, respectively. Biochemical values did not differ from other chelonians, but values were generally higher in females than in males.

Key words: Blood chemistry, blood profile, freshwater turtle, hematology, *Mauremys leprosa*, Mediterranean pond turtle.

Hematologic and blood biochemical parameters are normally used to evaluate the health of animals and humans. In reptiles, several studies have described the general characteristics of the blood profile (Dessauer, 1970; Duguy, 1970; Frye, 1991; Campbell, 1996; Stein, 1996), but there are many species for which reference values are still unknown or imprecise. In live individuals, blood profiles provide a minimally invasive tool that can support health evaluations, especially in relation to determining potential effects associated with such factors as pollution, disease, invasion by exotic species, among others. Such evaluations are dependent on reliable reference values in healthy animals. Blood parameters of reptiles may be influenced by many factors, such as age, sex, seasonality, and reproduction (Dessauer, 1970; Duguy, 1970; Frye, 1991; Wilkinson, 2003), and these parameters can vary through the annual cycle or even throughout the life of the individuals.

In this study, the blood profile of the Mediterranean pond turtle (*Mauremys*

leprosa), has been documented by analyzing data from free-living males and females throughout the annual cycle. A prior study on seasonal changes in hematology and blood chemistry of this species (Pagés et al., 1992), and a study that analyzed seasonal changes in lymphoid distribution (Muñoz and De la Fuente, 2004) have been reported, but were limited to freshwater turtles kept in captivity. This manuscript reveals the wide variation in hematologic and biochemical parameters obtained from the turtles under natural conditions and establishes the reference values necessary for the evaluation of individuals from wild populations.

Mauremys leprosa inhabits the north of Africa, the Iberian Peninsula, and some areas of southern France (Keller and Busack, 2001). Although it is globally considered to be species of a least concern conservation status (Cox et al., 2006), *M. leprosa* is ranked as a vulnerable species in Spain. Population declines, and disappearances of several local populations, have occurred because of habitat fragmentation, pollution, and exotic species introduction (Pleguezuelos et al., 2002).

From March 2003 through November 2004, 114 healthy, adult freshwater turtles (56 males and 58 females) were captured at the Doñana Biological Reserve (Huelva, southwestern Spain). For each turtle, blood (0.5–0.7 ml) was collected from the occipital venous sinus (Martínez-Silvestre and Marco, 2002) using a disposable sterile syringe with a 0.6-gauge (23G) needle. Blood smears for differential leukocyte counts (heterophils, lymphocytes, basophils, eosinophils, and monocytes) were stained with a commercial stain (Diff

Quick); 0.1 ml was placed in a tube containing lithium heparin for hematologic analyses. Red blood cell (RBC) counts and total white blood cells (WBC) counts were done using the methodology of Campbell (1996). Hematocrit or packed cell volume (PCV) was measured after centrifuging whole blood in hematocrit tubes containing ammonium-heparin at $13,700 \times G$ for 120 sec (Stat Spin VT-RH 12). Mean cell volume (MCV) was estimated using RBC and PCV values. The remaining blood (0.4–0.6 ml) was placed in a tube for separation of serum and, after 60 min, was centrifuged at $12,000 \times G$ for 150 sec (Stat Spin VT-RT 12). Serum was kept at -80 C for 14 days before analysis. Biochemical analysis including calcium, phosphorus, glucose, cholesterol, total protein, uric acid, sodium, creatinine, and potassium concentrations and aspartate aminotransferase (AST), creatine phosphokinase (CK), lactate dehydrogenase (LDH), and alkaline phosphatase (ALP) activities were performed on a Roche/Hitachi Modular Analytics automated chemistry analyzer (Roche Diagnostics, Mannheim, Germany). All individuals were released 2–3 hr after capture.

For males and females, means, standard deviations, and 95% confidence intervals were calculated for the hematologic and blood biochemistry profiles. For nonnormally distributed variables, reference intervals were defined by median and minimum and maximum ranges.

Hematologic and biochemical blood profiles for adult males and females of *M. leprosa* are presented in Table 1. The leukocyte differential determined in this study does not follow the pattern previously described for this species. In captive *M. leprosa*, the percentage of lymphocytes ranged between 64.9 and 57.8 (Muñoz and De la Fuente, 2004). Although the prevalence of lymphocytes is variable, the low frequency found for free-living *M. leprosa* (4.2–7.7%) does not coincide with values reported for other

chelonians, in which these cells represent over 50% of the leukocyte differential count (Duguy, 1967, 1970; Taylor and Jacobson, 1982; Wood and Ebanks, 1984; Martínez-Silvestre et al., 2001). Heterophils were the most common leukocytes and were two times more prevalent than eosinophils. Similar high frequencies have been described for *Trachemys scripta* (ISIS, 2002), *Testudo radiata* (Marks and Citino, 1990), and *Testudo graeca* and *Testudo hermanni* (Lawrence, 1986). The prevalences of eosinophils, averaging 35 and 32% in male and female *M. leprosa*, respectively, were higher than reported in other species of chelonians (Lawrence, 1986; Marks and Citino, 1990; Anderson et al., 1997; ISIS, 2002). Monocyte and lymphocyte percentages were less than 7%, and basophils were generally absent from the differential count. A similar leukocyte differential was obtained for another fresh water turtle (*Emys orbicularis*) sampled in the same locality (Hidalgo-Vila, 2006) and suggests the influence of environmental factors in the study area.

The PCV, RBC and WBC values were higher in males than in females, as has been reported for RBC data in other reptiles (Duguy, 1967; Duguy, 1970; Frye, 1991). In contrast, MCV values were higher in females (Table 1). Wide reference intervals were obtained for MCV; similar results were attributed to inherent errors in manual RBC counting for desert tortoises, *Gopherus agassizii* (Christopher et al., 1999). However, in *M. leprosa*, the variation could be also explained by seasonal differences in the volume of the erythrocytes (Hidalgo-Vila, unpubl. data).

The biochemical reference intervals of *M. leprosa* are presented in Table 1, and all were within the common ranges for chelonians (Dessauer, 1970; Rosskopf, 1982; Taylor and Jacobson, 1982; Marks and Citino, 1990; Bolten and Bjorndal, 1992; Kölle, 1999; Martínez-Silvestre et al., 2001). With the exception of LDH and creatinine, all the biochemical values analyzed were higher in females than in

TABLE 1. Hematologic and biochemical blood profiles for adult from 56 male and 58 female Mediterranean pond turtles (*Mauremys leprosa*) captured from the wild in different periods of their annual cycle.^a

Parameter	Male				Female			
	n	Mean median ^b	SD	95% C.I. minimum–maximum ^c	n	Mean median ^b	SD	95% C.I. minimum–maximum ^c
PCV (%)	53	20.5	5.04	19.1–21.9	58	18.5	4.9	17.2–19.8
MCV (fL)	31	513.6 ^b		292.9–2,411.8 ^c	31	550.4 ^b		259.2–2149.0 ^c
RBC ($\times 10^6/\mu\text{l}$)	29	0.42	0.21	0.34–0.50	31	0.33	0.14	0.27–0.38
WBC ($\times 10^3/\mu\text{l}$)	29	4.58	2.34	3.69–5.47	31	4.40	2.52	3.47–5.33
Heterophils (%)	26	53.8	17.1	46.9–60.8	29	58.5	17.3	52.0–65.1
Eosinophils (%)	26	35.3	16.6	28.5–42.0	29	32.6	17.5	25.9–39.2
Lymphocytes (%)	26	6.3	3.5	4.9–7.7	29	5.8	4.3	4.1–7.4
Monocytes (%)	26	4.3	3.4	2.3–6.5	29	2 ^b		0–11 ^c
Basophils (%)	26	0 ^b		0–2 ^c	29	0 ^b		0–1 ^c
Calcium (mmol/L)	55	1.7	0.4	2.3–2.6	56	4.3	1.4	3.4–4.7
Phosphorus (mmol/L)	56	0.7	0.3	0.6–0.8	56	1.1	0.5	0.9–1.2
Glucose (mmol/L)	55	3.7	2.1	3.1–4.2	54	4.5	2.1	3.9–5.1
Cholesterol (mmol/L)	55	2.3	1.5	1.9–2.7	55	3.6	1.6	3.2–4.0
Total protein (g/dl)	54	3.1	1.3	2.8–3.5	55	3.4	1.3	3.1–3.8
Uric acid ($\mu\text{mol/L}$)	55	91.0	64.0	73.7–108.3	56	93.2	56.3	78.1–108.3
Sodium (mmol/L)	52	132.4	2.9	131.6–133.2	55	132.5	3.5	131.6–133.5
Potassium (mmol/L)	53	3.4	0.5	3.2–3.5	55	3.4	0.4	3.3–3.6
AST (IU/L)	54	149 ^b		27–531 ^c	54	161	84	138–184
CK (IU/L)	56	1,429	882	1,193–1,665	54	1,738	1,254	1,396–2,080
LDH (IU/L)	54	949	402	839–1,059	56	934	445	815–1,023
ALP (IU/L)	55	37	20	31–42	56	47	30	39–55
Creatinine ($\mu\text{mol/L}$)	55	22.0	7.0	20.2–24.0	56	22.0	7.4	20.0–24.0

^a 95% C.I. = 95% confidence interval; PCV = packed cell volume; MCV = mean cell volume; RBC = red blood cell; WBC = white blood cells; AST = aspartate aminotransferase; CK = creatine phosphokinase; LDH = lactate dehydrogenase; ALP = alkaline phosphatase.

^b For nonnormally distributed variables, reference intervals were defined by the median rather than the mean.

^c For nonnormally distributed variables, reference intervals were defined by minimum and maximum ranges rather than the 95% C.I.

males. High levels of calcium, phosphorus, and cholesterol are common in female reptiles during egg development and vitellogenesis (Dessauer, 1970; Campbell, 2004), and in the present study, samples from females were collected during the breeding season. These sex-related differences were not found in the previous study of captive *M. leprosa* (Pagés et al., 1992) and the smaller size of the turtles included in that study suggests that those animals were subadults. Glucose, uric acid, sodium, calcium, and phosphorus values obtained in wild *M. leprosa* are dissimilar to those reported for captive *M. leprosa* (Pagés et al., 1992). For calcium, this different can be explained by the high

number of gravid females. Differences in the parameters could be associated with the influence of captivity on physiology or even to the younger age of turtles in the previous study.

We are grateful for the encouragement provided by R. Soriguer, and to X. Ruiz, M. Florencio, and A. Portheault for their field and laboratory assistance. J. Hidalgo-Vila was financed by research grant I3P-CSIC. This study was supported by Consejería Medio Ambiente-Junta de Andalucía (UE cofinanced project).

LITERATURE CITED

ANDERSON, N. L., R. F. WACK, AND R. HATCHER. 1997. Hematology and clinical chemistry reference

- ranges for clinically normal, captive New Guinea snapping turtle (*Elseya novaguineae*) and the effects of temperature, sex, and sample type. *Journal of Zoo and Wildlife Medicine* 28: 394–403.
- BOLTEN, A. B., AND K. A. BJORNDALE. 1992. Blood profiles for a wild population of green turtles (*Chelonia mydas*) in the southern Bahamas: Size-specific and sex-specific relationships. *Journal of Wildlife Diseases* 28: 407–413.
- CAMPBELL, T. W. 1996. Clinical pathology. In *Reptile medicine surgery*, D. R. Mader (ed.). W. B. Saunders Company Ltd, Philadelphia, Pennsylvania, pp. 248–257.
- . 2004. Clinical chemistry of reptiles. In *Veterinary hematology clinical chemistry*, D. B. Troy (ed.). Lippincott Williams and Wilkins, Baltimore, Maryland, pp. 493–498.
- CHRISTOPHER, M. M., K. H. BERRY, I. R. WALLIS, K. A. AGY, B. T. ENEN, AND C. C. PETERSON. 1999. Reference intervals and physiologic alterations in hematologic and biochemical values of free-ranging desert tortoises in the Mojave Desert. *Journal of Wildlife Diseases* 35: 212–238.
- COX, N., J. CHANSON, AND S. TUART. 2006. The status and distribution of reptiles and amphibians of the Mediterranean basin. IUCN, Gland, Switzerland, and Cambridge, UK, pp. 42.
- DESSAUER, H. C. 1970. Blood chemistry of reptiles: Physiological and evolutionary aspects. In *Biology of the reptilia*, Vol. 3, C. Gans and T. Parson (eds.). Academy Press, London, UK, pp. 1–72.
- DUGUY, R. 1967. Le cycle annuel des éléments figurés dus sang chez *Emys orbicularis* L., *Lacerta muralis* Laur., et *Natrix maura* L. *Bulletin de la Société Zoologique de France* 92: 23–37.
- . 1970. Numbers of blood cells and their variation. In *Biology of the reptilia*, Vol. 3, C. Gans and T. Parson (eds.). Academy Press, London, UK, pp. 93–110.
- FRYE, F. L. 1991. Hematology as applied to clinical reptile medicine. In *Biomedical surgical aspects of captive reptile husbandry*, Vol. 1. F. L. Frye (ed.). Krieger Publishing Co., Malabar, Florida, pp. 209–280.
- HIDALGO-VILA, J. 2006. Hematología y perfil bioquímico sanguíneo en las especies de galápagos *Mauremys leprosa* y *Emys orbicularis*. Aspectos sanitarios y evaluación de la introducción de la especie exótica *Trachemys scripta elegans* sobre la fauna autóctona. Ph. D. Thesis. Universidad de Sevilla, España, 308 pp.
- [ISIS] INTERNATIONAL SPECIES INFORMATION SYSTEM. 2002. *Trachemys scripta* common slider. Physiological reference values—American units. CD-ROM. Apple Valley, Minnesota.
- KELLER, C., AND S. D. BUSACK. 2001. *Mauremys leprosa* (Schweigger, 1812) Maurische Bachschildkröte. In *Handbuch der reptilien und amphibien Europas. Schildkröten (Testudines)*, Vol. I. U. Fritz (ed.). Aula, Wiebelsheim, pp. 57–88.
- KÖLLE, P., H. LAMNEK, AND R. HOFFMANN. 1999. Blutwerte bei der Europäischen Sumpfschildkröte (*Emys orbicularis*). *Tierärztl Prax*, 27: 198–201.
- LAWRENCE, K. 1986. Seasonal variations in haematological data from Mediterranean tortoises (*Testudo graeca* and *Testudo hermanni*) in captivity. *Research in Veterinary Science* 40: 225–230.
- MARKS, S., AND B. CITINO. 1990. Hematology and serum chemistry of the radiated tortoise (*Testudo radiata*). *Journal of Zoo and Wildlife Medicine* 21: 342–344.
- MARTÍNEZ-SILVESTRE, A., AND I. MARCO. 2002. Venipuncture technique of the occipital venous sinus in freshwater aquatic turtles. *Journal of Herpetological Medicine and Surgery* 12: 31–32.
- , S. LAVIN, I. MARCO, J. MONTANO, J. RAMON LÓPEZ, AND J. SOLER MASSANA. 2001. Haematology and plasma chemistry of captive *Testudo marginata*. *Chelonii* 3: 187–189.
- MUÑOZ, F. J., AND M. DE LA FUENTE. 2004. Seasonal changes in lymphoid distribution of the turtle *Mauremys caspica*. *Copeia* 1: 178–183.
- PAGÉS, T., V. I. PEINADO, AND G. VISCOR. 1992. Seasonal changes in hematology and blood chemistry of the freshwater turtle *Mauremys caspica leprosa*. *Comparative Biochemistry and Physiology* 103A: 275–278.
- PLEGUEZUELOS, J. M. 2002. In *Atlas y libro rojo de los anfibios y reptiles de España*. (2nd Edition), J. M. Pleguezuelos, R. Márquez and M. Lizana (eds.). Dirección General de la Conservación de la Naturaleza-Asociación Herpetológica, España, Madrid, pp. 501–532.
- ROSSKOPF, W. J. 1982. Normal hemogram and blood chemistry values for California desert tortoises. *Veterinary Medicine/Small Animal Clinician* 77: 85–87.
- STEIN, G. 1996. Hematologic and blood chemistry values in reptiles. In *Reptile medicine and surgery*, D. R. Mader (ed.). W. B. Saunders Company Ltd., Philadelphia, Pennsylvania, pp. 473–483.
- TAYLOR, R. W., AND E. R. JACOBSON. 1982. Hematology and serum chemistry of the gopher tortoise, *Gopherus polyphemus*. *Comparative Biochemistry and Physiology* 72A: 425–428.
- WILKINSON, R. 2003. Clinical Pathology. In *Medicine and surgery of tortoises and turtles*, S. McArthur, R. Wilkinson and J. Meyer (eds.). Blackwell Publishing, Oxford, UK, pp. 141–186.
- WOOD, F. E., AND G. K. EBANKS. 1984. Blood cytology and hematology of the green sea turtle, *Chelonia mydas*. *Herpetologica* 40: 331–336.

Received for publication 9 January 2007.