

Epidemiologic Survey for *Brucella suis* Biovar 2 in a Wild Boar (*Sus scrofa*) Population in Northwest Italy

Authors: Bergagna, Stefania, Zoppi, Simona, Ferroglio, Ezio, Gobetto, Mario, Dondo, Alessandro, et al.

Source: Journal of Wildlife Diseases, 45(4) : 1178-1181

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-45.4.1178>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Epidemiologic Survey for *Brucella suis* Biovar 2 in a Wild Boar (*Sus scrofa*) Population in Northwest Italy

Stefania Bergagna,¹ Simona Zoppi,¹ Ezio Ferroglio,^{2,5} Mario Gobetto,³ Alessandro Dondo,² Elisabetta Di Giannatale,⁴ Maria Silvia Gennero,¹ and Carla Grattarola¹ ¹Istituto Zooprofilattico Sperimentale di Piemonte, Liguria e Valle d'Aosta, Torino, Italy; ²Dipartimento di Produzioni Animali, Epidemiologia ed Ecologia, Università degli Studi di Torino, Grugliasco, Italy; ³Servizio Veterinario Area "C" TO4, Torino, Italy; ⁴National Reference Centre for Brucellosis, Teramo, Italy, Studi di Torino, Grugliasco, Italy; ⁵Corresponding author (email: ezio.ferroglio@unito.it)

ABSTRACT: In 2001, antibodies to *Brucella* spp. were detected in the wild boar (*Sus scrofa*) population of Regional Park in the Piedmont, northwest Italy. This was the first report of swine brucellosis in Italian wildlife, and in response, we conducted a survey on wild boars culled in the park to evaluate the presence and epidemiologic pattern of *Brucella*. In total, 2,267 serum samples and 1,841 tissue samples were collected and tested from 2001 to 2007. Differences in antibody and infection prevalence among sampling years were statistically significant ($\chi^2=28.2$, $P<0.0001$ and $\chi^2=27.8$, $P<0.0001$, respectively). Serologic and culture results also differed between gender and age groups, and for serology, a positive age trend was observed in both male and female age classes ($\chi^2_{(\text{trend})}=90.9$, $P<0.00005$ OR: 1, 2.5, 4.9, 7.5; $\chi^2_{(\text{trend})}=43.1$, $P<0.00002$, OR: 1, 2.5, 2.9, 4.8, respectively).

Key words: *Brucella suis* 2, Italy, swine brucellosis, wild boar.

Apart from classical swine fever, swine brucellosis, caused by *Brucella suis* biovar 2, is one of the most important endemic diseases in the wild boar (*Sus scrofa*) populations in Central Europe (Garin-Bastuji et al., 2000; Godfroid, 2002; Godfroid and Käsbohrer, 2002; Cvetnic et al., 2003; Al Dahouk et al., 2005; Vengust et al., 2006; Leuenberger et al., 2007). In Italy, *B. suis* biovar 2 was detected in hares (*Lepus europaeus*) that were imported from Hungary in 1995, but in spite of an increased sampling effort, brucellosis was not detected in endemic wildlife until the end of the 1990s (Quaranta et al., 1995; Ebani et al., 2003). In northwest Italy, brucellosis has been reported in the Piedmont in alpine ibex (*Capra ibex*;

Ferroglio et al., 1998) and in chamois (*Rupicapra rupicapra*; Ferroglio et al. 2003). In 2001, antibodies to *Brucella* spp. were detected in wild boars in the "Mandria di Venaria Reale" Regional Park (Piedmont, northwest Italy 45°8'7"N, 7°37'31"E; Gennero et al., 2006). This park was established in 1978, includes 3,600 ha of fenced area, and is characterized by plain forest with open meadows and streams; it is inhabited by red deer (*Cervus elaphus*, 200–500 animals) and an estimated 500 wild boar. In this study, we conducted a survey of wild boars culled in the park in order to evaluate the presence and epidemiologic pattern of *Brucella* in the wild boar population.

Blood samples were collected from 2,267 culled wild boars from 2001 to 2007. Blood was allowed to clot, and serum was separated by centrifugation and stored at –20 C until testing. After 2002, tissue samples (spleen and genital organs) also were collected from 1,841 animals. Gender and age were determined for all animals. Age was estimated based on tooth eruption pattern; four age categories were used (A: <10 mo; B: 11–18 mo; C: 19–24 mo; D: >25 mo; Briedermann, 1986).

Serum samples were tested by rose Bengal test (RBT) and complement fixation test (CFT) according to Alton et al. (1988), and 20 g of tissue (uterus, spleen, and testicles) were macerated and inoculated onto *Brucella* medium according to Farrell (1974). Samples were inoculated

onto modified Thayer Martin Medium (Martin et al., 1996), and media plates were enriched with 10% horse serum and selective supplements; cultures were incubated for at least 10 days. If *Brucella* was suspected during Stamp's staining, colonies were identified to genus by classical techniques (Alton et al., 1988) and typed by means of biochemical (Alton et al., 1988) and molecular tests (Corbel et al., 1983; Bricker and Halling, 1994; Cloeckert et al., 1995) by the laboratories of the National Reference Centre of Brucellosis (Teramo, Italy). Epi Info™ 6 for DOS Current Version 6.04d (Centers for Disease Control and Prevention, Atlanta, Georgia, USA) was used to test for associations between serologic and culture results with year of sampling, season, gender, and age by chi-square tests. Differences were considered significant when $P < 0.05$. Agreement between serology and bacteriology test results were calculated using WinEpiScope 2.0 (N. de Blas, C. Ortega, K. Frankena, J. Noordhuizen, M. Thrusfield: <http://www.clive.ed.ac.uk/winepiscope/>).

Antibodies to *Brucella* spp. were detected with RBT and CFT in 448 of 2,267 serum samples. Of 1,841 animals tested by culture, 448 tested positive (Table 1). All isolates were classified as *Brucella suis* biovar 2.

Differences observed in antibody prevalence among sampling years were statistically significant ($\chi^2 = 28.2$, $P < 0.0001$); prevalence of infection as determined by culture differed by years ($\chi^2 = 27.8$, $P < 0.0001$). For both serology and culture, prevalence decreased significantly in 2005 and 2006 ($\chi^2 = 12.1$, $P < 0.0005$ and $\chi^2 = 12.6$, $P < 0.0005$, respectively) and increased in 2007 (compared to 2004 and 2005; $\chi^2 = 9.9$, $P < 0.002$ and $\chi^2 = 13.3$, $P < 0.0003$, respectively). Differences in the prevalences of antibody- and culture-positive animals also were detected between some gender and age groups. An increased antibody prevalence with age was observed in both males and females

($\chi^{2(\text{trend})} = 90.9$, $P < 0.00005$ OR: 1, 2.5, 4.9, 7.5 and $\chi^{2(\text{trend})} = 43.1$, $P < 0.00002$, OR: 1, 2.5, 2.9, 4.8, respectively). These age differences were not observed with culture results. Differences among antibody prevalences in males and females were only detected within the C age class ($\chi^2 = 8.9$, $P < 0.003$).

Antibody prevalence was significantly lower ($\chi^2 = 10.3$, $P = 0.0003$) from August (8%) to November and did not significantly vary during the other months of the year. Prevalence estimates based on culture were significantly lower ($\chi^2 = 38.9$, $P = 0.00000$) from July to November, and the lowest prevalence occurred in August (2%).

In comparisons of serology and culture results for the 1,762 wild boars for which both serum and tissue samples were available, the test agreement was low (Kappa = 0.37), and 26.8% (51/190) of samples positive on bacteriology were negative on both serologic tests.

Our data confirm the presence of *B. suis* 2 in wild boar in Italy, and it appears that the prevalence based on serology ($19.8 \pm 1.6\%$) and culture ($10.8 \pm 1.4\%$) is high. These prevalence estimates are consistent with estimates from traditionally (Godfroid et al., 1994; Al Dahouk et al., 2005) and nontraditionally endemic areas (Hars et al., 2000; Ruiz-Fons et al., 2006). Annual prevalence estimates (antibodies and culture) suggest that *B. suis* 2 was maintained in this population from 2001 to 2004 at a relatively constant level. In 2005 and 2006, prevalence significantly decreased, and in 2007, it returned to previous 2001–2004 levels. The cause of this variation is not known, and data related to wild boar density are not available to evaluate potential population-dependent causes. Considering the low agreement among serologic and culture results, it is recommended that both be used for future epidemiologic surveys.

The increased antibody prevalence with age observed with both males and females may be attributable to sexual maturity. In

TABLE 1. Serologic and bacteriologic results (positives/examined; prevalence with 95% confidence interval [CI95%] in parentheses) on sera and organs from wild boar (*Sus scrofa*) collected from 2001 to 2007 in the “Mandria di Venaria Reale” Regional Park, Piedmont, Italy (RBT = rose Bengal test; CFT = complement fixation test; M = male; F = female; n.d. = not done).

Years ^a	Sex	RBT/CFT	Bacteriologic test
2001	M	11/180 [6.11% CI95%: 2.61–9.61]	n.d.
	F	8/201 [3.98% CI95%: 1.28–6.68]	n.d.
2002	M	44/151 [29.14% CI95%: 21.89–36.39]	17/141 [12.06% CI95%: 6.68–17.43]
	F	25/134 [18.66% CI95%: 12.06–25.25]	9/124 [7.26% CI95%: 2.69–11.83]
2003	M	42/130 [32.31% CI95%: 24.27–40.35]	20/144 [13.89% CI95%: 8.24–19.54]
	F	45/187 [24.06% CI95%: 17.94–30.19]	25/205 [12.20% CI95%: 7.72–16.68]
2004	M	39/144 [27.08% CI95%: 19.83–34.34]	22/148 [14.87% CI95%: 9.13–20.60]
	F	43/136 [31.62% CI95%: 23.80–39.43]	22/148 [14.87% CI95%: 9.13–20.60]
2005*	M	43/214 [20.09% CI95%: 14.73–25.46]	19/220 [8.64% CI95%: 4.92–12.35]
	F	42/250 [16.08% CI95%: 12.17–21.43]	14/254 [5.51% CI95%: 2.71–8.32]
2006*	M	25/167 [14.97% CI95%: 9.56–20.38]	8/135 [5.93% CI95%: 1.94–9.91]
	F	32/193 [16.58% CI95%: 11.33–21.83]	12/155 [7.74% CI95%: 3.54–11.95]
2007	M	31/104 [29.81% CI95%: 21.02–38.60]	19/98 [19.39% CI95%: 11.56–27.22]
	F	18/76 [23.68% CI95%: 14.13–33.24]	11/69 [15.94% CI95%: 7.30–24.58]
Subtotal	M	235/1090 [21.56% CI95%: 19.12–24.00]	105/886 [11.85% CI95%: 9.72–13.98]
	F	213/1177 [18.10 CI95%: 15.90–20.30]	93/955 [9.74% CI95%: 7.86–11.62]
Total		448/2267 [19.76% CI95%: 18.12–21.40]	198/1841 [10.75% CI95%: 9.34–12.17]

^a Asterisk denotes significant values ($P < 0.05$).

females, antibody prevalence increased sharply from class A (<10 mo) to class B (>11 mo; $\chi^2 = 9.6$, $P < 0.002$), while it did not increase significantly from class B to class C. Difference in antibody prevalence between genders was only detected in age class C ($\chi^2 = 8.9$, $P < 0.003$), while differences in other age classes were not significant. No significant differences were observed between males and females in bacteriology, even if a positive correlation seemed to emerge for class C (19–24 mo; $\chi^2 = 3.3$, $P = 0.06$). The higher prevalence of antibody- and culture-positive animals observed from December to June could be related to reproductive season; estrus generally starts in late autumn until February, while partum occurs mainly from February until June (Durio et al., 1995).

Brucella suis has not been previously detected in Italy, from either wild boar or domestic swine, and it is possible that it was introduced by hares imported from European countries where it is endemic (Dedek, 1983; Hubalek et al., 1993; Quaranta et al., 1995; Pilaszek et al.,

1996). As suggested by Lanfranchi et al. (2003), wildlife disease management must be based on avoiding introduction of new pathogens, and a health risk assessment should be included in all future translocation plans.

The authors gratefully acknowledge the game warden of the “Mandria di Venaria Reale” Regional Park for invaluable help in sample collection. We are also grateful to all laboratory technicians for their excellent technical assistance. The research was funded by grant from the Ministry of Health, Rome, and from the Regional Public Health Department.

LITERATURE CITED

- AL DAHOUK, S., K. NOCKLER, H. TOMASO, W. D. SPLETTSTOESSER, G. JUNGENSEN, U. RIBER, T. PETRY, D. HOFFMANN, H. C. SCHOLZ, A. HENSEL, AND H. NEUBAUER. 2005. Seroprevalence of brucellosis, tularemia, and yersiniosis in wild boars (*Sus scrofa*) from north-eastern Germany. *Journal of Veterinary Medicine Series B* 52: 444–455.
- ALTON, G. G., M. L. JONES, R. D. ANGUS, AND J. M. VERGER. 1988. In: *Bacteriologic Methods and Serologic Methods. Techniques for the Brucellosis Laboratory*. Institut National de la Recherche Agronomique, Paris, France, pp. 13–61, 82–113.

- BRICKER, B., AND S. M. HALLING. 1994. Differentiation of *Brucella abortus* bv. 1, 2 and 4, *Brucella melitensis*, *Brucella ovis* and *Brucella suis* bv. 1 by PCR. *Journal of Clinical Microbiology* 32: 2660–2666.
- BRIEDERMANN, L. 1986. Schwarzwild. Neumann & Neudamm Verlag, Berlin, 529 pp.
- CLOECKAERT, A., J. M. VERGER, M. GRAYON, AND O. GREPINET. 1995. Restriction site polymorphism of the genes encoding the major 25 kDa and 36 kDa outer membrane proteins of *Brucella*. *Microbiology* 141: 2111–2121.
- CORBELL, M. J., K. P. W. GILL, E. L. THOMAS, AND D. MC. L. F. D. HENDRY. 1983. Methods for the identification of *Brucella*. MAFF Publications, Booklet 2085. Alnwick, Northumberland, U.K., 36 pp.
- CVETNIC, Z., M. MITAK, M. OCEPEK, M. LOJKIC, S. TERZIC, L. JEMERSIC, A. HUMSKI, B. HABRUN, B. SOSTARIC, M. BRSTILO, B. KRT, AND B. GARIN-BASTUJI. 2003. Wild boars (*Sus scrofa*) as reservoirs of *Brucella suis* biovar 2 in Croatia. *Acta Veterinaria Hungarica* 51: 465–473.
- DEDEK, J. 1983. Zur Epizootiologie der Schweinebrucellose unter besonderer Berücksichtigung von Erregerreservoirern. *Monatshefte für Veterinärmedizin* 38: 852–856.
- DURIO, P., U. GALLO-ORSI, E. MACCHI, AND A. PERRONE. 1995. Monthly birth distribution and structure of an Alpine population of wild boar (*Sus scrofa*) in north-west Italy. In *Ongules/Ungulates* 91, F. Spitz, G. Janeau, G. González and S. Aulagnier (eds.), S.F.E.P.M.-I.R.G.M., Paris-Toulouse, France, pp. 395–397.
- EBANI, V. V., D. CERRI, A. POLI, AND E. ANDREANI. 2003. Prevalence of *Leptospira* and *Brucella* antibodies in wild boars (*Sus scrofa*) in Tuscany, Italy. *Journal of Wildlife Diseases* 39: 718–722.
- FARRELL, I. D. 1974. The development of new selective medium for the isolation of *Brucella abortus* from contaminated source. *Research in Veterinary Science* 16: 280–286.
- FERROGLIO, E., F. TOLARI, E. BOLLO, AND B. BASSANO. 1998. First isolation of *Brucella melitensis* from alpine ibex. *Journal of Wildlife Diseases* 34: 400–402.
- , S. GENNERO, L. ROSSI, AND F. TOLARI. 2003. Epidemiologia della brucellosi nel Camoscio alpino. *Journal of Mountain Ecology* 7: 229–232.
- GARIN-BASTUJI, B., J. HARS, D. CALVEZ, M. THIEBAUD, AND M. ARTOIS. 2000. Brucellose du porc domestique et du sanglier sauvage due à *Brucella suis* biovar 2 en France. *Epidémiologie et Santé Animale* 38: 1–5.
- GENNERO, M. S., C. GRATTAROLA, S. BERGAGNA, S. ZOPPI, A. BARBARO, AND A. DONDO. 2006. Trend of *Brucella suis* infection in wild boar in Piedmont region (2002–2005). *Epidémiologie et Santé Animale* 49: 59–62.
- GODFROID, J. 2002. Brucellosis in wildlife. *Revue Scientifique et Technique* 21: 277–286.
- , AND A. KÄSBOHRER. 2002. Brucellosis in the European Union and Norway at the turn of the twenty-first century. *Veterinary Microbiology* 90: 135–145.
- , P. MICHEL, L. UYTTERHAEGEN, C. DE SMEDT, F. RASSENEUR, F. BOELAERT, C. SAEGERMAN, AND X. PATIGNY. 1994. Brucellose enzootique à *Brucella suis* biotype 2 chez le sanglier (*Sus scrofa*) en Belgique. *Annales de Médecine Vétérinaire* 138: 263–268.
- HARS, J., M. VALERY, F. CHADUC, B. GARIN-BASTUJI, O. PINGUET, AND S. ROSSI. 2000. Surveillance de la Brucellose du sanglier et du lièvre dans le département de l'Allier. *Bulletin d'Information sur la Pathologie des Animaux Sauvages en France* 23: 121–138.
- HUBALEK, Z., Z. JURICOVA, I. SVOBODOVA, AND J. HALOUZKA. 1993. A serologic survey for some bacterial and viral zoonoses in game animals in the Czech Republic. *Journal of Wildlife Diseases* 29: 604–607.
- LANFRANCHI, P., E. FERROGLIO, G. POGLAYEN, AND V. GUBERTI. 2003. Wildlife veterinarian, conservation and public health. *Veterinary Research Communication* 27: 567–574.
- LEUENBERGER, R., P. BOUJON, B. THÜR, R. MISEREZ, B. GARIN-BASTUJI, J. RÜFENACHT, AND K. D. STÄRK. 2007. Prevalence of classical swine fever, Aujeszky's disease and brucellosis in a population of wild boar in Switzerland. *Veterinary Record* 160: 362–368.
- MARTIN, C. M., M. P. JIMENEZ DE BAGUES, M. BARBERAN, AND J. M. BLASCO. 1996. Comparison of two selective media for the isolation of *Brucella melitensis* from naturally infected sheep and goats. *Veterinary Record* 27: 409–411.
- PILASZEK, J., K. SZULOWSKI, AND M. TRUSZCZYNSKI. 1996. Diagnosis of brucellosis in hares. *Zycie Weterynaryjne* 71: 180–182.
- QUARANTA, V., R. FARINA, A. POLI, D. CERRI, AND L. PALAZZO. 1995. Sulla presenza di *Brucella suis* biovar 2 nella lepre in Italia. *Selezione Veterinaria* 36: 953–958.
- RUZ-FONS, F., J. VICENTE, D. VIDAL, U. HÖFLE, D. VILLANÚA, C. GAUSS, J. SEGALÉS, S. ALMERÍA, V. MONTORO, AND C. GORTÁZAR. 2006. Seroprevalence of six reproductive pathogens in European wild boar (*Sus scrofa*) from Spain: The effect on wild boar female reproductive performance. *Theriogenology* 65: 731–743.
- VENGUST, G., Z. VALENCAK, AND A. BIDOVEC. 2006. A serological survey of selected pathogens in wild boar in Slovenia. *Journal of Veterinary Medicine Series B* 53: 24–27.

Received for publication 16 September 2008.