

PHYSIOLOGIC VALUES OF STONE SHEEP

Author: FRANZMANN, ALBERT W.

Source: Journal of Wildlife Diseases, 7(3): 139-141

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-7.3.139

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

| Journal of Wildlife Diseases Vol. 7, July, 1971 | 139 |
|---|-----|
| Huizinga, H. W., and G. E. Cosgrove, PULMONARY ARTERIAL FILARI- ASIS IN THE HOUSE SPARROW | 205 |
| Plumb, J. A., TISSUE DISTRIBUTION OF CHANNEL CATFISH VIRUS | 213 |
| Kocan, R. M., and A. Sprunt IV, THE WHITE-CROWNED PIGEON: A FRUIT-EATING PIGEON AS A HOST FOR TRICHOMONAS | |
| GALLINAE | 217 |
| ANNOUNCEMENTS | |
| Training in Wildlife Diseases in North America | 219 |
| Letter to the Editor | 221 |
| Special Publications of the Index-Catalogue of Medical and Veterinary | |
| Zoology | 221 |
| Book Review | 222 |
| 1971 Conservation Directory | 223 |
| Journal to Adopt Metric (SI) Units in 1972 | 223 |
| Attention Members — 1972 Dues | 223 |
| PROGRAM FOR THE ANNUAL WDA CONFERENCE | 224 |

PHYSIOLOGIC VALUES OF STONE SHEEP

ALBERT W. FRANZMANN, Idaho Cooperative Wildlife Research Unit, University of Idaho, Moscow, Idaho 83843

Abstract: Physiologic values obtained from 10 stone sheep (*Ovis dalli stonei*) were reported, providing a start in establishing base line values for the species. The influence of excitability, diet, and stress of the sheep at sampling upon certain values was recognized.

INTRODUCTION

Physiologic values are necessary criteria for disease study of a species. Most domestic animals and some wild species have established values. The stone sheep (Ovis Dalli Stonei) has no published values to my knowledge. This report presents physiologic values obtained from 10 captive stone sheep at the Okanagan Game Farm, Penticton, British Columbia on October 27, 1970.

MATERIALS AND METHODS

The sheep were in a 3 acre enclosure and were fed good quality alfalfa hay and protein supplement. Salt and water were available at all times. Each individual's condition in the herd at time of sampling was rated 9 on a 10 point grading system modified for stone sheep. There was no clinical evidence of disease and the herd's reproductive rate has been above 80% the past 2 years. The sheep were highly excited at time of handling.

A field laboratory was established utilizing the Unitest System (Biodynamics Inc., Indianapolis, Ind.) to obtain these values: blood glucose, blood urea nitrogen, cholesterol, serum glutamic oxalacetic transaminase, hemoglobin, and total protein.

Blood mineral values were determined by using the facilities of the Washington State University, School of Veterinary Medicine Clinical Pathology Laboratory,



Pullman, Washington. The calcium determinations were done by microtitration (Oxford Laboratories, San Mateo, California), and the phosphorus determination by Phospho-Pak (Uni Tech Chemical Mfg. Co., Sun Valley, California). Erythrccyte and leukocyte counts were obtained using the Coulter model B electronic particle counter courtesy of The Endoparasite Vector Pioneering Research Laboratory, Agriculture Research Service, Pullman, Washington.

RESULTS

Physiologic values obtained from the stone sheep are shown in Tables 1 and 2 with the individuals listed in order of handling and bleeding. There are some wide variations in values between individuals and comparing these with published values from Rocky Mountain bighorn sheep^{1,4} and domestic sheep² there are notable differences as well as similarities between groups. With small sample sizes and many potential variables

Table 1. Blood serum chemistry values for stone sheep.

| Sex/Age | BUN* Cholesterol (mg/100 ml) | | Total Glucase Protein (g/100 ml) | | Ca P (mg/100 ml) | | SGOT** |
|---------|---------------------------------|-----|--|------|---------------------|-----|--------|
| M 6 mo. | 37 | 95 | 180 | 10.6 | 8.3 | 4.7 | 175 |
| F 6 mo. | 80 | 140 | 245 | 8.9 | 8.7 | 5.9 | 130 |
| F 6 mo. | 30 | 110 | 180 | 8.7 | 10.7 | 3.3 | 220 |
| F 6 mo. | 90 | 100 | 270 | 10.1 | 8.3 | 4.4 | 175 |
| F1yr. | 32 | 90 | 305 | 7.0 | 8.0 | 3.7 | 180 |
| F 4 yr. | 40 | 95 | 250 | 6.0 | 7.6 | 1.6 | 150 |
| M 6 mo. | 37 | 100 | 450 | 3.1 | 10.7 | 4.6 | 30 |
| F 5 yr. | 44 | 90 | 200 | 7.1 | 10.7 | 4.2 | 110 |
| F 6 yr. | 38 | 110 | 240 | 3.6 | 9.7 | 3.3 | 160 |
| F 5 yr. | 40 | 115 | 310 | 3.5 | 8.3 | 2.5 | 110 |

* Blood urea nitrogen

** Serum glutamic oxalacetic transaminase (Karmen units)

| Table 2. | Hematologic and physiologic values for stone sheep. |
|----------|---|
| | |

| Age/Sex | * WBC (10 ³) | ** RBC (10 ^a) | *** PCV (%) | Hemoglobin g/100 ml | Heart Rate/min | Respiratory Rate/min | Rectal Temp. (F) |
|---------|-----------------------------|------------------------------|----------------|------------------------|-------------------|-------------------------|---------------------|
| M 6 mo. | 6.4 | 5.09 | 54 | 22.0 | 120 | 80 | 105.6 |
| F 6 mo. | 8.6 | 8.91 | 53 | 20.5 | 120 | 50 | 104.2 |
| F 6 mo. | 6.7 | 7.08 | 55 | 20.5 | 88 | 36 | 103.8 |
| F 6 mo. | | _ | 54 | 18.0 | 120 | 40 | 104.6 |
| F1yr. | 6.1 | 4.54 | 52 | 20.0 | 140 | 72 | 104.2 |
| F 4 yr. | 5.4 | 7.51 | 51 | 19.0 | 110 | 40 | 103.8 |
| M 6 mo. | 11.7 | 8.71 | 56 | 21.0 | 220 | 48 | 104.6 |
| F 5 yr. | 3.6 | 7.59 | 50 | 15.5 | 112 | 36 | 103.8 |
| F 6 yr. | 5.1 | 8.27 | 48 | 18.0 | 148 | 60 | 102.8 |
| F 5 yr. | 6.8 | 6.30 | 52 | 20.0 | 148 | 52 | 103.2 |

* leukocytes

** erythrocytes

*** packed cell volume

affecting each group, a meaningful interpretation of differences between groups would not be valid, but a discussion of the possible influence of some known variables in this sample may aid in comparing results.

DISCUSSION

There are several factors such as excitability, high protein intake, and stress inherent with sampling this group of sheep which influenced the results obtained.

Excitability can influence physiologic values and in particular such values as glucose, rectal temperature, respiratory and heart rates.² The stone sheep blood glucose values are very high compared to domestic sheep¹ as are the rectal

temperature, respiratory and heart rates.⁸ Excitability could also have had an influence on high blood urea nitrogen (BUN) levels, but perhaps more important was the fact that these sheep were on a high protein diet which directly affects BUN.²

Serum glutamic oxalacetic transaminase (SGOT) values reflect cell necrosis⁵ and this value increases with muscular activity and stress. This corresponds with these results.

Some of the particularly variant individual values such as for total protein, and the generally low phosphorus values cannot be explained from the limited values available, but this does manifest the need for accumulated physiologic data on this and other wild species to establish base line values for useful interpretation of results.

Acknowleigement

I am most grateful to Mr. Ed Lacey and the organization at the Okanagan Game Farm for their cooperation and assistance, and to the College of Forestry, Wildlife, and Range Science and the Idaho Cooperative Wildlife Research Unit for their support.

LITERATURE CITED

- CAMPBELL, L. A., and D. S. KRONFIELD. 1961. Estimation of low concentrations of plasma glucose using glucose oxidase. Am. J. Vet. Res. 22: 587.
- 2. COLES, E. H. 1967. Veterinary Clinical Pathology. 455 pp. W. B. Saunders, Philadelphia.
- 3. DUKES, H. H. 1955. The Physiology of Domestic Animals. 817 pp. Cornell University Press, Ithaca, N.Y.
- 4. FRANZMANN, A. W., and E. T. THORNE. 1970. Physiologic values in wild bighorn sheep (*Ovis canadensis canadensis*) at capture, after handling, and after captivity. J.A.V.M.A. 157: 647-650.
- WHANGER, P.O., P. H. WESWIG, O. H. MUTH, and J. E. OLDFIELD. 1969. Tissue lactic dehydragenase, glutamic-oxalacetic transaminase, and peroxidase changes of selenium-defecient myopathic lambs. J. Nutr. 99: 331-337.
- 6. WOOLF, A., and D. C. KRADEL. 1970. Hematologic values of captive Rocky Mountain bighorns. J. Wildl. Dis. 6: 67-68.

Received for publication December 21, 1970