

EVALUATION OF METAL PARTITIONING ACROSS HUMBOLDT PENGUIN (SPHENISCUS HUMBOLDTI) EGG COMPONENTS

Authors: Daniels-Abdulahad, Mya, Giarikos, Dimitrios, Adkesson, Michael J., and Hirons, Amy C.

Source: Journal of Wildlife Diseases, 60(2) : 474-489

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/JWD-D-22-00176>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Evaluation of Metal Partitioning across Humboldt Penguin (*Spheniscus humboldti*) Egg Components

Mya Daniels-Abdulahad,^{1,5} Dimitrios Giarikos,^{2,3} Michael J. Adkesson,⁴ and Amy C. Hiron^{1,3}

¹ Department of Marine and Environmental Sciences, Halmos College of Arts and Sciences, Nova Southeastern University, 8000 N. Ocean Dr., Dania Beach, Florida 33004, USA

² Department of Chemistry and Physics, Halmos College of Arts and Sciences, Nova Southeastern University, 3301 College Ave., Fort Lauderdale, Florida 33314, USA

³ SECLER: Study of Environmental Conservation through Leading-Edge Research, Nova Southeastern University, 3301 College Ave., Fort Lauderdale, Florida 33314, USA

⁴ Chicago Zoological Society, Brookfield Zoo, 3300 Golf Rd., Brookfield, Illinois 60513, USA

⁵ Corresponding author (email: md1961@nova.edu)

ABSTRACT: Humboldt Penguin (*Spheniscus humboldti*) population declines are attributable to several multifaceted anthropogenic impacts. At present, the exposure of Humboldt Penguins to high concentrations of heavy metals in the marine environment is a preeminent concern, due to mining along the Peruvian coast near key rookery sites. Metal and selenium concentrations were determined in eggs collected from September 2020 to April 2021 from a managed-care penguin population at the Brookfield Zoo to establish reference values for health indices conducted on wild populations. Concentrations of 16 elements, with emphasis on those found in mine efflux—arsenic, cadmium, copper, lead, mercury, selenium, and zinc—were assessed via inductively coupled plasma mass spectrometry in yolk, albumen, and eggshell. Data analyses indicate a clear delineation between egg constituents, with lipid-rich yolk displaying notably higher concentrations ($\mu\text{g/g}$) of arsenic (0.20 ± 0.064), chromium (0.086 ± 0.03), cobalt (0.01 ± 0.003), iron (238.65 ± 54.72), lead (0.32 ± 0.97), manganese (2.71 ± 0.66), molybdenum (0.57 ± 0.14), tin (3.29 ± 0.99), and zinc (64.03 ± 13.01) than other components (albumen and eggshell). These data confirm that heavy metals are partitioned differently across Humboldt Penguin egg components, which provides insight into the potential connection between embryonic nutrient source contamination and subsequent chick viability.

Key words: Albumen, eggshell, heavy metals, Humboldt Penguin, yolk.

INTRODUCTION

Seabirds play an integral role in the establishment and preservation of stable marine ecosystems through their extensive trophic diversity and global ubiquity (Colabuono et al. 2016). However, their populations are often destabilized by anthropogenic activities. The Humboldt Penguin (*Spheniscus humboldti*) has been recognized for nearly 2 decades as a species vulnerable to extinction, largely because of human activity (BirdLife International 2020). Population declines throughout the species' range of Peru and Chile are attributed to guano harvesting (Duffy 1983; Paredes and Zavalaga 2001), resource competition from commercial fisheries (Thiel et al. 2007), trophic restructuring following spatiotemporally cyclical El Niño Southern Oscillation events (Taylor et al. 2008), and industrial mining (Zavalaga and Paredes 1997). Although coastal

mining operations have long threatened the species with habitat degradation and direct anthropogenic disturbance, including mass culls organized to remove the “nuisance” birds (Zavalaga and Paredes 1997; Paredes et al. 2002, 2003), recent mining expansions near the species' largest Peruvian rookery, Punta San Juan Reserve (PSJ, $15^{\circ}22'S$, $75^{\circ}11'W$), have made heavy metal exposure a significant concern (Adkesson et al. 2019).

Peru is the world's second largest producer of copper (Cu), zinc (Zn), and silver (Ag); it is also South America's leading contributor of gold (Au) and lead (Pb), third largest source of tin (Sn), and fourth largest source of molybdenum (Mo; Ministerio de Energía y Minas 2020). The country's largest open-pit iron (Fe) ore mine, producing 4 million tons annually, is within 20 km of PSJ, and an expanding copper mine is within 6 km (Adkesson et al. 2018).

Contaminants from local mines may reach the PSJ rookery directly via runoff from waste retention ponds where flocculated debris is consolidated before removal and disposal. Additionally, the northward flow of the Humboldt Current in the Pacific Ocean off the coast of South America may facilitate transport of contaminant runoff from Tacna, a mining community in southern Peru that manages the world's fifth largest copper mine, as well as copper tailings from dumping sites along Chile's northern coast (Castilla and Correa 1997).

Many metals, such as Cu, Fe, selenium (Se), and Zn, are essential micronutrients required to facilitate many physiologic processes; however, excess amounts, particularly of acutely toxic heavy metals, can lead to deleterious effects (Richards 1997). Heavy metals are a group of high-atomic-mass metals and metalloids that are toxic at low concentrations (Fergusson 1990), and previous research on Humboldt Penguins at PSJ confirmed heavy metal exposure in this protected population (Adkesson et al. 2019). Although heavy metal toxicity is dependent on dose, chemical form (e.g., organic vs. inorganic metal compounds), route of exposure, and physiologic parameters (e.g., sex, age, body condition, immune status), chronic exposure can cause severe physiologic detriment, including endocrine dysfunction, neurologic disruption, and immunohematologic complications (Scheuhammer 1987). Lipophilic metals, such as Cu, manganese (Mn), and Zn (Tjälve and Gottofrey 1991), exhibit distribution partiality once ingested and absorbed by the body, resulting in a disproportionate accumulation of contaminants in adipose-rich tissues, such as the brain. Metals also tend to accumulate in organs responsible for toxin elimination, such as the liver and kidneys (Das et al. 2003). Avian-specific effects of heavy metal exposure and bioaccumulation include decreased egg production, behavioral abnormalities (Fry 1995), and teratogenesis (Scheuhammer 1987).

Mechanisms of heavy metal elimination in birds include excretion (e.g., via feces), feather deposition, and egg offloading (Burger 1994; Ackerman et al. 2016, 2017, 2019). Although

maternal offloading is well established as a physiologic pathway for toxin elimination in fowl (Holcman and Stibilj 1997; Grace and MacFarlane 2016; Kabeer et al. 2021), passerines (Mora 2003), and shorebirds (Burger 2002; Burger and Gochfeld 2004), heavy metal research on penguin egg components is scarce and, when present, almost entirely comprises data from Antarctic species (Metcheva et al. 2006, 2011; Smichowski et al. 2006; Jerez et al. 2011, 2013a, b; Celis et al. 2014). We aimed to quantify concentrations of heavy metals in the egg components (shell, yolk and white) of Humboldt Penguins, a South American species, using eggs from a managed-care population at the Brookfield Zoo, Chicago, Illinois, USA.

MATERIALS AND METHODS

Sample collection

Whole eggs were collected from nine breeding pairs of Humboldt Penguins under managed care at the Brookfield Zoo, Chicago, Illinois, USA, from September 2020 to April 2021. Eggs were refrigerated in plastic specimen containers (4 C) until shipment to Nova Southeastern University (Fort Lauderdale, Florida, USA) for analysis. Analyzed samples included eggshell ($n=18$), yolk ($n=14$), and albumen ($n=14$). Internal components for four eggs were omitted from analyses because of structural degradation.

Sample preparation and analysis

Whole eggs were rinsed with ultrapure deionized water (18.2 megohms) from a Barnstead water (Lake Balboa, California, USA) purification system, and gently scrubbed with a nylon cleaning brush. Eggs were then cracked along the equatorial axis and separated into eggshell, albumen, and yolk. Individual egg components were placed in acid-washed glass beakers and dried for a minimum of 24 h in an isotherm vacuum oven (Thermo Fisher Scientific, Waltham, Massachusetts, USA) at 100 C, with pressure reduced below 10^{-2} torr using a vacuum pump (14008-01 model Welch 1400 DuoSeal, Welch, Mt. Prospect, Illinois, USA). Approximately 0.2 g (dry mass) of each sample was placed in a microwave acid digestion vessel (Parr Instrument Company, Moline, Illinois, USA) with 4 mL of 30% hydrogen peroxide (Sigma-Aldrich, St. Louis, Missouri, USA) and 1 mL of concentrated trace-metal-grade nitric

acid (Sigma-Aldrich). The vessel was heated in a conventional microwave oven at 400 W for 35 s to achieve internal device pressure up to 1200 psi and reactant temperatures up to 250 °C, ensuring total sample digestion, then cooled and depressurized for 45 min.

Digested samples were shipped to the University of Southern Mississippi Center for Trace Analysis (Hattiesburg, Mississippi, USA) for analysis of 16 elements (15 metals plus selenium) via inductively coupled plasma-mass spectrometry (ICP-MS): aluminum (Al), arsenic (As), cadmium (Cd), chromium (Cr), cobalt (Co), copper (Cu), iron (Fe), lead (Pb), manganese (Mn), mercury (Hg), molybdenum (Mo), nickel (Ni), selenium (Se), tin (Sn), vanadium (V), and zinc (Zn). Before analysis, digested samples were diluted fivefold in 0.64 M ultrapure nitric acid containing 2 ppb indium as an internal standard and held in acid-washed Teflon autosampler vials (Thermo Fisher). Mass spectrometer scans were performed at low (Cd-111, Hg-199, 200, 201, 202, Pb-208), medium (Al-27, V-51, Cr-52, Mn-55, Fe-56, Co-59, Ni-60, Cu-63, Zn-66), and high (As-75, Se-77, 82) resolution, depending on the isotope. To correct for molybdenum oxide interference on Cd, Mo-98 was monitored. External standards were used with a high standard and a blank rerun every eight samples. Two US Geological Survey reference water concentrations were also assessed as part of each analytical run to verify standardization. In several cases, sample calibration was also verified by standard additions. Blanks of ultrapure deionized water and trace metal basis nitric acid (3%, 4%, 5%) were used for quality control purposes. The ICP-MS detection limits of each element are provided in Supplementary Materials Table S1. No certified reference materials were used because none are available for heavy metal concentrations in egg components.

Statistical analyses were performed using Excel (version 16.60; Microsoft Corporation, Redmond, Washington, USA) and R statistical software (R 4.2.0 GUI 1.78 High Sierra build [8075]; R Core Team 2022). Geometric mean was used rather than arithmetic mean, to mitigate the impact of data outliers. A Shapiro-Wilk test was used to determine if the data were normally distributed, and a Bartlett's test was used to determine homogeneity of variances. Nonparametric analyses were used because of variable data distributions and lack of homogeneity. Kruskal-Wallis and post-hoc tests were performed to determine if metal concentrations

varied among egg components. Potential correlations in heavy metal concentrations within egg components were assessed via Kendall's tau correlation (for $n < 30$) and plotted as correlation matrices (Supplementary Materials Figs. S1–S3). Values below the detection limit (DL) were proxied as one-half of the DL. Statistical significance was identified when $P < 0.05$. Selenium binds to organic Hg at a molar ratio of 1:1; therefore the molar ratio was calculated for each sample using

$$\text{Molar Ratio} = \frac{[\text{Se}]/78.96 \text{ g/mol}}{[\text{Hg}]/200.59 \text{ g/mol}},$$

where 78.96 g/mol and 200.59 g/mol represent the atomic masses of Se and Hg, respectively.

RESULTS

Summary statistics for egg component element data are presented by sample type (eggshell, albumen, and yolk) in Supplementary Materials Tables S2–S4. All 15 metals and Se were detected in yolk ($n=14$). Albumen ($n=14$) and eggshell ($n=18$) contained detectable levels of 13 of the 16 elements: Al, Cd, Co, Cr, Cu, Fe, Mn, Ni, Pb, Se, Sn, V, and Zn. Arsenic, Hg, and Mo were below the detection limit in eggshells, and Mo could not be detected in albumen. Element distributions for each sample type (yolk, albumen, eggshell) are displayed in Fig. 1.

Molar Se:Hg ratios exceeded 1:1 for all yolk and albumen samples (Supplementary Materials Table S5). Mercury was not detected in eggshell, precluding molar Se:Hg ratio calculation for this component.

Yolk exhibited the highest geometric mean concentrations (µg/g) for As (0.20 ± 0.064), Co (0.01 ± 0.003), Cr (0.086 ± 0.03), Fe (238.65 ± 54.72), Mn (2.71 ± 0.66), Mo (0.57 ± 0.14), Pb (0.32 ± 0.97), Sn (3.29 ± 0.99), and Zn (64.03 ± 13.01 ; Table 1). Albumen had the highest concentrations of Al (4.19 ± 2.52), Cd (0.0009 ± 0.001), Cu (4.02 ± 1.22), Hg (0.62 ± 0.22), Se (1.52 ± 0.45), and V (0.0015 ± 0.03 ; Table 1). Eggshell had the highest concentration of Ni (0.31 ± 1.29 ; Table 1).

Statistically significant differences between eggshell and yolk occurred for all metals except




FIGURE 1. Geometric mean concentrations ($\mu\text{g/g}$) for aluminum (Al), arsenic (As), cadmium (Cd), chromium (Cr), cobalt (Co), copper (Cu), iron (Fe), lead (Pb), manganese (Mn), mercury (Hg), molybdenum (Mo), nickel (Ni), selenium (Se), tin (Sn), vanadium (V), and zinc (Zn) across egg component types for eggs from Humboldt Penguins (*Spheniscus humboldti*). Concentration data outliers are represented as circles above the maximum value.

Al, Cd, and Pb. Eggshell and albumen statistical differences were found for all metals except Al, As, Fe, Mo, and Zn. Arsenic, Co, Fe, Hg, Mn, Mo, and Zn concentrations differed significantly between yolk and albumen (Table 2). Aluminum was the only metal to show no variation among egg components.

Correlation matrices (Supplementary Materials Figs. S1–S3) represent associations between heavy metals within egg components. In yolk, highly positive ($r > 0.7$) correlations were observed between Se and Sn, Se and Mn, Hg and Ni, Cr and Ni, Cd and Ni, Fe and Zn, and Cu and Zn; highly negative correlations ($r > -0.7$) were observed between Sn and Mo, Ni and Mo, Mo and Hg, and Cr and Mo. Albumen had highly positive correlations between Zn and Mn, Zn and Fe, Sn and Cr, Mn and Fe, Pb and Co, Ni and Co, and Pb

and Cd; highly negative correlations were not observed. Eggshell produced highly positive correlations between Se and Ni, V and Mn, Zn and Fe, Ni and Cd, and Pb and Al; highly negative correlations were not observed.

DISCUSSION

It is well established that wild penguins may act as reliable bioindicators of inorganic environmental pollution (Celis et al. 2014; Finger et al. 2015; Adkesson et al. 2018, 2019), but contaminant data are scarce from penguins in managed care. We therefore draw comparisons between our data and heavy metal concentrations reported for eggs and tissues from other bird species. The value of intertissue and interspecies heavy metal concentration comparisons in seabirds has been given by Ackerman et al.

TABLE 1. Summary of geometric mean and standard deviation for metal and selenium concentrations (μg/g) determined through inductively coupled plasma-mass spectrometry analysis of egg components of Humboldt Penguins (*Spheniscus humboldti*) from a managed-care population at the Brookfield Zoo, Chicago, Illinois, USA. The highest concentration for each element is denoted with boldface text.

Element ^a	Sample type		
	Eggshell	Albumen	Yolk
Al	3.61 ± 1.79	4.19 ± 2.52	3.86 ± 0.94
As	ND ^b	0.0164 ± 0.0328	0.195 ± 0.064
Cd	0.0007 ± 0.002	0.0009 ± 0.001	0.0006 ± 0.0007
Co	0.0024 ± 0.0023	0.0038 ± 0.0031	0.0097 ± 0.003
Cr	0.035 ± 0.015	0.063 ± 0.018	0.0864 ± 0.031
Cu	1.32 ± 0.33	4.02 ± 1.22	3.77 ± 0.52
Fe	4.04 ± 9.23	10.4 ± 16.03	239 ± 54.7
Hg	ND	0.618 ± 0.217	0.079 ± 0.061
Mn	0.498 ± 0.22	0.171 ± 0.219	2.71 ± 0.66
Mo	ND	ND	0.575 ± 0.144
Ni	0.314 ± 1.29	0.269 ± 0.688	0.0908 ± 0.058
Pb	0.0154 ± 0.008	0.0541 ± 0.071	0.322 ± 0.972
Se	0.304 ± 0.083	1.52 ± 0.45	1.37 ± 0.18
Sn	0.798 ± 0.085	2.78 ± 1.29	3.29 ± 0.99
V	0.0003 ± 0.0005	0.0015 ± 0.002	0.0021 ± 0.0025
Zn	3.00 ± 1.96	3.99 ± 4.59	64.0 ± 13.0

^a Aluminum (Al), arsenic (As), cadmium (Cd), chromium (Cr), cobalt (Co), copper (Cu), iron (Fe), lead (Pb), manganese (Mn), mercury (Hg), molybdenum (Mo), nickel (Ni), selenium (Se), tin (Sn), vanadium (V), and zinc (Zn).
^b ND = not detected.

(2016). Across three seabird species, Forster’s Terns (*Sterna forsteri*), Black-necked Stilts (*Himantopus mexicanus*), and American Avocets (*Recurvisrostra americana*), metal content in eggs was highly correlated with concentrations in maternal blood, kidney, muscle, liver, and head feathers. Although such relationships have yet to be quantified for Humboldt Penguins, Ackerman et al. (2016) validate the relevance of drawing comparisons between eggs and tissues from adult birds across taxa.

Selenium is not a metal, but selenium concentrations were measured because Hg exhibits a high binding affinity for Se and acts to inhibit Se-dependent enzymes vital to metabolic pathways in the brain and neuroendocrine system (Ralston and Raymond 2010). As such, ratios exceeding 1:1 may indicate a protective effect of Se against Hg toxicity (Berry and Ralston 2008). Alternatively, a molar ratio ≤1 would be indicative of all available Se being bound to Hg, potentially resulting in oxidative stress risk if any unbound Hg existed (Cáceres-Saez et al. 2013).

However, increased Se tissue concentrations (i.e., Se:Hg >1) preserves Se-dependent enzyme activity, subsequently mitigating Hg toxicity (Berry and Ralston 2008; Ralston et al. 2008; Ralston and Raymond 2010). As molar Se:Hg ratios exceeded 1:1 for all yolk and albumen samples, indicating that Hg present in these egg components was fully bound to Se, it is posited that the potential for oxidative stress due to mercury exposure during chick embryogenesis is reduced for Humboldt Penguins in our managed-care population.

Yolk

Of all egg components, yolk exhibited the highest concentrations for 10/16 elements; Mo was detected only in yolk and not in albumen or eggshell. The range of elements concentrated in yolk can be largely explained by the metabolomics of avian vitellogenesis. Lipid is the largest yolk constituent by dry mass (Nys and Guyot 2011). Certain metals, such as Mn and

TABLE 2. Kruskal-Wallis test outputs for aluminum (Al), arsenic (As), cadmium (Cd), chromium (Cr), cobalt (Co), copper (Cu), iron (Fe), lead (Pb), manganese (Mn), mercury (Hg), molybdenum (Mo), nickel (Ni), selenium (Se), tin (Sn), vanadium (V), and zinc (Zn) concentrations ($\mu\text{g/g}$) in egg components of Humboldt Penguins (*Spheniscus humboldti*) from a managed-care population at the Brookfield Zoo, Chicago, Illinois, USA. For comparisons where a significant difference in element content was found, the sample with higher concentration is denoted with boldface text.

Element ^a	Sample comparison	<i>H</i> value	<i>P</i> value	Significant difference?
Al	Eggshell + albumen	0.577	0.4474	No
	Eggshell + yolk	1.48	0.2242	No
	Yolk + albumen	0.475	0.4907	No
As	Eggshell + albumen	1.05	0.3051	No
	Eggshell + yolk	22.9	<0.0001	Yes
	Yolk + albumen	20.3	<0.0001	Yes
Cd	Eggshell + albumen	7.48	0.0062	Yes
	Eggshell + yolk	3.54	0.0601	No
	Yolk + albumen	0.762	0.3827	No
Co	Eggshell + albumen	3.9	0.0482	Yes
	Eggshell + yolk	20.1	<0.0001	Yes
	Yolk + albumen	14.5	0.0001	Yes
Cr	Eggshell + albumen	15.9	0.0007	Yes
	Eggshell + yolk	20.1	<0.0001	Yes
	Yolk + albumen	3.72	0.0536	No
Cu	Eggshell + albumen	22.2	<0.0001	Yes
	Eggshell + yolk	22.9	<0.0001	Yes
	Yolk + albumen	0.931	0.3346	No
Fe	Eggshell + albumen	2.55	0.1106	No
	Eggshell + yolk	22.9	<0.0001	Yes
	Yolk + albumen	20.3	<0.0001	Yes
Pb	Eggshell + albumen	6.67	0.0098	Yes
	Eggshell + yolk	1.67	0.1965	No
	Yolk + albumen	0.357	0.5503	No
Mn	Eggshell + albumen	13	0.0003	Yes
	Eggshell + yolk	22.9	<0.0001	Yes
	Yolk + albumen	20.3	<0.0001	Yes
Hg	Eggshell + albumen	22.9	<0.0001	Yes
	Eggshell + yolk	22.9	<0.0001	Yes
	Yolk + albumen	20.3	<0.0001	Yes
Mo	Eggshell + albumen	0.468	0.4941	No
	Eggshell + yolk	22.9	<0.0001	Yes
	Yolk + albumen	20.3	<0.0001	Yes
Ni	Eggshell + albumen	17.1	<0.0001	Yes
	Eggshell + yolk	17.1	<0.0001	Yes
	Yolk + albumen	0.076	0.7828	No
Se	Eggshell + albumen	22.9	<0.0001	Yes
	Eggshell + yolk	22.9	<0.0001	Yes
	Yolk + albumen	2.03	0.1543	No
Sn	Eggshell + albumen	16.8	<0.0001	Yes
	Eggshell + yolk	22.9	<0.0001	Yes
	Yolk + albumen	1.54	0.2140	No
V	Eggshell + albumen	9.01	0.0025	Yes
	Eggshell + yolk	12.8	0.0004	Yes
	Yolk + albumen	1.43	0.2322	No

TABLE 2. Continued.

Element ^a	Sample comparison	H value	P value	Significant difference?
Zn	Eggshell + albumen	0.175	0.6761	No
	Eggshell + yolk	22.9	<0.0001	Yes
	Yolk + albumen	20.3	<0.0001	Yes

^aAluminum (Al), arsenic (As), cadmium (Cd), chromium (Cr), cobalt (Co), copper (Cu), iron (Fe), lead (Pb), manganese (Mn), mercury (Hg), molybdenum (Mo), nickel (Ni), selenium (Se), tin (Sn), vanadium (V), and zinc (Zn).

Zn, are able to form lipophilic metal complexes that promote absorption and tissue penetration, increasing contaminant retention in yolk (Tjälve and Gottfroy 1991). These metals also tend to accumulate in organs responsible for toxin elimination, such as the liver (Das et al. 2003). The lipids vitellogenins and yolk-targeted very-low-density lipoproteins (VLDL_y) are synthesized in the liver, secreted into the circulatory system, and taken up within the ovary's follicles for generation of yolk, the nutrient source for the developing embryo (Walzem 1996; Vezina et al. 2003). Once transported to the oocyte, vitellogenin is catabolized to phosvitin and lipovitellin, the predominant proteins found in yolk (Hermier et al. 1989; Speake et al 1998). Phosvitin exhibits a high capacity to bind essential metals (e.g., Ca and Fe), hence their enrichment in yolk (Strixner and Kulozik 2011).

During yolk development, ~17 d in Humboldt Penguins (Ancel et al. 2013), hepatic lipid production increases as lipoprotein synthesis shifts to generate the smaller, lipid-rich VLDL_y molecules required to support an embryo. As plasma vitellogenin concentrations increase, a corresponding rise in trace mineral concentrations occurs due to vitellogenin-metal binding (Hill 1974; Panic et al. 1974; Richards 1989). Increases in VLDL_y and vitellogenin production by the liver necessitate a corresponding increase in hepatocyte metabolism (Vezina et al. 2003). Richards (1997) attributes yolk heavy metal content almost entirely to vitellogenesis-driven increases in hepatic metabolism, which in turn facilitate maternal transfer of contaminants to the developing egg via the circulatory system.

The trends in yolk metal accumulation that we found have been seen in other birds, as

vitellogenesis mechanisms are conserved across avian species. Duck eggs from commercial farms in Thailand had yolk concentrations of Fe, Mn, and Zn an order of magnitude greater than those in albumen (Aendo et al. 2018), a trend reflected in our Humboldt Penguin eggs (Table 3). Humboldt Penguin yolk Cu, Hg, and Pb concentrations (Table 1) exceeded wet weight concentrations in Taiwanese duck eggs (Jeng and Yang 1994) by an order of magnitude of 19. However, concentrations in commercial chicken eggs (Demirulus 2013; Sarkar et al. 2018), were consistently higher than those we observed in Humboldt Penguin eggs; this might be attributable to herbicide or pesticide contamination of feed residues (Sarker et al. 2017; Table 3). Yolk Pb concentrations (0.322 µg/g; Table 1) in our study were similar to those found in the homogenized internal egg components of Great Cormorants (*Phalacrocorax carbo*) in Greece (Goutner et al. 2001). Although values for yolk of Pied Avocets (*Recurvirostra avosetta*), Yellow-legged Gulls (*Larus cachinnans michahellis*), and Mediterranean Gulls (*Larus melanocephalus*) from the Evros and Axios deltas in Greece were an order of magnitude greater than those penguin yolk, these birds also occupy habitats with considerable pollutant input from freshwater discharge generated by local agricultural complexes and commercial aquaculture ponds (Goutner et al. 2001). Compared to homogenized egg contents collected from Yellow-breasted Chats (*Icteria virens*) and Willow Flycatchers (*Empidonax traillii*) from Arizona, USA (Mora 2003), Humboldt Penguin yolk values exceeded wet weight concentrations of Cu, Mn, and Zn but were lower in Se (Table 3). This may be attributable to the managed-care Humboldt Penguins occupying a higher trophic niche than either

shorebird species, with contaminant biomagnification subsequently resulting in greater metal accumulation in the penguins' egg components.

Mean As (0.20 µg/g), Mn (2.71 µg/g), Pb (0.322 µg/g), and Zn (64.03 µg/g; Table 1) concentrations in Humboldt Penguin yolk fell within the range of mean concentrations reported for feathers from the Antarctic penguins Adelie Penguins (*Pygoscelis adeliae*; Dos Santos et al. 2006), Chinstrap Penguins (*Pygoscelis antarctica*; Jerez et al. 2013b), and Gentoo Penguins (*Pygoscelis papua*; Metcheva et al. 2006; Jerez et al. 2011, 2013a;). Although As, Mn, and Zn concentrations in kidney, liver, and muscle of these species exceed managed-care Humboldt Penguin yolk concentrations by up to an order of magnitude (Metcheva et al. 2010; Jerez et al. 2013a, b), the same tissues are significantly lower than yolk for Pb. Yolk Pb values were also similar to those in feathers of Magellanic Penguins (*Spheniscus magellanicus*) from the coast of southern Brazil (Kehrig et al. 2015), and yolk As and Zn concentrations were similar to those in feathers of Little Penguins (*Eudyptula minor*) from three island rookeries near Melbourne, Australia (Finger et al. 2015).

Yolk was an order of magnitude higher in Hg and two orders of magnitude higher in Zn than serum from wild Humboldt Penguins at PSJ but comparable in Al concentrations. Wild Humboldt Penguin feathers at PSJ had significantly higher concentrations of Al, As, Cu, Fe, Hg, and Mn than yolk from the eggs that we tested (Adkesson 2019). Humboldt Penguin excreta from three Chilean rookeries had mean As and Pb concentrations 40 times greater and mean Zn eight times higher (Celis et al. 2014) than in the yolks we tested (Table 4). Heightened metal concentrations in wild birds are to be expected because of the prevalence of mines and generalized anthropogenic activity throughout coastal Peru and Chile (Adkesson 2019).

Data for other heavy metals that we found to be highly concentrated in the yolk (Co [0.01 µg/g], Cr [0.086 µg/g], Mo [0.57 µg/g], and Sn [3.29 µg/g]; Table 1) are not well established in the literature for penguins.

Albumen

We found the highest concentrations of Al, Cd, Cu, Hg, and Se in albumen, a lipid-free fluid comprising primarily water and proteins, as well as vitamins and minerals required for embryonic development (Strixner and Kulozik 2011). Unlike yolk constituents, which are derived from components synthesized remotely (i.e., in the liver), albumen proteins are synthesized and secreted locally by the magnum of the oviduct (Nys and Guyot 2011). Ovalbumin and ovotransferrin, the two principal albumen proteins, form metal complexes that offer protection from bacterial infections (Schiafone and Barroeta 2011). Ovalbumin binds both Se and Hg (Magat and Sell 1979; Latshaw and Biggert 1981), as well as Cu, Mn, and Zn (Goux and Venkatasubramanian 1986). Ovotransferrin is principally responsible for Fe binding (Schiafone and Barroeta 2011), but Al, Cd, Cr, Co, Cu, Mn, and Zn bind in vitro (Burley and Vadehra 1989). The highly positive correlation between Mn and Fe in Humboldt Penguin albumen may result from Mn interacting with Fe cofactors on enzymes and impinging on transferrin receptor expression, limiting Fe endocytosis into red blood cell precursors, hepatocytes, and endothelial cells (Gupta et al. 2007).

Mercury levels in birds are well established due to the toxicity associated with its bioaccumulation and biomagnification. Humboldt Penguin mean Hg concentrations in albumen (0.62 µg/g; Table 1) were five times higher than levels found in wild Gentoo Penguin albumen (Brasso et al. 2012). This likely results from managed-care birds receiving higher trophic prey items than those foraged in the wild, because Hg levels in albumen from eggs laid by managed-care Gentoo Penguins are higher than albumen from wild birds (Brasso et al. 2012).

Notably, Hg in Humboldt Penguin albumen (0.62 µg/g; Table 1) far exceeded values in wild Magellanic Penguin feathers (0.033 µg/g; Frias et al. 2012) despite the species' nearly identical trophic dynamics and ecological niche (Simeone et al. 2009). Additionally, Humboldt Penguin albumen is comparable to values found in

TABLE 3. Mean metal concentrations (μg/g) in the egg components of various bird species compared to those in egg components of Humboldt Penguins (*Spheniscus humboldti*) from a managed-care population at the Brookfield Zoo, Chicago, Illinois, USA.

Species	Sample type	Al ^a	As ^a	Cd ^a	Cu ^a	Hg ^a	Mn ^a	Pb ^a	Zn ^a	Literature source
Domestic Duck (<i>Anas platyrhynchos domesticus</i>)	Albumen	— ^b	—	0.18	3.38	—	0.10	2.21	1.66	Aendo et al. (2018)
	Yolk	—	—	0.14	1.79	—	1.37	1.85	50.05	
Chicken (<i>Gallus gallus domesticus</i>)	Albumen	—	—	0.0018	0.83	0.0178	—	0.0136	—	Jeng and Yang (1995)
	Yolk	—	—	0.0038	0.00136	0.0097	—	0.0847	—	
	Albumen	—	—	0.78	4.2	—	3.2	—	5.8	Demirulus (2013)
	Yolk	—	—	0.79	6.6	—	4.3	—	38.9	
Willow Flycatcher (<i>Empidonax traillii</i>)	Albumen	—	0.160	0.039	0.899	—	—	0.106	—	Sarkar et al (2018)
	Yolk	—	0.157	0.042	0.882	—	—	0.123	—	
Yellow-breasted Chat (<i>Icteria virens</i>)	Homogenized internal components	—	<0.5	—	2.5	—	1.9	<0.5	38.5	Mora (2003)
	Homogenized internal components	—	<0.5	—	3.2	—	2.7	<0.5	52.4	
Great Cormorant (<i>Phalacrocorax carbo</i>)	Homogenized internal components	—	—	—	0.003	—	—	0.008	—	Gonthier et al. (2001)
	Homogenized internal components	—	—	—	0.006	—	—	0.07	—	
Pied Avocet (<i>Recurvirostra avosetta</i>)	Homogenized internal components	—	—	—	0.005	—	—	0.068	—	
Yellow-legged Gull (<i>Larus cachinnans michahellis</i>)	Homogenized internal components	—	—	—	0.003	—	—	0.020	—	
	Homogenized internal components	—	0.224	0.001	6.583	0.137	1.493	0.014	51.38	Lam et al. (2005)
Black-crowned Night Heron (<i>Nycticorax nycticorax</i>)	Shell	—	—	0.006	1.602	0.071	4.164	0.152	9.636	
	Homogenized internal components	—	—	—	6.286	0.520	1.720	0.007	39.87	
Bridled Tern (<i>Sterna anaethetus</i>)	Shell	—	—	0.008	1.116	0.056	1.031	0.030	5.390	
	Homogenized internal components	—	1.38	0.002	3.920	0.562	2.636	0.010	47.62	
	Shell	—	0.397	0.002	1.235	0.004	1.112	0.060	2.351	

TABLE 3. Continued.

Species	Sample type	Al ^a	As ^a	Cd ^a	Cu ^a	Hg ^a	Mn ^a	Pb ^a	Zn ^a	Literature source
Humboldt Penguin (<i>Spheniscus humboldti</i>)	Shell	—	3.61	0.0007	1.32	—	0.498	0.0154	3.0	Present study
	Albumen	0.0164	4.19	0.0009	4.02	0.618	0.171	0.0541	3.99	
	Yolk	0.195	3.86	0.0006	3.77	0.079	2.71	0.322	64	

^a Arsenic (As), aluminum (Al), cadmium (Cd), copper (Cu), mercury (Hg), lead (Pb), and zinc (Zn).

^b — = not tested for this element.

Adelie (Carravieri et al. 2016) and Gentoo Penguin feathers (Dos Santos et al. 2006; Table 4).

Concentrations of Cu (4.02 µg/g), Hg (0.618 µg/g), and Pb (0.0541 µg/g) in Humboldt Penguin albumen were greater than those found in Thai (3.38 µg/g for Cu, 2.21 µg/g for Pb; Aendo et al. 2018), and Taiwanese duck eggs (0.00136 µg/g, 0.0178 µg/g, and 0.0146 µg/g, respectively; Jeng and Yang 1994). Additionally, penguin albumen had higher Cd levels than yolk, a trend also observed in ducks (Aendo et al. 2018) that is likely attributable to ovotransferrin binding (Burley and Vadehra 1989). However, much like for yolk, heavy metal concentrations in commercial chicken egg albumen have been found to be significantly greater than those in Humboldt Penguin egg albumen (Demirulus 2013; Sarkar et al 2018; Table 3).

The mean albumen Al (4.19 µg/g) that we found resembled concentrations in the kidneys and livers of Adelie, Chinstrap, and Gentoo Penguins (Jerez et al. 2013a), while mean Cd levels (0.0009 µg/g) were lower than values in these penguins across tissue types (Jerez et al. 2013a; Table 4). While mean Al concentrations in feathers from Humboldt Penguin at PSJ far exceeded mean egg albumen levels in the managed-care population, serum Al levels in the wild birds were approximately half that seen in albumen (Adkesson et al. 2019). Albumen also contained up to two orders of magnitude higher Hg and Zn than serum collected from wild Humboldt Penguins at PSJ. Concentrations of Se (1.52 µg/g) were highest in albumen; however, much like Co, Cr, Mo, Sn, and V in yolk, Se concentrations are not reported for penguins (Table 4).

Eggshell

Eggshell had the lowest concentration of all analyzed metals except Ni (0.31 µg/g). This was unexpected, as certain divalent metals, specifically Cd and Pb, mimic and replace Ca during shell biomineralization (Strixner and Kulozik 2011). The replacement of calcium by its mimics results in the inhibition of Ca precipitation in calcified tissues (Wada et al. 1995; Bridges and Zalups 2005), resulting in metal increases in eggshells (O’Flaherty 1991;

TABLE 4. Mean metal concentrations (µg/g) in the eggs and tissues of various penguin species compared to those in egg components of Humboldt Penguins (*Spheniscus humboldti*) from a managed-care population at the Brookfield Zoo, Chicago, Illinois, USA. For comparisons where an element was not tested, a dash is used in lieu of a concentration value.

Region	Species	Sample type	Al	As	Cd	Cu	Hg	Mn	Pb	Zn	Literature source
Antarctica	Adelie Penguin (<i>Pygoscelis adeliae</i>)	Feather	43.4	0.17	0.13	19.3	0.66	2.01	0.32	83.9	Jerez et al. (2013a, b); Carravieri et al. (2016)
		Kidney	4.09	0.44	0.2	14.8	—	11.18	0.05	234	Jerez et al. (2013a)
		Bone	—	0.13	0.01	0.96	—	8.31	0.04	227	
		Liver	6.81	0.6	0.06	92.1	—	12.01	0.04	133	
	Chinstrap Penguin (<i>Pygoscelis antarctica</i>)	Muscle	—	—	0.01	6.4	—	1.5	0.12	104	Smichowski et al. (2006)
		Feather	26	0.01	0.1	20.3	—	2.25	1.73	97.3	Jerez et al. (2011)
		Kidney	10.93	0.5	0.54	13.6	—	10.19	0.14	92.8	Jerez et al. (2013b)
		Bone	—	0.08	<0.001	0.71	—	12.5	0.14	235	Jerez et al. (2013a)
		Liver	15.52	0.47	0.11	132	—	11.42	0.18	132	Jerez et al. (2013b)
		Muscle	—	—	0.01	6.82	—	2.55	0.2	105	
		Feather	46	0.07	0.21	17	0.54	2.6	0.1	106	Dos Santos et al. (2006); Metcheva et al. (2006); Jerez et al. (2011)
	Gentoo Penguin (<i>Pygoscelis papua</i>)	Kidney	6.91	0.4	0.2	14.3	—	7.54	<0.001	125	Jerez et al. (2013a)
		Bone	—	0.19	0.001	1.15	—	11.01	<0.001	245	Barbosa et al. (2013)
		Liver	2.12	0.45	0.08	142	—	10.51	<0.001	52.9	Jerez et al. (2013a)
		Muscle	—	—	0.01	—	—	1.46	<0.001	106	
		Eggshell	28.9	—	<0.05	1.24	0.05	0.82	0.68; 0.75	4.07	Yin et al. (2008); Metcheva et al. (2011)
		Feather	—	—	—	—	1.98	—	—	—	Scheifler et al. (2005)
	King Penguin (<i>Aptenodytes patagonicus</i>)	Feather	—	—	—	—	2.1	—	—	—	Carravieri et al. (2016)
		Northern Rockhopper Penguin (<i>Eudyptes moseleyi</i>)	—	—	—	—	1.96	—	—	—	
		Southern Rockhopper Penguin (<i>Eudyptes chrysocome</i>)	—	—	—	—	2.24	—	—	—	Carravieri et al. (2013)
		Macaroni Penguin (<i>Eudyptes chrysolophus</i>)	—	—	—	—	—	—	—	—	

TABLE 4. Continued.

Region	Species	Sample type	Al	As	Cd	Cu	Hg	Mn	Pb	Zn	Literature source
Australia	Little Penguin (<i>Eudyptula minor</i>)	Feather	40.4	0.18	0.06	—	4.13	—	0.42	84.7	Finger et al. (2015)
South America	Magellanic Penguin (<i>Spheniscus magellanicus</i>)	Feather	—	—	—	—	0.033	—	0.14	—	Frias et al. (2012)
		Kidney	—	—	—	—	—	—	0.55	—	
		Liver	—	—	7.25	—	—	—	5.7	—	
Humboldt Penguin (<i>Spheniscus humboldti</i>)		Excreta	—	7.86	—	—	—	—	12.79	487	Celis et al. (2014)
		Serum	2.14	—	—	—	0.0024	—	—	0.57	Adkesson et al. (2019)
		Feather	67	5.0	—	10.5	1.21	5.7	—	48	
		Shell	3.61	—	0.0007	1.32	—	0.498	0.0154	3.00	Present study
		Albumen	4.19	0.0164	0.0009	4.02	0.618	0.171	0.0541	3.99	
		Yolk	3.86	0.195	0.0006	3.77	0.079	2.71	0.322	64	

^aAluminum (Al), arsenic (As), cadmium (Cd), chromium (Cr), cobalt (Co), copper (Cu), iron (Fe), lead (Pb), manganese (Mn), mercury (Hg), molybdenum (Mo), nickel (Ni), selenium (Se), tin (Sn), vanadium (V), and zinc (Zn).

Teodorova et al. 2003; Scheuhammer 1996). Accordingly, Cd and Pb are expected to concentrate in the eggshell but were found in greater concentrations in albumen and yolk.

Humboldt Penguin heavy metal concentrations in eggshell were comparable to Gentoo Penguin eggshells for Al, Cu, Mn and Zn; Gentoo Penguin eggshell had higher metal concentrations (Metcheva et al. 2011). Likewise, all analyzed metals were found in higher concentrations in the feathers and serum of wild penguins at PSJ than in our Humboldt Penguin eggs (Adkesson et al. 2019; Table 4).

However, our Humboldt Penguin eggshells were higher in As (3.61 µg/g) than eggshells collected from Bridled Terns (*Sterna anaethetus*; 0.397 µg/g) inhabiting the Pearl River Delta manufacturing complex near Hong Kong (Lam et al. 2005). Eggshells of Little Egrets (*Egretta garcetta*) and Black-crowned Night Herons (*Nycticorax nycticorax*) inhabiting the same industrial complex had higher quantities of Cd, Mn, and Pb, while Zn concentrations were within the same range (2.351 µg/g to 9.636 µg/g) as our Humboldt Penguin eggshells (3.00 µg/g; Table 3). Managed-care birds represent a population with a significantly reduced risk of environmental contaminant exposure, yet the penguins assessed in this study nevertheless had accumulated metals in their eggshells to an extent comparable to findings in the wild birds assessed by Lam et al. (2005). Thus, when considering Humboldt Penguin populations inhabiting rookeries near coastal mines in Peru and Chile, we conjecture that there is an immense potential for significant contaminant accumulation in wild penguin eggshells, because of the introduction of metals and other environmental contaminants from local ore generation.

Implications for wild Humboldt Penguins

A recent ecosystem health assessment of Peruvian marine protected areas (MPAs), Loaiza et al. (2022) designated Punta San Juan, home to the largest Peruvian Humboldt Penguin rookery, as having the highest incidence of adverse biological effects and highest metal concentrations for As, Cd, Cu, Ni, Pb,

and Zn in sediment and seston (particles suspended in water) out of all surveyed MPAs, because of its proximity to mines. Heavy metal exposure during embryonic development has been heavily implicated in teratogenesis and developmental deformities in birds (Gilani and Alibhai 1990). Accordingly, the potential for reductions in offspring viability due to heavy metal contamination in eggs is a concern for species exhibiting multidecadal population declines, such as the Humboldt Penguins. Our findings serve to substantiate the value of assessing eggs when conducting assessments of environmental contamination in these birds.

It is important to note that the penguins in this study remained in a controlled environment (i.e., consistent diet and environment) throughout oogenesis and egg laying. Variance in habitat, diet, and environmental exposure risk experienced by wild birds significantly influences their assimilation of inorganic contaminants, and this must be taken into consideration. Additionally, most published studies assessing heavy metal contamination in eggs have presented data in terms of eggshell and homogenized internal components (i.e., combined yolk and albumen), limiting the extent to which some comparisons could be made. In the future, quantification of heavy metal concentrations in wild Humboldt Penguin egg components should be pursued to fully assess concerns regarding heavy metal exposure and contamination to the chick embryo. By determining the extent of heavy metal offloading that occurs via egg production, linkages among contaminants, embryo viability, and the likelihood of post-hatch chick mortality may be established. Such data would significantly inform existing strategies for the management of wild Humboldt Penguins, facilitating the conservation of this species.

ACKNOWLEDGMENTS

The authors thank the Chicago Zoological Society veterinary and animal care staff at Brookfield Zoo for their assistance with sample collection. Heavy metal ICP-MS analyses were conducted by Alan M. Shiller and Melissa Gilbert at the University of Southern Mississippi's Center for Trace Analysis. Project

funding was provided by the Chicago Zoological Society's Chicago Board of Trade Endangered Species Fund.

SUPPLEMENTARY MATERIAL

Supplementary material for this article is online at <http://dx.doi.org/10.7589/JWD-D-22-00176>.

LITERATURE CITED

- Ackerman JT, Eagles-Smith CA, Herzog MP, Hartman CA. 2016. Maternal transfer of contaminants in birds: Mercury and selenium concentrations in parents and their eggs. *Environ Pollut* 210:145–154.
- Ackerman JT, Hartman CA, Herzog MP. 2017. Maternal transfer of mercury to songbird eggs. *Environ Pollut* 230:463–468.
- Ackerman JT, Herzog MP, Evers DC, Cristol DA, Kenow KP, Heinz GH, Lavoie RA, Brasso RL, Mallory ML, et al. 2019. Synthesis of maternal transfer of mercury in birds: Implications for altered toxicity risk. *Environ Sci Technol* 54:2878–2891.
- Adkesson MJ, Levensgood JM, Scott JW, Schaeffer DJ, Langan JN, Cárdenas-Alayza S, de la Puente S, Majluf P, Yi S. 2018. Assessment of polychlorinated biphenyls, organochlorine pesticides, and polybrominated diphenyl ethers in the blood of Humboldt Penguins (*Spheniscus humboldti*) from the Punta San Juan Marine Protected Area, Peru. *J Wildl Dis* 54:304–314.
- Adkesson MJ, Levensgood JM, Scott JW, Schaeffer DJ, Panno B, Langan JN, Cárdenas-Alayza S, James-Yi S. 2019. Analysis of toxic and essential elements in the blood and feathers of Humboldt Penguins (*Spheniscus humboldti*) at Punta San Juan, Peru. *J Wildl Dis* 55:438–443.
- Aendo P, Netvichian R, Tippayalak S, Sanguankiat A, Khuntamoon T, Songserm T, Tulayakul P. 2018. Health risk contamination of heavy metals in yolk and albumen of duck eggs collected in central and western Thailand. *Biol Trace Elem Res* 184:501–507.
- Ancel A, Beaulieu M, Gilbert C. 2013. The different breeding strategies of penguins: A review. *C R Biol* 336:1–12.
- Barbosa A, De Mas E, Benzal J, Diaz J, Motas M, Jerez S, Pertierra L, Benayas J, Justel A, et al. 2013. Pollution and physiological variability in gentoo penguins at two rookeries with different levels of human visitation. *Antarct Sci* 25:329–338.
- Berry MJ, Ralston NVC. 2008. Mercury toxicity and the mitigating role of selenium. *Ecohealth* 5:456–459.
- BirdLife International. 2020. *Spheniscus humboldti*. The IUCN Red List of Threatened Species 2020: e.T22697817A182714418. doi: <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T22697817A182714418.en>. Accessed November 2023.
- Brasso RL, Abel S, Polito MJ. 2012. Pattern of mercury allocation into egg components is independent of dietary exposure in Gentoo penguins. *Arch Environ Contam Toxicol* 62:494–501.

- Bridges CC, Zalups RK. 2005. Molecular and ionic mimicry and the transport of toxic metals. *Toxicol Appl Pharmacol* 204:274–308.
- Burger J. 1994. Heavy metals in avian eggshells: Another excretion method. *J Toxicol Environ Health* 41:207–220.
- Burger J. 2002. Food chain differences affect heavy metals in bird eggs in Barnegat Bay, New Jersey. *Environ Res* 90:33–39.
- Burger J, Gochfeld M. 2004. Marine birds as sentinels of environmental pollution. *EcoHealth* 1:263–274.
- Burley RW, Vadehra DV. 1989. *The avian egg: Chemistry and biology*. John Wiley and Sons, New York, New York.
- Cáceres-Saez I, Dellabianca NA, Goodall RNP, Cappozzo HL, Guevara SR. 2013. Mercury and selenium in subantarctic Commerson's dolphins (*Cephalorhynchus c. commersonii*). *Biol Trace Elem Res* 151:195–208.
- Carravieri A, Bustamante P, Churlaud C, Cherel Y. 2013. Penguins as bioindicators of mercury contamination in the Southern Ocean: Birds from the Kerguelen Islands as a case study. *Sci Total Environ* 454–455:141–148.
- Carravieri A, Cherel Y, Jaeger A, Churlaud C, Bustamante P. 2016. Penguins as bioindicators of mercury contamination in the southern Indian Ocean: Geographical and temporal trends. *Environ Pollut* 213:195–205.
- Castilla J, Correa J. 1997. Copper tailing impacts in coastal ecosystems of northern Chile: From species to community responses. Copper. Report of an International Meeting 20–21 June 1996, Brisbane, Australia, Moore M, Imray P, Dameron C, Callan P, Langley A, Mangas S, editors. National Environmental Health Forum, Metal Series No. 3, Brisbane, Australia, pp. 71–80.
- Celis J, Espejo W, González-Acuña D, Jara S, Barra R. 2014. Assessment of trace metals and porphyrins in excreta of Humboldt Penguins (*Spheniscus humboldti*) in different locations of the northern coast of Chile. *Environ Monit Assess* 186:1815–1824.
- Colabuono FI, Vander Pol SS, Huncik KM, Taniguchi S, Petry MV, Kucklick JR, Montone RC. 2016. Persistent organic pollutants in blood samples of Southern Giant Petrels (*Macronectes giganteus*) from the South Shetland Islands, Antarctica. *Environ Poll* 216:38–45.
- Das K, Debacker V, Pillet S, Bouquegneau J. 2003. Heavy metals in marine mammals. In: *Toxicology of marine mammals*, Vos JG, Bossart G, Fournier M, O'Shea T, editors. Taylor and Francis, London, UK, pp. 135–154.
- Demirulus H. 2013. The heavy metal content in chicken eggs consumed in Van Lake territory. *Ekoloji Dergisi* 22:19–25.
- Dos Santos IR, Silva-Filho EV, Schaefer C, Sella SM, Silva CA, Gomes V, Passos MJACR, Van Ngan P. 2006. Baseline mercury and zinc concentrations in terrestrial and coastal organisms of Admiralty Bay, Antarctica. *Environ Poll* 140:304–311.
- Duffy DC. 1983. Competition for nesting space among Peruvian guano birds. *Auk* 100:680–688.
- Ferguson JE. 1990. *The heavy elements: Chemistry, environmental impact and health effects*. Pergamon Press, Oxford, UK.
- Finger A, Lavers JL, Dann P, Nugegoda D, Orbell JD, Robertson B, Scarpaci C. 2015. The little penguin (*Eudyptula minor*) as an indicator of coastal trace metal pollution. *Environ Poll* 205:365–377.
- Frias JE, Gil MN, Esteves JL, García Borboroglu P, Kane OJ, Smith JR, Boersma PD. 2012. Mercury levels in feathers of Magellanic penguins. *Mar Pollut Bull* 64:1265–1269.
- Fry DM. 1995. Reproductive effects in birds exposed to pesticides and industrial chemicals. *Environ Health Perspect* 103(Suppl. 7):165–171.
- Gilani SH, Alibhai Y. 1990. Teratogenicity of metals to chick embryos. *J Toxicol Environ Health* 30:23–31.
- Goutner V, Papagiannis I, Kalfakakou V. 2001. Lead and cadmium in eggs of colonially nesting waterbirds of different position in the food chain of Greek wetlands of international importance. *Sci Total Environ* 267:169–176.
- Goux WJ, Venkatasubramanian PN. 1986. Metal binding properties of hen ovalbumin and S-ovalbumin: Characterization of the metal ion binding site by ³¹P NMR and water protein relaxation rate enhancements. *Biochemistry* 25:84–94.
- Grace EJ, MacFarlane GR. 2016. Assessment of the bioaccumulation of metals to chicken eggs from residential backyards. *Sci Total Environ* 563–564:256–260.
- Gupta S, Stroh GM, Hassouna H. 2007. Positive influence of supplemental manganese on iron incorporation in brain and red blood cells. *J Investig Med* 55:355.
- Hermier D, Folgez P, Williams J, Chapman MJ. 1989. Alterations in plasma lipoproteins and apolipoproteins associated with oestrogen-induced hyperlipidemia in the laying hen. *Eur J Biochem* 184:109–118.
- Hill R. 1974. Changes in circulating copper, manganese and zinc with the onset of lay in the pullet. In: *Trace Elements in Man and Animals. Vol. 2. Proceedings of the second International Symposium on Trace Element Metabolism in Animals*, Madison, Wisconsin, 18–22 June 1973; Hoekstra WG, Suttie JW, Ganther HE, Mertz W, editors. University Park Press, Baltimore, MD, pp. 632–634.
- Holcman A, Stibilj V. 1997. Arsenic residues in eggs from laying hens fed with a diet containing arsenic (III) oxide. *Arch Environ Contamin Toxicol* 32:407–410.
- Jeng SL, Yang CP. 1995. Determination of lead, cadmium, mercury and copper concentrations in duck eggs in Taiwan. *Poultry Sci* 74:187–193.
- Jerez S, Motas M, Benzal J, Diaz J, Barbosa A. 2013a. Monitoring trace elements in Antarctic penguin chicks from south Shetland Islands, Antarctica. *Mar Pollut Bull* 69:67–75.
- Jerez S, Motas M, Benzal J, Diaz J, Vidal V, D'Amico V, Barbosa A. 2013b. Distribution of metals and trace elements in adult and juvenile penguins from the Antarctic peninsula area. *Environ Sci Pollut Res Int* 20:3300–3311.
- Jerez S, Motas M, Palacios MJ, Valera F, Cuervo JJ, Barbosa A. 2011. Concentration of trace elements in feathers of three Antarctic penguins: Geographical and interspecific differences. *Environ Pollut* 159:2412–2419.
- Kabeer MS, Hameed I, Kashif SU, Khan M, Tahir A, Anum F, Khan S, Raza S. 2021. Contamination of heavy metals in poultry eggs: A study presenting relation between heavy metals in feed intake and eggs. *Arch Environ Occup Health* 76:220–232.
- Kehrig HA, Hauser-Davis RA, Seixas TG, Fillmann, G. 2015. Trace-elements, methylmercury and metallothionein levels in Magellanic penguin (*Spheniscus*

- magellanicus*) found stranded on the southern Brazilian coast. *Mar Poll Bull* 96:450–455.
- Lam JCW, Tanabe S, Lam MHW, Lam PKS. 2005. Risk to breeding success of waterbirds by contaminants in Hong Kong: Evidence from trace elements in eggs. *Environ Pollut* 135:481–490.
- Latshaw JD, Biggert M. 1981. Incorporation of selenium into egg proteins after feeding selenomethionine or sodium selenite. *Poultry Sci* 60:1309–1313.
- Loaiza I, De Boeck G, De Troch M. 2022. Peruvian marine ecosystems under metal contamination: First insights for marine species consumption and sustainable management. *Sci Total Environ* 826:154132.
- Magat W, Sell JL. 1979. Distribution of mercury and selenium in egg components and egg-white proteins. *Proc Soc Exp Biol Med* 161:458–463.
- Metcheva R, Yurukova L, Teodorova SE. 2011. Biogenic and toxic elements in feathers, eggs, and excreta of gentoo penguin (*Pygoscelis papua ellsworthii*) in the Antarctic. *Environ Monit Assess* 182:571–585.
- Metcheva R, Yurukova L, Teodorova S, Nikolova E. 2006. The penguin feathers as bioindicator of Antarctica environmental state. *Sci Total Environ* 362:259–265.
- Ministerio de Energía y Minas. 2020. Peru: Mining country. <https://mineria.minem.gob.pe/en/institucional/peru-mining-country/>. Accessed October 2022.
- Mora MA. 2003. Heavy metals and metalloids in egg contents and eggshells of passerine birds from Arizona. *Environ Poll* 125:393–400.
- Nys Y, Guyot N. 2011. Chapter six: Egg formation and chemistry. In: *Improving the safety and quality of eggs and egg products*, Nys Y, Bain M, Immerseel FV, editors. Woodhead Publishing, Cambridge, UK, pp. 83–132.
- O'Flaherty EJ. 1991. Physiologically based models for bone-seeking elements. II. Kinetics of lead disposition in rats. *Toxicol Appl Pharmacol* 111:313–331.
- Panic B, Bezbradica L, Nedeljkov N, Istwari AG 1974. Some aspects of trace element metabolism in poultry. In: *Trace element metabolism in animals. Volume 2. Proceedings of the second International Conference on Trace Element Metabolism in Animals*. Hoekstra WG, Suttie JW, Ganther HE, Mertz W, editors. University Park Press, Baltimore, Maryland, pp. 635–637.
- Paredes R, Zavalaga CB. 2001. Nesting sites and nest types as important factors for the conservation of Humboldt Penguins (*Spheniscus humboldti*). *Biol Conserv* 100:199–205.
- Paredes R, Zavalaga CB, Bastistine G, Majluf P, McGill P. 2003. Status of the Humboldt Penguin in Peru 1999–2000. *Waterbirds* 26:129–138.
- Paredes R, Zavalaga CB, Boness DJ. 2002. Patterns of egg laying and breeding success in Humboldt Penguins (*Spheniscus humboldti*) at Punta San Juan, Peru. *Auk* 119:244–250.
- Ralston NVC, Raymond LJ. 2010. Dietary selenium's protective effects against methylmercury toxicity. *Toxicology* 278:112–123.
- Ralston NVC, Raymond CR, Blackwell JL 3rd, Raymond LJ. 2008. Dietary and tissue selenium in relation to methylmercury toxicity. *Neurotoxicology* 29:802–811.
- R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>. Accessed April 2022.
- Richards MP. 1997. Trace mineral metabolism in the avian embryo. *Poult Sci* 76:152–164.
- Richards MP. 1989. Influence of egg production on zinc, copper and iron metabolism in the turkey hen (*Meleagris gallopavo*). *Comp Biochem Physiol A Comp Physiol* 93:811–817.
- Sarkar A, Hossain MM, Sarkar AC. 2018. Heavy metal residue in eggs of chicken (*Gallus gallus domesticus*) available in Sylhet. In: Fifth International Conference on Chemical Engineering, Dhaka, Bangladesh, 20–22 December 2017, pp. 146–154.
- Sarker MS, Quadir QF, Hossen MZ, Nazneed T, Rahman A. 2017. Evaluation of commonly used fertilizers, fish and poultry feeds as potential sources of heavy metals contamination in food. *Asian Australas J Food Saf Secur* 1:74–81.
- Scheifler R, Gauthier-Clerc M, Le Bohec C, Crini N, Cœurdassier M, Badot PM, Giraudeau P, Le Maho Y. 2005. Mercury concentrations in king penguin (*Aptenodytes patagonicus*) feathers at Crozet Islands (sub-Antarctic): temporal trend between 1966–1974 and 2000–2001. *Environ Toxicol Chem* 24:125–128.
- Scheuhammer AM. 1987. The chronic toxicity of aluminum, cadmium, mercury, and lead in birds: A review. *Environ Poll* 46:263–295.
- Scheuhammer A. 1996. Influence of reduced dietary calcium on the accumulation and effects of lead, cadmium and aluminum in birds. *Environ Poll* 94:337–343.
- Schiavone A, Barroeta AC. 2011. Egg enrichment with vitamins and trace minerals. In: *Improving the safety and quality of eggs and egg products. Vol. 2. Egg safety and nutritional quality*, Nys Y, Bain M, Immerseel FV, editors. Woodhead Publishing, Cambridge, UK, pp. 289–320.
- Simeone A, Hiriart-Bertrand L, Reyes-Arriagada R, Halpern M, Dubach J, Wallace R, Pütz K, Lüthi B. 2009. Heterospecific pairing and hybridization between wild Humboldt and Magellanic Penguins in southern Chile. *Condor* 11:544–550.
- Smichowski P, Vodopivec C, Muñoz-Olivas R, Gutierrez AM. 2006. Monitoring trace elements in selected organs of Antarctic penguin (*Pygoscelis adeliae*) by plasma-based techniques. *Microchem J* 82:1–7.
- Speake BK, Noble RC, Murray AMB. 1998. The utilization of yolk lipids by the chick embryo. *World's Poult Sci J* 54:319–334.
- Strixner T, Kulozik U. 2011. Egg proteins. In: *Handbook of food proteins*, Phillips GO, Williams PA, editors. Woodhead Publishing, Cambridge, UK, pp. 150–209.
- Taylor MH, Tam J, Blaskovic V, Espinosa P, Ballón RM, Wosnitza-Mendo C, Argüelles J, Diaz E, Purca, S, et al. 2008. Trophic modeling of the Northern Humboldt Current Ecosystem, Part II: Elucidating ecosystem dynamics from 1995–2004 with a focus on the impact of ENSO. *Prog Oceanogr* 79:266–378.
- Teodorova SE, Metcheva R, Topashka-Ancheva M. 2003. Bioaccumulation and damaging action of polymetal industrial dust on laboratory mice *Mus musculus alba* I. Analysis of Zn, Cu, Pb and Cd disposition and mathematical model for Zn and Cd bioaccumulations. *Environ Res* 91:85–94.

- Thiel M, Macaya EC, Acuña E, Arnts WE, Bastias H, Brokordt K, Camus PA, Castilla JC, Castro L, et al. 2007. The Humboldt current system of northern and central Chile: Oceanographic processes, ecological interactions and socioeconomic feedback. *Oceanogr Mar Biol* 45:195–344.
- Tjälve H, Gottofrey J. 1991. Effects of lipophilic complex formation on the uptake and distribution of some metals in fish. *Pharmacol Toxicol* 69:430–439.
- Vezina F. 2003. The metabolic cost of avian egg formation: Possible impact of yolk precursor production? *J Exp Biol* 206:4443–4451.
- Wada N, Yamashita K, Umegaki T. 1995. Effects of divalent cations upon nucleation, growth and transformation of calcium carbonate polymorphs under conditions of double diffusion. *J Cryst Growth* 148:297–304.
- Walzem RL. 1996. Lipoproteins and the laying hen: Form follows function. *Poult Avian Biol Rev* 7:31–64.
- Yin X, Xia L, Sun L, Luo H, Wang Y. 2008. Animal excrement: A potential biomonitor of heavy metal contamination in the marine environment. *Sci Total Environ* 399:179–185.
- Zavalaga CB, Paredes S. 1997. Humboldt Penguins at Punta San Juan, Peru. *Penguin Conserv* 10:6–8.

Submitted for publication 22 December 2022.

Accepted 20 October 2023.