

SEROPREVALENCE OF ERYSIPELOTHRIX RHUSIOPATHIAE IN BEAUFORT SEA POLAR BEARS (URSUS MARITIMUS) IS LINKED TO RINGED SEAL (PUSA HISPIDA) DEMOGRAPHICS

Authors: Biddlecombe, Brooke A., Pilfold, Nicholas W., Richardson, Evan S., Kutz, Susan, Mavrot, Fabien, et al.

Source: Journal of Wildlife Diseases, 61(1): 76-87

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/JWD-D-24-00078

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commmercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Seroprevalence of *Erysipelothrix rhusiopathiae* in Beaufort Sea Polar Bears (*Ursus maritimus*) is Linked to Ringed Seal (*Pusa hispida*) Demographics

Brooke A. Biddlecombe,^{1,5} Nicholas W. Pilfold,² Evan S. Richardson,³ Susan Kutz,⁴ Fabien Mavrot,⁴ Angela Schneider,⁴ and Andrew E. Derocher¹

¹Biological Sciences Centre, University of Alberta, 11455 Saskatchewan Drive, Edmonton, Alberta T6G 2E9, Canada

² Conservation Science and Wildlife Health, San Diego Zoo Wildlife Alliance, Escondido, California 92027, USA

³ Wildlife Research Division, Science and Technology Branch, Environment and Climate Change Canada, 150-123 Main Street, Winnipeg, Manitoba R3C 1M8, Canada

⁴ Department of Ecosystem and Public Health, Faculty of Veterinary Medicine, University of Calgary, 2500 University

Drive NW, Calgary, Alberta T2N 1N4, Canada

⁵ Corresponding author (email: bbiddlec@ualberta.ca)

ABSTRACT: Polar bear (Ursus maritimus) life history is intimately associated with the distribution of sea ice and their prey in Arctic ecosystems. These ecosystems are changing in response to climate warming, resulting in the increased prevalence of pathogens in polar bears. Erysipelothrix rhusiopathiae has a long history of infection in domestic species and more recently in wildlife in the Canadian Arctic. As a result of increasing reports of E. rhusiopathiae causing morbidity and mortality in Arctic terrestrial mammals, we tested the seroprevalence of E. rhusiopathiae in Beaufort Sea polar bears sampled in 1985-87, 1992, 1994, and 2003-11. Our sample of 180 polar bears (117 females, 61 males, two unknown) with a median age of 9 yr (range 1-26 yr) had a seropositivity of 27.2% (49/ 180 individuals). We used binomial logistic regressions to investigate biotic and abiotic factors that may be linked to seropositivity. The resulting top model found that increased predation on adult ringed seals (*Pusa [Phoca] hispida*) and negative winter Arctic Oscillation Index (AOI) years were associated with a higher probability of seropositivity. Ringed seals may be a reservoir for *E*. rhusiopathiae via their consumption of infected prey, as the pathogen can persist in marine fish, molluscs, and crustaceans. Negative winter AOIs in our data set reflected high ice volume years, which reduced ringed seal natality, resulting in fewer seal pups available as prey. Our results suggest that exposure to E. rhusiopathiae in Beaufort Sea polar bears is modulated by a predator-prey mechanism.

Key words: Beaufort Sea, Erysipelothrix rhusiopathiae, emerging disease, polar bears, ringed seals.

INTRODUCTION

Erysipelothrix rhusiopathiae, an opportunistic and generalist bacterium, is most widely known for infecting domestic pigs and poultry but has also been observed in many wildlife species (Kutz et al. 2015; Forde et al. 2016; Mavrot et al. 2022). *Erysipelothrix rhusiopathiae* is a gram-positive bacillus with a wide distribution (Wang et al. 2010), infecting a diversity of both marine and terrestrial vertebrates (Conklin and Steele 1979). A unique strain of this bacterium has emerged in the Canadian Arctic Archipelago as an important cause of widespread mortality contributing to population declines in muskoxen (*Ovibos moschatus*) and has been detected in the feces

and carcasses of multiple Arctic foxes (Vulpes lagopus), wolves (Canis lupus), seals (Phocidae), and caribou (Rangifer tarandus; Kutz et al. 2015; Forde et al. 2016; Seru 2023). Different, or uncharacterized, strains of E. rhusiopathiae have also been detected serologically or by culture from muskoxen, caribou, moose (Alces alces), and seals elsewhere in Canada and Alaska, and in Pribilof Island foxes (Vulpes lagopus ssp. pribilofensis; Forde et al. 2016; Spraker and White 2017; Mavrot et al. 2020; Aleuy et al. 2022; Sudlovenick et al. 2023; S. Kutz, unpubl. data). In domestic animals, death can occur < 6 d after infection, but the mortality rate varies across species and individuals (Bobrek et al. 2013). Transmission may occur via contact through damaged skin or mucous membranes (Bobrek et al. 2013); from contact with bacteria shed in feces, urine, and saliva (Brooke and Riley 1999); or via insects as fomites (Wood and Steele 1994). Signs of infection typically include spots or lesions on the skin and regions of hair loss in mammals, and as infection progresses it may lead to sepsis (Peterman 1944; Bobrek et al. 2013). *Erysipelothrix rhusiopathiae* infection may also be maintained in an asymptomatic carrier state (Forde et al. 2016).

The first record of *E. rhusiopathiae* in the Arctic was connected to multiple widespread and severe muskox mortalities on Banks Island and Victoria Island in northwestern Canada (Kutz et al. 2015), with subsequent outbreaks first detected on the Canadian islands of Prince Patrick (2017), Ellesmere (2021–present), and Axel-Heiberg (2022–present). Nevertheless, *E. rhusiopathiae* is not new to the Arctic, with muskoxen seropositive to the bacteria as far back as the earliest available samples: 1976 in Alaska and 1991 in Canada (Mavrot et al. 2020).

Polar bears scavenge muskox remains (Voorhees et al. 2014), so the presence of E. rhusiopathiae in the Beaufort Sea region muskoxen suggests that it could be transmitted to polar bears via scavenging. The bacterium was identified in 14 polar bears from a sample of 48 hunter-harvested individuals across Canada during the development of a new method of disease identification (Tschritter et al. 2023). Concerns about exposure to an agent with a pathogenesis such as E. rhusiopathiae in polar bears increased after alopecia and reduced body condition were observed in 28% of bears sampled in the southern Beaufort Sea in 2012 (Atwood et al. 2015; Bowen et al. 2015). Elevated numbers of dead and sick ringed seals near Alaska were also reported during the same period (NOAA 2012). Although the cause of these events was not determined, they were suspected to be linked, given the close ecological relationship between polar bears and ringed seals. There are no studies of temporal trends of *E. rhusiopathiae* in polar bears; thus exposure dynamics are unknown.

Marine fish, molluses, and crustaceans are hosts for E. rhusiopathiae, and it can persist in tissues for long periods, surviving in decaying tissues for up to 10 mo and surviving in the exterior mucoid slime of fish without killing them (Brooke and Riley 1999). The Canadian Arctic Archipelago strain of E. rhusiopathiae has been cultured from a seal found dead on Victoria Island (Seru 2023), and ringed seals in the eastern Canadian Arctic were seropositive for E. rhusiopathiae (Sudlovenick et al. 2023), suggesting that seals may be a source of infection for polar bears through predation. Numerically, ringed seals comprise up to 90% of the prey of polar bears in the Beaufort Sea (Thiemann et al. 2008; Pilfold et al. 2012). Transmission from ringed seals is potentially related to ringed seal age because adult seals consume more fish, which can be carriers of the pathogen (Brooke and Riley 1999). Ringed seals rely on subnivean lairs on the ice for parturition and nursing young, but heavy ice years reduce productivity and lowers pupping success (Smith 1987; Harwood et al. 2000; Nguyen et al. 2017). Pilfold et al. (2012) found that when polar bear seal kills were observed, years of low pup productivity showed a higher proportion of adult ringed seals killed in the Beaufort Sea, with half of the adult seals >21 yr of age, raising the potential that polar bears were exposed to E. rhusiopathiae during these periods.

The Arctic is warming at four times the global average, with consequences for both species and ecosystems (Burek et al. 2008; Mallory et al. 2010; Johannessen and Miles 2011; Jacobs et al. 2021; Rantanen et al. 2022). Many of the climate change effects are associated with range shifts in species and changes in habitats, which includes sea ice habitats (Melles et al. 2011; Comiso 2012; Afenyo et al. 2019). Polar bears in the Beaufort Sea are among the most affected by changes in sea ice; have declined up to 50% in abundance (Bromaghin et al. 2015); and have been predicted to undergo further declines (Hunter et al. 2010) because of sea ice decline with ongoing warming (Stern and



FIGURE 1. Locations of polar bears (*Ursus maritimus*) sampled in the Beaufort Sea, Canada, 1985–2011 for *Erysipelothrix rhusiopathiae*. Black dots are seropositive individuals, white dots are negative. The line shows the delineation between northern and southern Beaufort Sea polar bear populations.

Laidre 2016). Sea ice decline in this region has the potential to make these bears more sensitive to other stressors (Atwood et al. 2016a). An increased prevalence of pathogen exposure has been linked to climate warming (Altizer et al. 2013; Baker-Austin et al. 2013; Cohen et al. 2020; Kafle et al. 2020; Aleuy et al. 2022; Peacock et al. 2022). In the Western Hudson Bay subpopulation, seroprevalence of five different pathogens has increased in polar bears over the past 3 decades; all were correlated to climate-mediated changes in the environment (Pilfold et al. 2021; Buhler et al. 2023).

The objective of our study was to assess the trends in the seroprevalence of *E. rhusiopa-thiae* in Beaufort Sea polar bears in Canada from 1985 to 2011, and to investigate biotic and abiotic factors that may be linked to infection. We hypothesized that *E. rhusiopathiae* seroprevalence in polar bears would be related to a predator–prey mechanism associated with ringed seals and sea ice patterns. We predicted positive relationships between *E. rhusiopathiae* seropositivity in polar bears

and environmental variables that related to predation of adult ringed seals. We also predict relationships between seropositivity and variables that act as proxies for climate change, such as increased seropositivity after years of earlier sea ice breakup.

METHODS

Sample collection

Polar bear sampling occurred in April and May in the Beaufort Sea in 1985-87, 1992, 1994; and 2003-11 in Canada (Fig. 1). The study area included polar bears from the Southern Beaufort Sea and Northern Beaufort Sea subpopulations (Paetkau et al. 1999; Amstrup et al. 2000; Stirling et al. 2011). Polar bears were located and immobilized from a helicopter via remote administration of tiletamine-olazepam (Telazol, Virbac, Carros, France) following Stirling et al. (1989). Blood samples were collected from the femoral vein, centrifuged at $1500 \times G$, and the serum and cellular portions were then frozen at -80 C until analysis. Age was determined by extracting a vestigial premolar and counting cementum annuli (Calvert and Ramsay 1998); dependent offspring were aged by body size and tooth eruption patterns.

For each sampled bear, sex, body condition, and capture location were recorded. Body condition was ranked from 1 (very skinny) to 5 (very fat) following Stirling et al. (2008b). Capture and handling protocols were approved by the Environment and Climate Change Canada's Western and Northern Animal Care Committee and the University of Alberta Biological Sciences Animal Policy and Welfare Committee, in accordance with the Canadian Council on Animal Care guidelines.

Seroprevalence analysis

To detect anti- Erysipelothrix antibodies in the polar bear sera we used an indirect ELISA developed by Giménez-Lirola et al. (2012) and further adapted to wildlife (Mavrot et al. 2020). Commonly used for seroprevalence analyses, ELISAs are highly sensitive (>97%) for determining antibody presence (Trono et al. 2001; Opsteegh et al. 2010). The test outcome is a percent positivity value (PP) of the sample compared with a positive control (serum from a pig vaccinated against *E. rhusiopathiae*). As for many wildlife species, no cutoff value to discriminate between positive and negative samples has been established. We used a mixture-distribution modeling approach to determine the best cutoff value (Mavrot et al. 2020). The PP cutoff for seropositivity was 55.1 (95% confidence interval [CI]: 46.1 - 73.6), where individuals above the cutoff but within the CI were deemed borderline positive and included as a positive in analyses.

Data analyses

Binomial logistic regressions were used to investigate the relationship between E. rhusiopathiae seropositivity and potentially predictive variables using RStudio (version 1.2.5033; R Core Team 2020). Positive seroprevalence samples were denoted as 1 and negative samples as 0. We included seven biotic and six abiotic covariates in regression models (Table 1). Following Pilfold et al. (2021), environmental variables were included with a 1-yr lag, suggesting that exposure reflected past conditions rather than at the sampling. Three initial models were chosen a priori to include different covariates to investigate different expected relationships: a biological model, a temporospatial model, and a ringed seal consumption model. A global model with all variables

together was also run. All models were linear mixed-effects models, with bear identification (ID) as the random variable, fit with the Nelder Mead optimizer. Pearson correlation between all numerical variables was tested, and any variables with a correlation ≥ 0.50 were not included in the same model. The only correlated variables were year and prey age (described below), with a correlation of 0.64, which were in separate a priori models.

Biological model: Body condition rankings were dummy coded into two binary covariates: a poor body condition covariate with rankings 1 and 2, and a good body condition covariate with rankings 4 and 5, with normal ranking (3) acting as a reference (Table 1). The biological model to investigate biological correlates with seropositivity included bear age, sex, poor body condition, and good body condition as fixed effects, and bear ID as a random effect.

Temporospatial model: The model to investigate potential temporal and spatial patterns included capture year, latitude and longitude of capture, number of low ice days per year, and the date of sea ice breakup as fixed effects; bear ID was a random effect. The length of the sea ice season has been linked to polar bear survival, most often through access to ringed seals (Stirling and Parkinson 2006; Regehr et al. 2007; Molnar et al. 2010; Hamilton et al. 2014). The length of the low ice season corresponds to the amount of time polar bears have a reduced ability to hunt seals, and the sea ice breakup date denotes the start of the low ice season, reflecting the date bears are forced off the ice, their primary habitat.

Ringed seal consumption model: The ringed seal consumption model focused on the suspected effects of ringed seal consumption as a source of exposure, including ringed seal prey age class (prey age, hereafter), female bears with cubs, Arctic Ocean Oscillation Index (AOO), and winter Arctic Oscillation Index (AOI) as fixed effects, and bear ID as a random effect. Prey age is an annual value, calculated by Pilfold et al. (2012). This model uses a proportional width index (PWI) of growth layer groups measured from ringed seal canine teeth that were killed by polar bears and then converted into a body condition index (see Nguyen et al. 2017). Nguyen et al. (2017) found PWI to be positively correlated with ringed seal ovulation rate; thus it may

Name	Range	Description and source	
Biotic			
Age (yr)	1-26	Age of polar bear via tooth histology (Calvert and Ramsay 1998)	
Sex	1/0	Field determination with females as reference category (0)	
Poor ^a	1/0	Polar bears rated 1 or 2 on 5-point body condition index (Stirling et al. 2008b)	
Good ^a	1/0	Polar bears rated 4 or 5 on 5-point body condition index (Stirling et al. 2008b)	
PreyAge	0-0.75	Proportion of seal kills detected by aerial surveys that were adult, ≥ 5 yr old (Pilfold et al. 2012)	
PWI^{b}	0.267-0.293	Index of ringed seal (<i>Pusa hispida</i>) body condition assessed on seal kills (Nguyen et al. 2017)	
CUB	1/0	Females with a cub-of-the-year the previous year	
Abiotic			
LowIce	86–166 d	Number of days of sea ice concentration <50% as determined by SSM/I (Cavalieri et al. 1996)	
BreakUp	136–199	Ordinal date that sea ice concentration dropped and remained <50% as determined by SSM/I (Cavalieri et al. 1996)	
AOO	1/0	Index of Arctic wind-driven circulation regimes (Proshutinsky and Johnson 1997), measured annually. Converted to a binary variable, where values <0 are assigned 1	
Winter_AOI	1/0	Index of sea-level pressure anomalies north of 20°N (Thompson and Wallace 1998), mean for January–March. Converted to a binary variable where values <0 are assigned 1	
Latitude	69.25-76.68	Location of polar bear capture	
Longitude	117.36 - 140.03	Location of polar bear capture	

TABLE 1. Covariates used to model the likelihood of *Erysipelothrix rhusiopathiae* seropositivity in polar bears (*Ursus maritimus*) of the Beaufort Sea, Canada, 1985–2011. All temporal variables were taken from the year before serum sample acquisition to reflect antibodies from exposure in the past year.

^a 5-point body condition index dummy-coded, with "average" (3) forming the reference category.

^b PWI = proportional width index; SSM/I = special sensor microwave/imager; AOO = Arctic Ocean Oscillation Index; AOO = Arctic Oscillation Index.

be associated with ringed seal predation trends in polar bears. Proportional width index data were only available until 2007 (Nguyen et al. 2017), so a separate ringed seal consumption model was run with a truncated time series to test potential effects of PWI. Both winter AOI and AOO may affect both body condition in ringed seals (Nguyen et al. 2017) and polar bear predation (Pilfold et al. 2015). Winter AOI and AOO were converted to binary variables where negative values were denoted as 1 and positive values were 0 because negative AOO and winter AOI years reflected higher ice volume years and were of greater interest. Females with cubs-ofthe-year (cubs) were included as a separate group because compared with other polar bears they were expected to feed on a higher proportion of seal pups (Stirling 2002; Stirling et al. 2008a).

Winter AOI and AOO were included in the ringed seal consumption model because they were associated with ringed seal natality (Nguyen et al. 2017) and polar bear predation patterns (Pilfold et al. 2015). Pilfold et al. (2015) described relationships between polar bear kill rates, ringed seal reproduction, ice concentration, and largescale atmospheric circulation patterns (i.e., winter AOI and AOO) and concluded that kill rates were influenced by a combination of abiotic and biotic factors.

Model selection: There is a recognized temporal pattern in sea ice cover that correlates with ringed seal natality and polar bear predation on ringed seals (Stirling 2002; Pilfold et al. 2015; Nguyen et al. 2017), so interpretation of model results considered relationships between these variables before determining a best-fit model. Temporal trends in prey age were investigated against temporal trends in



FIGURE 2. Comparison of *Erysipelothrix rhusiopathiae* seropositivity and ringed seal (*Pusa hispida*) prey age in Beaufort Sea, Canada, polar bears (*Ursus maritimus*) sampled 1985–2011 during years of normal and heavy sea ice volume.

seropositivity because of the correlation between year and prey age. When seropositivity was plotted over time, peaks and troughs closely matched peaks and troughs in prey age data, which correlated with periods of normal and heavy sea ice cover, respectively. This pattern was evident when the comparison between ice conditions was plotted (Fig. 2) and suggested that an effect from year could be explained, in part, by prey age, reflecting that the time series began in heavy ice years, where heavy ice reduced ringed seal productivity and ended in years of high ringed seal productivity (Nguyen et al. 2017). Because of the correlation between covariates and this identified trend, year was not included in the top model. $P \leq 0.05$ was considered statistically significant, and P = 0.05-0.10 was considered marginally significant for interpreting covariate effect. On the basis of the results from the initial models and Akaike information criterion with a correction for small sample size (AICc) values, where $\Delta AICc \geq 2$ lower than another model equated to better model fit, a top model was chosen.

RESULTS

Of the 180 polar bears sampled, the median age was 9 yr (range 1–26 yr) and the proportion of females was 0.66. Seropositivity was found in 49/180 (27.2% of samples; 95% CI 21.0–34.4%). Of those that were seropositive, 20 were classified as borderline seropositive, with a PP above the cutoff but within the CI.

The biological model found a marginally significant relationship with sex but no other significant effects (Supplementary Material Table S1). The temporospatial model found a significant negative relationship with year but no other significant effects (Supplementary Material Table S2). The ringed seal consumption model found a significant positive relationship with prey age, as well as marginal significance from winter AOI (Supplementary Material Table S3). The ringed seal model with subset data to include PWI had a marginally significant relationship with winter AOI and a positive significant relationship with prey age (Supplementary Material Table S4). The global model with all covariates included did not converge.

Comparing across all models with the full sample size, the ringed seal consumption model had the lowest AICc value (Table 2). The reduced sample size model (Supplementary Material Table S4) had a lower TABLE 2. Comparing model-fit Akaike information criterion with a correction for small sample size (AICc) values to inform model selection for models estimating the likelihood of *Erysipelothrix rhusiopathiae* seropositivity in polar bears (*Ursus maritimus*) of the Beaufort Sea, Canada, 1985–2011.

	_
Biological 212.	00
Temporospatial 209.	07
Ringed seal (<i>Pusa hispida</i>) consumption 171.	00
PWI ^a (reduced sample size) 124.	11
Final 167.	69

^a PWI = proportional width index.

AICc value but could not be compared directly because of differences in sample size; model results outside of AICc do not suggest that this was the best-fit model. The top model included prey age and winter AOI from the ringed seal consumption model, whereby prey age had a significant positive effect, and winter AOI was marginally significant (Table 3). This top model had a lower AICc (Δ AICc = 3.31) than the original ringed seal consumption model. A model variant was tested with prey age alone to compare with the final model, which had a similar but slightly reduced fit, with a Δ AICc increase of 1.43.

DISCUSSION

When considering infectious diseases, Arctic ecosystems remain data deficient, and as such, lack baseline data (Hoberg et al. 2008; Kutz et al. 2014). This study provides insight on the prevalence of E. rhusiopathiae over time in polar bears within the Arctic marine food web. We detected E. rhusiopathiae seropositivity in Beaufort Sea polar bears from our earliest samples (1985), suggesting that polar bears in the western Arctic have been exposed for several decades. Seropositivity does not equate to active infection as antibodies often persist longer than a transient infection, although they reduce over time (Gilbert et al. 2013). Antibody persistence varies depending on multiple factors associated with both the host and the pathogen and usually ranges between a few weeks and several years (Secundino et al. 2006; Gilbert et al. 2013). Anti-Erysipelothrix antibody persistence in polar bears is unknown and seropositivity in our samples provided no information on when the animal was exposed. As such, although we considered temporal variables, past events and current events may be confounded.

Our time series was limited, with many years lacking samples, and notable previously quantified trends in ringed seal natality and

TABLE 3. Results from the preferred logistic regression model investigating the effects of covariates on the probability of *Erysipelothrix rhusiopathiae* seropositivity in polar bears (*Ursus maritimus*) of the Beaufort Sea, Canada, 1985–2011.

		Fixed effects		
Covariate	Estimate	SE	Z value	<i>P</i> value
Intercept	-2.593	0.549	-4.727	<0.001 ^a
Prey age ^b	2.797	0.971	2.881	0.004^{a}
Winter AOI ^c	0.752	0.440	1.709	0.087 ^d
		Random effect		
Covariate		SD		
Bear identification		1.05		

^a Significant values (P < 0.01).

^b Proportion of ringed seal (Pusa hispida) prey in adult age class.

^c Annual winter Arctic Oscillation Index.

 ${}^{d}P = 0.05-0.10$ was considered marginally significant.

ice volume, so an expected trend of increased infection over time, as was observed in Western Hudson Bay polar bears (Pilfold et al. 2021), was not present but might have been masked by these other effects. Increased seal mortality during the end of our study period (Ferguson et al. 2017; Nguyen et al. 2017) may have also influenced polar bear exposure to E. rhusiopathiae, because although ringed seal dynamics were correlated with atmospheric patterns, we speculate that they could also be associated with undocumented E. rhusiopathiae infection in the seals. Similarly, the observed E. rhusiopathiae muskox epidemic may have reflected a broader ecosystem-level epidemic (Kutz et al. 2015) and could have had an influence on polar bear exposure through scavenging of dead muskoxen and other species.

The initial models each focused on aspects of polar bear ecology. Most covariates related to individual bears had no quantifiable effects on seropositivity, which is similar to the negligible effect individual traits had on E. rhusio*pathiae* seropositivity in Arctic caribou (Aleuy et al. 2022). Most of the spatial and temporal covariates also had no effect on seropositivity. There was marginal significance from sex in the biological model, suggesting that males were more likely to be seropositive. The lack of significance for latitude and longitude suggests that E. rhusiopathiae infection was widespread. There was a significant effect from year in the temporospatial model, which suggested a decline in E. rhusiopathiae seropositivity over time when considered alone. When considered with ringed seal prey age, we believe that this apparent decline was an artifact of changes in predation associated with changes in ringed seal natality, which was in turn associated with sea ice dynamics (Nguyen et al. 2017; Harwood et al. 2020). The relationship identified between polar bear seropositivity, ringed seal prey age, and ice conditions (Fig. 2) supported choosing prey age over year (they could not both be included because of correlation) for the top model despite its significance in the temporospatial model, as it

was probably an artifact of an effect that was better explained by prey age. Interannual variability in *E. rhusiopathiae* infection in caribou has been found to be related to climatic conditions and some waxing and waning in herd immunity (Aleuy et al. 2022), which could be a potential factor in interannual variability in polar bear infection that we were not able to quantify.

The top model was informed by a predator-prey relationship between polar bears and ringed seals, and included the effect of winter AOI, which was linked to ringed seal population dynamics. Polar bears that preyed on a higher proportion of adult ringed seals had a significantly higher probability of seropositivity. Adult seals may have a higher chance of E. rhusiopathiae infection because they consume fish, a host for the pathogen, whereas juveniles feed on milk. In a case of infection, it is important to note that seals can also be exposed, infected, and recover without retaining transmissibility, so older seals may not necessarily have higher transmissibility of E. rhusiopathiae to polar bears (VanderWaal and Ezenwa 2016), but as an age class (as is included in our model) we can speculate that transmissibility is probably higher in adults versus pups and juveniles. Winter AOI in the model focused on the effects of negative winter AOI years, which reflected years of increased sea ice volume. As such, the positive winter AOI coefficient estimated by the model suggested that probability of seropositivity increased after years with high ice volume. High ice volume was related to reduced pupping success in ringed seals and the age distribution of ringed seals killed by polar bears reflected the availability of pups (Pilfold et al. 2012), so reduced pupping success was connected to an increase in adult seal consumption, and in turn consumption of prey that we can speculate had a higher likelihood of E. rhusiopathiae transmission. Ringed seal consumption has been linked to the protozoan parasite Toxoplasma gondii in polar bears; this parasite has multiple potential vectors, one of which is the transmission of oocysts from runoff water into filter-feeding fish, which are prey for ringed seals (Massie et al. 2010; Simon et al. 2013). Multiple-host pathogens are generalists, which allows them to be passed through food webs and succeed in multiple taxa (Woolhouse et al. 2001; Lafferty et al. 2008), a probable explanation for the apparent movement of *E. rhusiopathiae* through both the terrestrial and marine food webs.

Multiple aspects of how predator-prey relationships interact with pathogen transmission have been studied, including the effect of predation on pathogen persistence in host species, bioaccumulation in apex predators, and diseases crossing the species barrier from prey to predator (Hsieh and Hsiao 2008; Moore et al. 2010; Su and Hui 2011; Malmberg et al. 2021). In systems where a pathogen can cross from prey to predator, many outcomes are possible, including eradication of the pathogen in the prey from targeted predation, or the pathogen driving the predators to extinction from increased predator infection (Hsieh and Hsiao 2008). The high seroprevalence found in polar bears might indicate that as an apex predator they act as a bioaccumulator for the disease and could be a potential reservoir for it, releasing it in the marine environment. Our results suggest that the food web is a driver of E. rhusiopathiae exposure in polar bears, but we do not have enough information within the polar bear ringed seal system to predict how both species are interacting with E. rhusiopathiae or how this pathogen may affect predator-prey dynamics into the future. Further information on the occurrence of the bacterium in sympatric seal and fish populations might shed light on whether E. rhusiopathiae is maintained in the food web of Beaufort Sea polar bears, versus being a spillover into polar bears from a terrestrial system (through scavenging behavior).

Understanding the prevalence of *E. rhusio-pathiae* in polar bears and Arctic ecosystems would be positively affected by an improved surveillance system, inclusive of northern, community-based wildlife health surveillance. Tschritter et al. (2023) have developed sensitive disease surveillance tools for applications

in Arctic ecosystems, which included *E. rhusiopathiae* detection, but these methods determine the presence of bacterial DNA, which may not necessarily equate to infection. Various strains of *E. rhusiopathiae* have been found throughout the Arctic; further work to uncover which strain is infecting polar bears, or if there are multiple strains, would be valuable. Progress in detection and inclusive surveillance systems for *E. rhusiopathiae* would help us understand the extent of exposure across the Arctic, and potentially elucidate transmission patterns between regions and species.

ACKNOWLEDGMENTS

We thank Ian Stirling, Dennis Andriashek, Wendy Calvert, and Cheryl Spencer for their invaluable contributions to monitoring polar bears in the Beaufort Sea. Financial support for this study was provided by Banrock Station Environmental Fund, Canadian Association of Zoos and Aquariums, Canadian Wildlife Federation, Environment and Climate Change Canada, Government of the Northwest Territories, Hauser Bears, Natural Sciences and Engineering Research Council of Canada, Parks Canada, Polar Bears International, the Polar Continental Shelf Project of Natural Resources Canada, Quark Expeditions, San Diego Zoo Wildlife Alliance, US Department of the Interior (Bureau of Ocean Energy Management), US Geological Survey Alaska Science Centre, Polar Knowledge Canada, Morris Animal Foundation, and World Wildlife Fund Canada. We thank the reviewers for their time and valued insight.

SUPPLEMENTARY MATERIAL

Supplementary material for this article is online at http://dx.doi.org/10.7589/JWD-D-24-00078.

LITERATURE CITED

- Afenyo M, Jiang C, Ng AK. 2019. Climate change and Arctic shipping: A method for assessing the impacts of oil spills in the Arctic. *Transp Res D Transp Envi*ron 77:476–490.
- Aleuy OA, Anholt M, Orsel K, Mavrot F, Gagnon CA, Beckmen K, Côté SD, Cuyler C, Dobson A, et al. 2022. Association of environmental factors with seasonal intensity of *Erysipelothrix rhusiopathiae* seropositivity among Arctic caribou. *Emerg Infect Dis* 28:1650–1658.
- Altizer S, Ostfeld RS, Johnson PTJ, Kutz S, Harvell CD. 2013. Climate change and infectious diseases: From evidence to a predictive framework. *Science* 341:514–519.

- Amstrup SC, Durner GM, Stirling I, Lunn NJ, Messier F. 2000. Movements and distribution of polar bears in the Beaufort Sea. Can J Zool 78:948–966.
- Atwood T, Peacock E, Burek-Huntington K, Shearn-Bochsler V, Bodenstein B, Beckmen K, Durner G. 2015. Prevalence and spatio-temporal variation of an alopecia syndrome in polar bears (*Ursus maritimus*) of the southern Beaufort Sea. J Wildl Dis 51:48–59.
- Atwood TC, Marcot BG, Douglas DC, Amstrup SC, Rode KD, Durner GM, Bromaghin JF. 2016a. Forecasting the relative influence of environmental and anthropogenic stressors on polar bears. *Ecosphere* 7:e01370.
- Baker-Austin C, Trinanes JA, Taylor NGH, Hartnell R, Siitonen A, Martinez-Urtaza J. 2013. Emerging Vibrio risk at high latitudes in response to ocean warming. Nat Clim Change 3:73–77.
- Bobrek K, Gaweł A, Mazurkiewicz M. 2013. Infections with *Erysipelothrix rhusiopathiae* in poultry flocks. *World's Poult Sci J* 69:803–812.
- Bowen L, Miles AK, Stott J, Waters S, Atwood T. 2015. Enhanced biological processes associated with alopecia in polar bears (Ursus maritimus). Sci Total Environ 529:114–120.
- Bromaghin JF, McDonald TL, Stirling I, Derocher AE, Richardson ES, Regehr EV, Douglas DC, Durner GM, Atwood T, Amstrup SC. 2015. Polar bear population dynamics in the southern Beaufort Sea during a period of sea ice decline. *Ecol Appl* 25:634–651.
- Brooke CJ, Riley TV. 1999. Erysipelothrix rhusiopathiae: Bacteriology, epidemiology and clinical manifestations of an occupational pathogen. J Med Microbiol 48:789–799.
- Buhler KJ, Dibernardo A, Pilfold NW, Harms NJ, Fenton H, Carriere S, Kelly A, Schwantje H, Aguilar XF et al. 2023. Widespread exposure to mosquitoborne California serogroup viruses in caribou, Arctic fox, red fox, and polar bears, Canada. *Emerg Infect Dis* 29:54–63.
- Burek KA, Gulland FM, O'Hara TM. 2008. Effects of climate change on Arctic marine mammal health. *Ecol Appl* 18(2 Suppl):S126–S134.
- Calvert W, Ramsay MA. 1998. Evaluation of age determination of polar bears by counts of cementum growth layer groups. Ursus 10:449–453
- Cavalieri DJ, Parkinson CL, Gloersen P, Zwally HJ. 1996. Sea ice concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS passive microwave data, version 1. NASA National Snow and Ice Data Center Distributed Active Archive Center, Boulder, Colorado. https:// doi.org/10.5067/8GQ8LZQVL0VL. Accessed February 2024.
- Cohen JM, Sauer EL, Santiago O, Spencer S, Rohr JR. 2020. Divergent impacts of warming weather on wildlife disease risk across climates. *Science* 370: eabb1702.
- Comiso JC. 2012. Large decadal decline of the Arctic multiyear ice cover. J Clim 25:1176–1193.
- Conklin RH, Steele J. 1979. Erysipelothrix infections. In: CRC handbook series in zoonoses, Steele JH, editor. CRC Press, Boca Raton, Florida, pp. 327–337.
- Ferguson SH, Young BG, Yurkowski DJ, Anderson R, Willing C, Nielsen O. 2017. Demographic, ecological, and physiological responses of ringed seals to

an abrupt decline in sea ice availability. PeerJ5: e2957.

- Forde TL, Orsel K, Zadoks RN, Biek R, Adams LG, Checkley SL, Davison T, De Buck J, Dumond M, et al. 2016. Bacterial genomics reveal the complex epidemiology of an emerging pathogen in Arctic and boreal ungulates. *Front Microbiol* 7:1759.
- Gilbert AT, Fooks AR, Hayman DTS, Horton DL, Müller T, Plowright R, Peel A, Bowen R, Wood JLN, et al. 2013. Deciphering serology to understand the ecology of infectious diseases in wildlife. *EcoHealth* 10:298–313.
- Giménez-Lirola LG, Xiao CT, Halbur PG, Opriessnig T. 2012. Development and evaluation of an enzymelinked immunosorbent assay based on a recombinant SpaA protein (rSpaA415) for detection of anti-*Ery*sipelothrix spp. IgG antibodies in pigs. J Microbiol Methods 91:191–197.
- Hamilton SG, Castro de la Guardia L, Derocher AE, Sahanatien V, Tremblay B, Huard D. 2014. Projected polar bear sea ice habitat in the Canadian Arctic Archipelago. *Plos One* 9: e113746.
- Harwood LA, Smith TG, Alikamik J, Alikamik E, Lea EV, Stirling I, Wright H, Melling H, Zhu X. 2020. Longterm, harvest-based monitoring of ringed seal body condition and reproduction in Canada's western Arctic. Arctic 73:206–220.
- Harwood LA, Smith TG, Melling H. 2000. Variation in reproduction and body condition of the ringed seal (*Phoca hispida*) in western Prince Albert Sound, NT, Canada, as assessed through a harvest-based sampling program. Arctic 53:422–431.
- Hoberg EP, Polley L, Jenkins EJ, Kutz SJ, Veitch AM, Elkin BT. 2008. Integrated approaches and empirical models for investigation of parasitic diseases in northern wildlife. *Emerg Infect Dis* 14:10–17.
- Hsieh Y-H, Hsiao C-K. 2008. Predator–prey model with disease infection in both populations. *Math Med Biol* 25:247–266.
- Hunter CM, Caswell H, Runge MC, Regehr EV, Amstrup SC, Stirling I. 2010. Climate change threatens polar bear populations: A stochastic demographic analysis. *Ecology* 91:2883–2897.
- Jacobs P, Lenssen N, Schmidt G, Rohde R. 2021. The Arctic is now warming four times as fast as the rest of the globe. In: AGU Fall Meeting Abstracts, American Geophysical Union, New Orleans, Louisiana, 13–17 December [Abstract].
- Johannessen OM, Miles MW. 2011. Critical vulnerabilities of marine and sea ice-based ecosystems in the high Arctic. *Reg Environ Change* 11(Suppl 1):S239– S248.
- Kafle P, Peller P, Massolo A, Hoberg E, Leclerc L-M, Tomaselli M, Kutz S. 2020. Range expansion of muskox lungworms track rapid arctic warming: Implications for geographic colonization under climate forcing. *Sci Rep* 10:17323.
- Kutz S, Bollinger T, Branigan M, Checkley S, Davison T, Dumond M, Elkin B, Forde T, Hutchins W, et al. 2015. Erysipelothrix rhusiopathiae associated with recent widespread muskox mortalities in the Canadian Arctic. Can Vet J 56:560–563.

- Kutz SJ, Hoberg EP, Molnár PK, Dobson A, Verocai GG. 2014. A walk on the tundra: Host–parasite interactions in an extreme environment. Int J Parasitol Parasites Wildl 3:198–208.
- Lafferty KD, Allesina S, Arim M, Briggs CJ, De Leo G, Dobson AP, Dunne JA, Johnson PT, Kuris AM, et al 2008. Parasites in food webs: The ultimate missing links. *Ecol Lett* 11:533–546.
- Mallory ML, Gaston AJ, Gilchrist HG, Robertson GJ, Braune BM. 2010. Effects of climate change, altered sea-ice distribution and seasonal phenology on marine birds. In: A little less Arctic, Ferguson SH, Loseto LL, Mallory ML, editors. Springer, Dordrecht the Netherlands, pp. 179–195.
- Malmberg JL, White LA, VandeWoude S. 2021. Bioaccumulation of pathogen exposure in top predators. *Trends Ecol Evol* 36:411–420.
- Massie GN, Ware MW, Villegas EN, Black MW. 2010. Uptake and transmission of *Toxoplasma gondii* oocysts by migratory, filter-feeding fish. *Vet Parasitol* 169:296–303.
- Mavrot F, Aleuy OA, Forde T, Kutz SJ. 2022. Erysipelas in Arctic and northern regions. In: Arctic one health: Challenges for northern animals and people, Tryland M, editor. Springer, Cham, Switzerland, pp. 363–375.
- Mavrot F, Orsel K, Hutchins W, Adams LG, Beckmen K, Blake JE, Checkley SL, Davison T, Di Francesco J, et al. 2020. Novel insights into serodiagnosis and epidemiology of *Erysipelothrix rhusiopathiae*, a newly recognized pathogen in muskoxen (Ovibos moschatus). PLoS One 15:e0231724.
- Melles SJ, Fortin MJ, Lindsay K, Badzinski D. 2011. Expanding northward: Influence of climate change, forest connectivity, and population processes on a threatened species' range shift. *Glob Change Biol* 17:17–31.
- Molnar PK, Derocher AE, Thiemann GW, Lewis MA. 2010. Predicting survival, reproduction and abundance of polar bears under climate change. *Biol Conserv* 143:1612–1622.
- Moore SM, Borer ET, Hosseini PR. 2010. Predators indirectly control vector-borne disease: Linking predator– prey and host–pathogen models. J R Soc Interface 7:161–176.
- Nguyen L, Pilfold NW, Derocher AE, Stirling I, Bohart AM, Richardson E. 2017. Ringed seal (*Pusa hispida*) tooth annuli as an index of reproduction in the Beaufort Sea. *Ecol Indic* 77:286–292.
- NOAA (National Oceanic and Atmospheric Administration). 2012. Northern pinnipeds (ice seals & walruses) unusual mortality event (UME) March 2012 Update. 1–3 pp. https://media.fisheries.noaa.gov/dam-migration/ume-fact sheet031912-akr.pdf. Accessed August 2024.
- Opsteegh M, Teunis P, Mensink M, Züchner L, Titilincu A, Langelaar M, van der Giessen J. 2010. Evaluation of ELISA test characteristics and estimation of *Toxoplasma gondii* seroprevalence in Dutch sheep using mixture models. *Prev Vet Med* 96:232–240.
- Paetkau D, Amstrup SC, Born EW, Calvert W, Derocher AE, Garner GW, Messier F, Stirling I, Taylor MK, et al. 1999. Genetic structure of the world's polar bear populations. *Mol Ecol* 8:1571–1584.

- Peacock SJ, Kutz SJ, Hoar BM, Molnár PK. 2022. Behaviour is more important than thermal performance for an Arctic host–parasite system under climate change. *R Soc Open Sci* 9:220060.
- Peterman JE. 1944. Diagnostic chart of swine erysipelas. J Am Vet Med Assoc 105:10–11.
- Pilfold NW, Derocher AE, Stirling I, Richardson E. 2015. Multi-temporal factors influence predation for polar bears in a changing climate. *Oikos* 124:1098–1107.
- Pilfold NW, Derocher AE, Stirling I, Richardson E, Andriashek D. 2012. Age and sex composition of seals killed by polar bears in the Eastern Beaufort Sea. *PLOS One* 7:e41429.
- Pilfold NW, Richardson ES, Ellis J, Jenkins E, Scandrett WB, Hernández-Ortiz A, Buhler K, McGeachy D, Al-Adhami B, et al. 2021. Long-term increases in pathogen seroprevalence in polar bears (*Ursus maritimus*) influenced by climate change. *Glob Change Biol* 27:4481-4497.
- Proshutinsky AY, Johnson MA. 1997. Two circulation regimes of the wind-driven Arctic Ocean. J Geophys Res C Oceans 102:12493–12514.
- R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org. Accessed March 2024.
- Rantanen M, Karpechko AY, Lipponen A, Nordling K, Hyvärinen O, Ruosteenoja K, Vihma T, Laaksonen A. 2022. The Arctic has warmed nearly four times faster than the globe since 1979. *Commun Earth Environ* 3:168.
- Regehr EV, Lunn NJ, Amstrup SC, Stirling I. 2007. Effects of earlier sea ice breakup on survival and population size of polar bears in western Hudson Bay. J Wildl Manag 71:2673–2683.
- Secundino I, López-Macías C, Cervantes-Barragán L, Gil-Cruz C, Ríos-Sarabia N, Pastelin-Palacios R, Villasis-Keever MA, Becker I, Puente JL, et al. 2006. Salmonella porins induce a sustained, lifelong specific bactericidal antibody memory response. *Immunology* 117:59–70.
- Seru LV. 2023. Genomic and virulence profiling of Erysipelothrix rhusiopathiae isolated from widespread muskox mortalities in Arctic Archipelago. Master's Thesis, University of Calgary, Calgary, Canada.
- Simon A, Bigras Poulin M, Rousseau AN, Dubey JP, Ogden NH. 2013. Spatiotemporal dynamics of *Toxoplasma* gondii infection in Canadian lynx (*Lynx canadensis*) in western Québec, Canada. J Wildl Dis 49:39–48.
- Smith TG. 1987. The ringed seal, *Phoca hispida*, of the Canadian western Arctic. *Can Bull Fish Aquat Sci* 216:1–81.
- Spraker TR, White PA. 2017. Shaggy lame fox syndrome in Pribilof Island Arctic foxes (Alopex lagopus pribilofensis), Alaska. Vet Pathol 54:258–268.
- Stern HL, Laidre KL. 2016. Sea-ice indicators of polar bear habitat. Cryosphere 10:2027–2041.
- Stirling I. 2002. Polar bears and seals in the eastern Beaufort Sea and Amundsen Gulf: A synthesis of population trends and ecological relationships over three decades. Arctic 55(Suppl 1):59–76.

- Stirling I, McDonald TL, Richardson ES, Regehr EV, Amstrup SC. 2011. Polar bear population status in the northern Beaufort Sea, Canada, 1971–2006. Ecol Appl 21:859–876.
- Stirling I, Parkinson CL. 2006. Possible effects of climate warming on selected populations of polar bears (Ursus maritimus) in the Canadian Arctic. Arctic 59:261–275.
- Stirling I, Richardson E, Thiemann GW, Derocher AE. 2008a. Unusual predation attempts of polar bears on ringed seals in the southern Beaufort Sea: Possible significance of changing spring ice conditions. Arctic 61:14–22.
- Stirling I, Spencer C, Andriashek D. 1989. Immobilization of polar bears (Ursus maritimus) with Telazol[®] in the Canadian Arctic. J Wildl Dis 25:159–168.
- Stirling I, Thiemann GW, Richardson E. 2008b. Quantitative support for a subjective fatness index for immobilized polar bears. J Wildl Manag 72:568–574.
- Su M, Hui C. 2011. The effect of predation on the prevalence and aggregation of pathogens in prey. *Biosys*tems 105:300–306.
- Sudlovenick E, Simonee J, Jones MEB, L'Hérault V, Hernández-Ortiz A, Jenkins E, Parker S, Mavrot F, Schneider A, et al. 2023. Surveillance for zoonotic pathogens and Inuit Qaujimajatuqangit of ringed seals (nattiit) (*Pusa hispida*) in Frobisher Bay and Eclipse Sound, Nunavut, Canada. Arctic 76:293–310.
- Thiemann GW, Iverson SJ, Stirling I. 2008. Polar bear diets and arctic marine food webs: Insights from fatty acid analysis. *Ecol Monogr* 78:591–613.

- Thompson DW, Wallace JM. 1998. The Arctic Oscillation signature in the wintertime geopotential height and temperature fields. *Geophys Res Lett* 25:1297–1300.
- Trono KG, Pérez-Filgueira DM, Duffy S, Borca MV, Carrillo C. 2001. Seroprevalence of bovine leukemia virus in dairy cattle in Argentina: Comparison of sensitivity and specificity of different detection methods. *Vet Microbiol* 83:235–248.
- Tschritter CM, de Groot PVC, Branigan M, Dyck M, Sun Z, Lougheed SC. 2023. A new multiplexed magnetic capture–droplet digital PCR tool for monitoring wild-life population health and pathogen surveillance. *Ecol Evol* 13:e10655.
- VanderWaal KL, Ezenwa VO. 2016. Heterogeneity in pathogen transmission mechanisms and methodology. *Funct Ecol* 30:1606–1622.
- Voorhees H, Sparks R, Huntington HP, Rode KD. 2014. Traditional knowledge about polar bears (Ursus maritimus) in Northwestern Alaska. Arctic 67:523–536.
- Wang Q, Chang BJ, Riley TV. 2010. Erysipelothrix rhusiopathiae. Vet Microbiol 140:405–417.
- Wood RL, Steele JH. 1994. Erysipelothrix infections. In: Handbook of zoonoses, 2nd Ed., Section A, Beran GW, Editor-in-Chief. CRC Press, Boca Raton, Florida, pp. 83–91.
- Woolhouse ME, Taylor LH, Haydon DT. 2001. Population biology of multihost pathogens. Science 292:1109–1112.

Submitted for publication 8 May 2024. Accepted 16 August 2024.