

10.1071/CP19509_AC

© CSIRO 2020

Supplementary Material: *Crop & Pasture Science*, 2020, **71**, 491–505.

Interactions between crop sequences, weed populations and herbicide use in Western Australian broadacre farms: findings of a six-year survey

Martin Harries^{A,B}, Ken C. Flower^B, Craig A. Scanlan^C, Michael T. Rose^D and Michael Renton^{E,B}

^ADepartment of Primary Industries and Regional Development (DPIRD), Government of Western Australia, 20 Gregory Street, Geraldton, WA 6530, Australia.

^BUWA School of Agriculture and Environment and UWA Institute of Agriculture, The University of Western Australia, Perth, Stirling Highway, WA 6009, Australia.

^CDepartment of Primary Industries and Regional Development (DPIRD), Government of Western Australia, 75 York Road, Northam, WA 6401, Australia.

^DNSW Department of Primary Industries, Bruxner Highway, Wollongbar, NSW 2477, Australia.

^ESchool of Biological Sciences, The University of Western Australia, Perth, Stirling Highway, WA 6009, Australia.

^FCorresponding author. Email martin.harries@dpird.wa.gov.au

Supplementary tables

Table S1. Number of paddocks monitored for herbicide data by land use and year.

Year	Wheat	Barley	Canola	Lupin	Pasture	Other crop or fallow	Total
2010	81						81
2011	83	8	15	12	12	5	135
2012	64	16	25	4	20	4	133
2013	65	10	25	9	16	6	131
2014	75	10	26	13	8	2	134
Total	368	44	91	38	56	17	614

Table S2. Number of paddocks monitored by year, land use, and region.

Year	Wheat	Canola	Barley	Lupin	Pasture	Other crop or fallow	Total
2010	142						142
2011	99	25	12	23	17	8	184
2012	85	29	23	6	35	6	184
2013	84	33	10	14	35	7	183
2014	101	31	12	15	18	5	182
2015	75	27	6	12	18	4	142
Total	586	145	63	70	123	30	1017
NAR	268	59	5	43	28	13	416
CAR	196	49	28	18	42	13	346
SAR	122	37	30	9	53	4	255
Total	586	145	63	70	123	30	1017

Supplementary figures




Figure S1. Cumulative frequency distributions of application rate (g/ha.year) for the four most commonly applied herbicide - glyphosate, trifluralin, MCPA and paraquat.




Figure S2. Frequency of applications of the four most commonly used herbicides in each month.