

SUPPLEMENTAL MATERIALS

PARTIAL INCUBATION AND HATCHING ASYNCHRONY IN THE  
RED-SHOULDERED HAWK (*BUTEO LINEATUS*)

SARA J. MILLER

*Arkansas State University, PO Box 599, State University, AR 72467 USA*


CHERYL R. DYKSTRA

*Raptor Environmental, 7280 Susan Springs Drive, West Chester, OH 45069 USA*

JEFFREY L. HAYS AND MELINDA M. SIMON

*RAPTOR, Inc., 961 Barg Salt Run Road, Milford, OH 45150 USA*

Supplemental Figure S1. Incubation behavior (24-hr) during egg-laying at 10 Red-shouldered Hawk nests with 3-egg clutches in Cincinnati, Ohio, 2011 and 2012. Two nests lacked data for the first two days and nights. The day the first egg was laid was designated as Day 1; on average, the second egg was laid on Day 4, and the third on Day 7.



Supplemental Figure S2. Incubation behavior (24-hr) during egg-laying at four Red-shouldered Hawk nests with 4-egg clutches in Cincinnati, Ohio, 2011 and 2012. One nest lacked data for both the day and night of Day 6; another nest lacked data for the first two days and nights. The day the first egg was laid was designated as Day 1; on average, the second egg was laid on Day 4, the third on Day 7, and the fourth on Day 10.

